



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 06:16 PM EDT

PDB ID : 3UBP / pdb_00003ubp
Title : DIAMIDOPHOSPHATE INHIBITED BACILLUS PASTEURII UREASE
Authors : Benini, S.; Rypniewski, W.R.; Wilson, K.S.; Miletto, S.; Mangani, S.; Ciurli, S.
Deposited on : 1998-12-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	101	85% 13% .
2	B	126	% 78% 15% . .
3	C	570	2% 82% 16% . .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6903 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (UREASE GAMMA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	101	Total	C	N	O	S	8	0	0
			781	494	133	148	6			

- Molecule 2 is a protein called PROTEIN (UREASE BETA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	16	0	0
			951	589	171	190	1			

- Molecule 3 is a protein called PROTEIN (UREASE ALPHA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	19	0	0
			4323	2714	743	843	23			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	variant	UNP P41020
C	28	TRP	-	insertion	UNP P41020
C	29	ILE	GLY	variant	UNP P41020
C	36	THR	TYR	variant	UNP P41020
C	37	THR	TYR	variant	UNP P41020
C	38	TYR	LEU	variant	UNP P41020
C	263	LEU	VAL	variant	UNP P41020
C	420	ILE	MET	variant	UNP P41020

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is DIAMIDOPHOSPHATE (CCD ID: 2PA) (formula: $\text{H}_5\text{N}_2\text{O}_2\text{P}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	N	O	P	0	0
			5	2	2	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	124	Total	O	0	0
			124	124		
6	B	186	Total	O	0	0
			186	186		
6	C	531	Total	O	0	0
			531	531		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (UREASE GAMMA SUBUNIT)

Chain A: 




- Molecule 2: PROTEIN (UREASE BETA SUBUNIT)

Chain B: 



- Molecule 3: PROTEIN (UREASE ALPHA SUBUNIT)

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.53Å 131.53Å 188.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	18.00 – 2.00 18.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (18.00-2.00) 99.8 (18.00-2.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.158 , 0.200 0.161 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6903	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 2PA, ACE, NI, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/790	1.39	6/1065 (0.6%)
2	B	0.80	1/963 (0.1%)	1.51	10/1296 (0.8%)
3	C	0.71	2/4392 (0.0%)	1.47	29/5955 (0.5%)
All	All	0.72	3/6145 (0.0%)	1.47	45/8316 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	4
3	C	0	3
All	All	0	7

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	ASN	C-N	-10.85	1.17	1.33
3	C	1	MET	N-CA	5.61	1.56	1.46
3	C	519	ASN	CG-OD1	5.16	1.33	1.23

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	111	GLU	N-CA-CB	12.89	130.14	110.42
2	B	111	GLU	N-CA-C	-10.96	99.42	113.12
3	C	4	ASN	CA-CB-CG	10.18	122.78	112.60
3	C	327	GLN	OE1-CD-NE2	-9.77	112.83	122.60
3	C	251	ASP	CA-CB-CG	9.34	121.94	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	110	LYS	CA-C-O	9.19	130.54	119.97
3	C	547	LYS	CG-CD-CE	8.78	131.49	111.30
3	C	201	HIS	CA-CB-CG	-8.33	105.47	113.80
2	B	5	ASN	CA-C-N	8.18	133.13	121.02
2	B	5	ASN	C-N-CA	8.18	133.13	121.02
3	C	542	GLU	CA-C-N	-8.04	108.35	122.26
3	C	542	GLU	C-N-CA	-8.04	108.35	122.26
1	A	100	SER	CA-C-O	-7.46	108.12	120.80
3	C	352	HIS	CA-CB-CG	-6.71	107.09	113.80
3	C	327	GLN	CG-CD-NE2	6.64	126.36	116.40
3	C	542	GLU	CA-C-O	-6.54	112.13	120.31
3	C	302	ASN	N-CA-C	6.52	124.22	109.81
3	C	191	PRO	N-CA-CB	6.45	107.12	102.28
3	C	164	PRO	N-CA-CB	6.32	106.96	102.79
3	C	566	ARG	NE-CZ-NH2	-6.03	113.78	119.20
2	B	111	GLU	CA-C-O	6.00	127.04	119.78
1	A	81	GLN	CG-CD-NE2	5.92	125.28	116.40
3	C	5	ARG	NE-CZ-NH2	-5.83	113.95	119.20
3	C	303	PRO	N-CA-CB	5.58	108.87	103.51
3	C	463	ASP	CA-CB-CG	5.57	118.17	112.60
3	C	335	PHE	CA-CB-CG	5.56	119.36	113.80
2	B	22	ASN	CB-CG-ND2	5.53	124.69	116.40
3	C	249	HIS	CA-CB-CG	5.48	119.28	113.80
1	A	64	LEU	CA-C-O	-5.46	114.38	120.66
1	A	100	SER	N-CA-C	5.46	126.28	111.00
3	C	329	ILE	CB-CA-C	-5.42	104.94	111.18
3	C	366	ALA	CA-C-N	5.33	131.73	121.54
3	C	366	ALA	C-N-CA	5.33	131.73	121.54
3	C	566	ARG	CD-NE-CZ	5.33	131.87	124.40
3	C	540	ASN	CA-C-N	5.32	124.83	119.19
3	C	540	ASN	C-N-CA	5.32	124.83	119.19
2	B	105	GLY	CA-C-O	5.24	125.30	120.79
3	C	279	ALA	CA-C-O	5.21	126.07	120.55
1	A	99	ILE	CA-C-N	-5.19	112.35	121.70
1	A	99	ILE	C-N-CA	-5.19	112.35	121.70
3	C	307	PHE	CA-CB-CG	5.19	118.99	113.80
3	C	293	HIS	CA-C-N	5.13	124.74	119.56
3	C	293	HIS	C-N-CA	5.13	124.74	119.56
2	B	13	ARG	NE-CZ-NH2	5.09	123.78	119.20
2	B	39	PRO	N-CA-CB	5.05	108.03	103.33

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	109	ASN	Mainchain
2	B	110	LYS	Peptide
2	B	125	VAL	Mainchain
2	B	8	VAL	Mainchain
3	C	289	ALA	Mainchain
3	C	297	LEU	Mainchain
3	C	553	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	781	0	804	12	1
2	B	951	0	936	14	0
3	C	4323	0	4290	51	0
4	C	2	0	0	0	0
5	C	5	0	4	1	0
6	A	124	0	0	7	0
6	B	186	0	0	8	0
6	C	531	0	0	10	5
All	All	6903	0	6034	75	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:325:LEU:HA	3:C:326:LYS:HE2	1.61	0.82
3:C:536:ASP:HB3	6:C:1392:HOH:O	1.84	0.77
2:B:13:ARG:NE	6:B:280:HOH:O	2.21	0.73
2:B:82:GLU:CD	6:B:270:HOH:O	2.30	0.73
3:C:399:ASP:O	3:C:403:LEU:HG	1.90	0.71
2:B:109:ASN:ND2	6:B:269:HOH:O	2.24	0.71
3:C:396:ASN:O	3:C:398:SER:N	2.24	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:548:VAL:O	3:C:551:GLU:HG3	1.92	0.69
1:A:81:GLN:CD	6:A:133:HOH:O	2.39	0.65
2:B:13:ARG:NH1	6:B:200:HOH:O	2.30	0.65
3:C:512:ARG:HD2	6:C:1256:HOH:O	1.99	0.62
3:C:324:HIS:O	3:C:326:LYS:NZ	2.33	0.60
2:B:82:GLU:HG3	6:B:270:HOH:O	2.02	0.59
3:C:513:ARG:NH1	6:C:1220:HOH:O	2.31	0.59
1:A:61:LYS:HD3	6:A:217:HOH:O	2.03	0.59
1:A:81:GLN:HG2	6:A:133:HOH:O	2.01	0.58
3:C:273:SER:HB2	3:C:290:MET:HE3	1.84	0.58
2:B:82:GLU:CG	6:B:270:HOH:O	2.53	0.57
3:C:507:LYS:HD2	6:C:1333:HOH:O	2.06	0.56
1:A:12:GLN:HG3	6:A:223:HOH:O	2.07	0.55
3:C:287:ILE:HD12	3:C:290:MET:HG3	1.88	0.54
2:B:95:GLU:O	3:C:104:ASP:HB3	2.09	0.53
3:C:327:GLN:NE2	6:C:1404:HOH:O	2.11	0.53
3:C:325:LEU:CA	3:C:326:LYS:HE2	2.36	0.52
3:C:249:HIS:CE1	3:C:281:GLY:HA3	2.44	0.52
3:C:329:ILE:HB	3:C:332:ASP:HB2	1.92	0.52
3:C:81:GLY:HA2	3:C:404:LYS:HE2	1.93	0.51
2:B:39:PRO:HG2	6:B:176:HOH:O	2.10	0.51
1:A:81:GLN:NE2	6:A:180:HOH:O	2.44	0.50
2:B:53:LYS:HE2	2:B:122:TYR:CE1	2.46	0.50
3:C:396:ASN:C	3:C:398:SER:H	2.19	0.50
3:C:274:PHE:O	3:C:275:HIS:C	2.55	0.49
1:A:81:GLN:CG	6:A:133:HOH:O	2.58	0.49
1:A:32:TYR:HB3	1:A:33:PRO:HD3	1.94	0.49
3:C:175:PRO:HB3	3:C:199:LYS:HE3	1.95	0.49
3:C:6:GLN:NE2	6:C:1132:HOH:O	2.46	0.48
1:A:96:HIS:O	1:A:97:ASN:C	2.56	0.48
3:C:26:ASP:OD2	6:C:1141:HOH:O	2.20	0.48
3:C:470:THR:N	3:C:471:PRO:CD	2.76	0.47
3:C:472:GLN:HB3	3:C:473:PRO:HA	1.97	0.47
3:C:201:HIS:CE1	3:C:226:GLY:O	2.68	0.47
3:C:378:TRP:HB2	3:C:563:MET:HE1	1.96	0.47
2:B:57:PHE:O	2:B:124:GLY:HA3	2.15	0.46
3:C:362:THR:O	3:C:368:GLY:HA3	2.15	0.46
3:C:452:LYS:HG3	6:C:1140:HOH:O	2.15	0.46
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.97	0.46
3:C:282:GLY:O	3:C:283:HIS:C	2.58	0.46
3:C:139:HIS:CE1	5:C:902:2PA:HN42	2.35	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:522:ASN:HB3	6:C:1374:HOH:O	2.17	0.45
3:C:252:THR:HB	3:C:335:PHE:HE1	1.82	0.45
2:B:78:GLU:HB3	2:B:79:PRO:HD2	1.99	0.44
3:C:137:HIS:CE1	3:C:274:PHE:CD2	3.05	0.44
3:C:138:VAL:O	3:C:159:GLY:HA3	2.18	0.44
2:B:103:THR:HG23	2:B:113:ILE:HD11	1.99	0.44
3:C:384:MET:HA	3:C:384:MET:HE2	1.99	0.43
2:B:126:GLU:OE2	6:B:225:HOH:O	2.21	0.43
3:C:1:MET:HB3	3:C:1:MET:HE2	1.74	0.43
2:B:71:SER:OG	3:C:49:VAL:HG21	2.19	0.43
3:C:132:GLY:HA3	3:C:155:THR:OG1	2.19	0.43
3:C:259:LEU:HD11	3:C:290:MET:HG2	2.01	0.43
3:C:513:ARG:HE	3:C:513:ARG:HB3	1.68	0.43
3:C:301:THR:CG2	3:C:363:ASP:HB2	2.48	0.43
3:C:110:THR:HA	3:C:111:PRO:HD3	1.90	0.43
3:C:158:PHE:CZ	3:C:195:GLY:HA3	2.54	0.43
3:C:43:ASN:OD1	3:C:48:LYS:HB3	2.19	0.42
3:C:461:ILE:CG2	3:C:477:ARG:HD2	2.49	0.42
1:A:12:GLN:NE2	6:A:179:HOH:O	2.50	0.42
3:C:220:KCX:CX	3:C:222:HIS:HD2	2.32	0.42
3:C:437:LEU:HD12	3:C:437:LEU:N	2.34	0.42
1:A:73:VAL:N	1:A:74:PRO:CD	2.83	0.41
3:C:158:PHE:CE1	3:C:195:GLY:HA3	2.55	0.41
3:C:76:ILE:HG23	3:C:427:ILE:HD12	2.01	0.41
3:C:326:LYS:NZ	6:C:1427:HOH:O	2.35	0.41
1:A:32:TYR:HB3	1:A:33:PRO:CD	2.51	0.40
1:A:32:TYR:N	1:A:33:PRO:HD2	2.36	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:1228:HOH:O	6:C:1228:HOH:O[11_555]	1.23	0.97
6:C:1367:HOH:O	6:C:1367:HOH:O[10_665]	1.42	0.78
6:C:1379:HOH:O	6:C:1379:HOH:O[12_565]	1.58	0.62
6:C:1278:HOH:O	6:C:1278:HOH:O[10_665]	1.86	0.34
6:C:1433:HOH:O	6:C:1433:HOH:O[2_665]	1.93	0.27
1:A:18:GLU:OE2	1:A:22:ARG:NH1[2_665]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/101 (98%)	97 (98%)	2 (2%)	0	100	100
2	B	120/126 (95%)	114 (95%)	5 (4%)	1 (1%)	16	11
3	C	567/570 (100%)	541 (95%)	22 (4%)	4 (1%)	18	14
All	All	786/797 (99%)	752 (96%)	29 (4%)	5 (1%)	21	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	397	GLY
2	B	99	ILE
3	C	275	HIS
3	C	283	HIS
3	C	367	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	83 (98%)	2 (2%)	43	47
2	B	101/105 (96%)	99 (98%)	2 (2%)	48	54
3	C	460/460 (100%)	443 (96%)	17 (4%)	30	30
All	All	646/650 (99%)	625 (97%)	21 (3%)	33	34

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
1	A	100	SER
2	B	54	GLU
2	B	126	GLU
3	C	1	MET
3	C	42	VAL
3	C	77	LEU
3	C	158	PHE
3	C	249	HIS
3	C	253	LEU
3	C	305	ARG
3	C	326	LYS
3	C	328	ASN
3	C	329	ILE
3	C	337	ASP
3	C	395	LYS
3	C	398	SER
3	C	403	LEU
3	C	452	LYS
3	C	520	CYS
3	C	551	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
3	C	6	GLN
3	C	179	ASN
3	C	267	ASN
3	C	328	ASN
3	C	519	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	KCX	C	220	4,3	10,11,12	3.20	2 (20%)	6,12,14	2.54	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	220	4,3	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	220	KCX	CX-NZ	7.32	1.48	1.35
3	C	220	KCX	OQ1-CX	6.61	1.33	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	CE-NZ-CX	-5.23	113.10	121.98
3	C	220	KCX	CD-CE-NZ	-2.33	105.66	112.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	220	KCX	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	2PA	C	902	4	1,4,4	5.78	1 (100%)	0,6,6	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	902	2PA	P2-O1	5.78	1.56	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	902	2PA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	5:ASN	C	6:TYR	N	1.17

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/101 (99%)	-0.66	0 100 100	5, 12, 20, 26	4 (4%)
2	B	121/126 (96%)	-0.52	1 (0%) 82 82	7, 14, 26, 45	3 (2%)
3	C	569/570 (99%)	-0.58	14 (2%) 58 58	6, 12, 24, 53	9 (1%)
All	All	790/797 (99%)	-0.58	15 (1%) 66 66	5, 12, 25, 53	16 (2%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	327	GLN	5.0
3	C	397	GLY	3.9
3	C	328	ASN	3.6
3	C	329	ILE	3.2
3	C	326	LYS	3.2
3	C	522	ASN	3.0
2	B	13	ARG	2.8
3	C	316	LEU	2.5
3	C	391	LEU	2.5
3	C	396	ASN	2.5
3	C	336	ALA	2.4
3	C	26	ASP	2.3
3	C	330	PRO	2.2
3	C	390	PRO	2.2
3	C	335	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	KCX	C	220	12/13	0.96	0.06	5,10,17,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	2PA	C	902	5/5	0.97	0.08	14,14,18,19	0
4	NI	C	901	1/1	0.99	0.03	16,16,16,16	0
4	NI	C	900	1/1	0.99	0.04	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.