



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:35 AM UTC

PDB ID : 5UBX / pdb_00005ubx
Title : Crystal structure of a mutant mIgG2b Fc heterodimer in complex with Protein A peptide analog Z34C
Authors : Armstrong, A.A.; Zwolak, A.; Gilliland, G.L.
Deposited on : 2016-12-21
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	9% 73% 10% 17%
2	B	236	4% 73% 12% 15%
3	Z	34	9% 97% .
4	C	6	50% 50%
5	D	8	50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-2B chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1447	921	243	276	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	GLY	-	expression tag	UNP P01867
A	219	SER	-	expression tag	UNP P01867
A	307	THR	PRO	engineered mutation	UNP P01867
A	309	LEU	GLN	engineered mutation	UNP P01867
A	370	LYS	VAL	engineered mutation	UNP P01867
A	409	ARG	LYS	engineered mutation	UNP P01867

- Molecule 2 is a protein called Ig gamma-2B chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	201	Total	C	N	O	S	0	0	0
			1521	968	256	290	7			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	212	HIS	-	expression tag	UNP P01867
B	213	HIS	-	expression tag	UNP P01867
B	214	HIS	-	expression tag	UNP P01867
B	215	HIS	-	expression tag	UNP P01867
B	216	HIS	-	expression tag	UNP P01867
B	217	HIS	-	expression tag	UNP P01867
B	218	GLY	-	expression tag	UNP P01867
B	219	SER	-	expression tag	UNP P01867
B	253	ASP	ILE	engineered mutation	UNP P01867
B	405	LEU	PHE	engineered mutation	UNP P01867

- Molecule 3 is a protein called Peptide analog of B-domain from Protein A - Z34C.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Z	34	Total	C	N	O	S	0	0	0
			269	164	55	47	3			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	6	Total	C	N	O	0	0	0
			75	42	3	30			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	D	8	Total	C	N	O	0	0	0
			99	56	4	39			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	22	Total	O	0	0
			22	22		
6	B	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig gamma-2B chain C region



- Molecule 2: Ig gamma-2B chain C region



- Molecule 3: Peptide analog of B-domain from Protein A - Z34C



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  50% 50%

NA61	NA62	BR43	MAN4	NA65	NA66	NA67	FUC8
------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.49Å 101.49Å 135.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.14 – 2.70 38.14 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.14-2.70) 99.7 (38.14-2.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.69Å)	Xtriage
Refinement program	PHENIX dev_1428	Depositor
R, R_{free}	0.209 , 0.257 0.222 , 0.258	Depositor DCC
R_{free} test set	978 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	74.1	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3453	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1479	0.76	0/2017
2	B	0.37	0/1558	0.76	0/2130
3	Z	0.31	0/273	0.69	0/366
All	All	0.38	0/3310	0.76	0/4513

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1447	0	1362	12	0
2	B	1521	0	1426	14	0
3	Z	269	0	238	1	0
4	C	75	0	64	0	0
5	D	99	0	85	0	0
6	A	22	0	0	0	0
6	B	20	0	0	1	0
All	All	3453	0	3175	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:378:SER:HB3	2:B:428:ARG:HB2	1.75	0.68
2:B:369:VAL:HB	2:B:406:ILE:HG22	1.77	0.65
1:A:293:ARG:HA	1:A:300:ILE:HA	1.80	0.63
1:A:259:VAL:HG23	1:A:308:ILE:HD13	1.83	0.59
1:A:354:PRO:HG2	1:A:357:GLN:HB2	1.87	0.57
2:B:259:VAL:HG23	2:B:308:ILE:HD13	1.89	0.55
2:B:385:GLY:N	6:B:603:HOH:O	2.40	0.54
1:A:246:ASN:O	1:A:250:VAL:HG23	2.08	0.53
2:B:354:PRO:HG2	2:B:357:GLN:HB2	1.91	0.52
2:B:351:LEU:HB2	2:B:366:THR:HB	1.97	0.47
2:B:368:LEU:HD13	2:B:407:TYR:CZ	2.49	0.47
2:B:240:VAL:HG22	2:B:263:VAL:HG22	1.98	0.46
2:B:399:ASP:OD1	2:B:400:SER:N	2.49	0.46
1:A:369:VAL:HB	1:A:406:ILE:HG22	1.98	0.45
1:A:437:LEU:HD12	1:A:438:LYS:H	1.83	0.43
2:B:356:GLU:H	2:B:356:GLU:CD	2.27	0.43
1:A:320:LYS:HA	1:A:335:THR:HA	2.00	0.43
2:B:346:PRO:HB3	2:B:372:PHE:HB3	2.02	0.42
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.54	0.42
1:A:382:THR:OG1	1:A:424:SER:HB2	2.20	0.42
2:B:367:CYS:HB2	2:B:381:TRP:CZ2	2.54	0.42
1:A:434:ASN:HA	3:Z:10:TYR:CG	2.55	0.42
1:A:258:LYS:HE2	1:A:258:LYS:HB2	1.88	0.41
1:A:409:ARG:HD3	2:B:405:LEU:HD13	2.02	0.41
2:B:365:LEU:HD12	2:B:365:LEU:HA	1.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/230 (80%)	182 (98%)	3 (2%)	0	100	100
2	B	197/236 (84%)	196 (100%)	1 (0%)	0	100	100
3	Z	32/34 (94%)	31 (97%)	1 (3%)	0	100	100
All	All	414/500 (83%)	409 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/214 (73%)	155 (99%)	1 (1%)	78	91
2	B	165/220 (75%)	161 (98%)	4 (2%)	43	72
3	Z	25/32 (78%)	25 (100%)	0	100	100
All	All	346/466 (74%)	341 (99%)	5 (1%)	59	82

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	284	VAL
2	B	264	VAL
2	B	303	VAL
2	B	304	SER
2	B	314	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	426	ASN
3	Z	14	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	C	1	4	14,14,15	0.42	0	17,19,21	0.38	0
4	BMA	C	2	4	11,11,12	0.99	1 (9%)	15,15,17	1.26	2 (13%)
4	MAN	C	3	4	11,11,12	0.88	1 (9%)	15,15,17	0.94	2 (13%)
4	NAG	C	4	4	14,14,15	0.31	0	17,19,21	0.40	0
4	MAN	C	5	4	11,11,12	0.80	1 (9%)	15,15,17	1.05	2 (13%)
4	NAG	C	6	4	14,14,15	0.27	0	17,19,21	0.52	0
5	NAG	D	1	5,2	14,14,15	0.44	0	17,19,21	0.64	0
5	NAG	D	2	5	14,14,15	0.55	0	17,19,21	0.49	0
5	BMA	D	3	5	11,11,12	1.05	1 (9%)	15,15,17	1.26	2 (13%)
5	MAN	D	4	5	11,11,12	0.66	0	15,15,17	1.02	2 (13%)
5	NAG	D	5	5	14,14,15	0.29	0	17,19,21	0.49	0
5	MAN	D	6	5	11,11,12	1.00	1 (9%)	15,15,17	0.93	1 (6%)
5	NAG	D	7	5	14,14,15	0.39	0	17,19,21	0.50	0
5	FUC	D	8	5	10,10,11	0.96	1 (10%)	14,14,16	1.01	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	4	-	0/6/23/26	0/1/1/1
4	BMA	C	2	4	-	0/2/19/22	0/1/1/1
4	MAN	C	3	4	-	0/2/19/22	0/1/1/1
4	NAG	C	4	4	-	2/6/23/26	0/1/1/1
4	MAN	C	5	4	-	0/2/19/22	0/1/1/1
4	NAG	C	6	4	-	2/6/23/26	0/1/1/1
5	NAG	D	1	5,2	-	0/6/23/26	0/1/1/1
5	NAG	D	2	5	-	2/6/23/26	0/1/1/1
5	BMA	D	3	5	-	0/2/19/22	0/1/1/1
5	MAN	D	4	5	-	1/2/19/22	0/1/1/1
5	NAG	D	5	5	-	2/6/23/26	0/1/1/1
5	MAN	D	6	5	-	0/2/19/22	0/1/1/1
5	NAG	D	7	5	-	0/6/23/26	0/1/1/1
5	FUC	D	8	5	-	-	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	3	BMA	O5-C1	-3.31	1.38	1.43
5	D	8	FUC	O5-C1	-2.72	1.39	1.43
5	D	6	MAN	C2-C3	2.47	1.56	1.52
4	C	2	BMA	C1-C2	2.44	1.58	1.52
4	C	3	MAN	O5-C1	-2.41	1.39	1.43
4	C	5	MAN	O5-C1	-2.12	1.40	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	3	BMA	C1-O5-C5	3.19	116.47	112.19
5	D	3	BMA	O2-C2-C3	-2.75	104.45	110.15
4	C	2	BMA	C1-O5-C5	2.71	115.82	112.19
4	C	5	MAN	O2-C2-C3	-2.69	104.58	110.15
5	D	4	MAN	C1-O5-C5	2.69	115.79	112.19
4	C	2	BMA	O2-C2-C3	-2.34	105.31	110.15
5	D	4	MAN	O2-C2-C3	-2.28	105.42	110.15
5	D	6	MAN	C1-O5-C5	2.26	115.22	112.19
4	C	5	MAN	C1-O5-C5	2.18	115.11	112.19
4	C	3	MAN	C1-O5-C5	2.16	115.08	112.19
4	C	3	MAN	O2-C2-C3	-2.15	105.69	110.15
5	D	8	FUC	O5-C5-C4	2.04	113.23	109.55

There are no chirality outliers.


All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	5	NAG	O5-C5-C6-O6
5	D	5	NAG	C4-C5-C6-O6
4	C	4	NAG	C4-C5-C6-O6
4	C	4	NAG	O5-C5-C6-O6
4	C	6	NAG	C4-C5-C6-O6
5	D	2	NAG	C4-C5-C6-O6
4	C	6	NAG	O5-C5-C6-O6
5	D	2	NAG	O5-C5-C6-O6
5	D	4	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/230 (83%)	0.43	21 (10%) 10 9	36, 65, 127, 151	0
2	B	201/236 (85%)	0.29	9 (4%) 38 34	45, 69, 107, 138	0
3	Z	34/34 (100%)	0.80	3 (8%) 15 13	66, 95, 127, 148	0
All	All	426/500 (85%)	0.39	33 (7%) 19 17	36, 70, 123, 151	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	ASN	5.6
2	B	444	SER	3.6
1	A	294	GLU	3.5
1	A	332	ILE	3.4
1	A	331	PRO	3.3
3	Z	1	PHE	3.1
2	B	324	ASN	3.1
3	Z	33	ASP	3.1
1	A	333	GLU	3.1
3	Z	34	CYS	3.1
1	A	442	SER	3.0
1	A	269	ASP	3.0
1	A	300	ILE	3.0
1	A	265	ASP	3.0
1	A	270	ASP	3.0
1	A	263	VAL	2.9
1	A	266	VAL	2.7
1	A	241	PHE	2.6
2	B	269	ASP	2.6
1	A	243	PHE	2.6
1	A	264	VAL	2.5
1	A	291	THR	2.5
1	A	288	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	271	PRO	2.4
1	A	273	VAL	2.4
2	B	270	ASP	2.4
2	B	331	PRO	2.3
2	B	323	VAL	2.3
1	A	302	VAL	2.3
1	A	359	SER	2.2
2	B	384	ASN	2.2
2	B	318	GLU	2.0
2	B	284	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	1	14/15	-	-	126,128,130,133	0
4	BMA	C	2	11/12	-	-	120,127,133,134	0
4	MAN	C	3	11/12	-	-	134,138,140,140	0
4	NAG	C	4	14/15	-	-	138,142,146,146	0
4	MAN	C	5	11/12	-	-	129,135,139,143	0
4	NAG	C	6	14/15	-	-	127,131,135,135	0
5	NAG	D	1	14/15	0.52	0.19	67,73,76,78	0
5	NAG	D	2	14/15	-	-	40,69,78,78	0
5	BMA	D	3	11/12	-	-	66,77,88,98	0
5	MAN	D	4	11/12	-	-	103,106,108,112	0
5	NAG	D	5	14/15	-	-	114,118,125,128	0
5	MAN	D	6	11/12	-	-	57,67,76,83	0
5	NAG	D	7	14/15	-	-	69,78,86,89	0
5	FUC	D	8	10/11	-	-	71,74,76,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.