



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:57 AM UTC

PDB ID : 7UBS / pdb_00007ubs
Title : Pfs230 D1 domain in complex with 230AS-26
Authors : Tang, W.K.; Tolia, N.H.
Deposited on : 2022-03-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	11% 83% 7% 10%
1	B	191	10% 84% 7% 9%
1	C	191	8% 73% • 24%
1	D	191	5% 73% • 23%
2	H	264	% 77% 10% 12%
2	I	264	2% 81% 7% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	J	264	2% 81% 8% 12%
2	K	264	% 83% 5% 12%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24542 atoms, of which 12159 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gametocyte surface protein P230.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	172	Total	C	H	N	O	S	0	0	0
			2745	877	1375	213	276	4			
1	B	173	Total	C	H	N	O	S	0	0	0
			2764	883	1386	214	277	4			
1	C	145	Total	C	H	N	O	S	0	0	0
			2371	756	1205	184	222	4			
1	D	147	Total	C	H	N	O	S	0	0	0
			2399	764	1218	187	226	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	585	GLN	ASN	conflict	UNP P68874
B	585	GLN	ASN	conflict	UNP P68874
C	585	GLN	ASN	conflict	UNP P68874
D	585	GLN	ASN	conflict	UNP P68874

- Molecule 2 is a protein called 230AS-26.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	232	Total	C	H	N	O	S	0	0	0
			3532	1135	1734	299	356	8			
2	I	233	Total	C	H	N	O	S	0	0	0
			3554	1141	1747	301	357	8			
2	J	233	Total	C	H	N	O	S	0	0	0
			3554	1141	1747	301	357	8			
2	K	233	Total	C	H	N	O	S	0	0	0
			3554	1141	1747	301	357	8			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total 4	O 4	0	0
3	B	3	Total 3	O 3	0	0
3	C	3	Total 3	O 3	0	0
3	H	12	Total 12	O 12	0	0
3	I	13	Total 13	O 13	0	0
3	J	19	Total 19	O 19	0	0
3	K	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gametocyte surface protein P230




- Molecule 1: Gametocyte surface protein P230



- Molecule 1: Gametocyte surface protein P230





- Molecule 1: Gametocyte surface protein P230











• Molecule 2: 230AS-26





• Molecule 2: 230AS-26



• Molecule 2: 230AS-26



• Molecule 2: 230AS-26



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.88Å 205.51Å 93.86Å 90.00° 103.01° 90.00°	Depositor
Resolution (Å)	19.98 – 2.50 19.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.5 (19.98-2.50) 93.3 (19.98-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.64 (at 2.50Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.209 , 0.250 (Not available) , 0.262	Depositor DCC
R_{free} test set	2782 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	24542	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.21	0/1396	0.47	0/1889
1	B	0.25	0/1404	0.47	0/1900
1	C	0.18	0/1189	0.40	0/1606
1	D	0.19	0/1204	0.38	0/1627
2	H	0.21	0/1840	0.40	1/2502 (0.0%)
2	I	0.20	0/1849	0.38	0/2513
2	J	0.25	0/1849	0.43	0/2513
2	K	0.22	0/1849	0.42	0/2513
All	All	0.22	0/12580	0.42	1/17063 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	250	ARG	CG-CD-NE	-5.03	100.94	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1370	1375	1373	8	0
1	B	1378	1386	1384	11	0
1	C	1166	1205	1205	5	0
1	D	1181	1218	1218	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1798	1734	1736	19	0
2	I	1807	1747	1749	14	0
2	J	1807	1747	1749	11	0
2	K	1807	1747	1749	8	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	H	12	0	0	0	0
3	I	13	0	0	0	0
3	J	19	0	0	0	0
3	K	15	0	0	0	0
All	All	12383	12159	12163	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:148:SER:OG	2:J:149:PRO:HD3	1.71	0.91
1:A:578:SER:OG	1:C:611:CYS:SG	2.34	0.86
2:I:148:SER:OG	2:I:149:PRO:HD3	1.83	0.78
1:D:615:VAL:HG13	1:D:620:ILE:HD12	1.66	0.77
2:H:148:SER:OG	2:H:149:PRO:HD3	1.86	0.75
2:I:232:VAL:HG22	2:I:250:ARG:HG2	1.72	0.71
2:J:232:VAL:HG22	2:J:250:ARG:HG2	1.73	0.70
2:H:180:LEU:HD22	2:H:218:PHE:CD2	2.28	0.68
1:A:567:THR:HG1	1:B:560:ILE:N	1.95	0.65
1:C:615:VAL:HG11	1:C:692:LEU:HD12	1.79	0.65
1:D:615:VAL:HG11	1:D:692:LEU:CD1	2.28	0.63
1:B:615:VAL:HG11	1:B:692:LEU:CD1	2.30	0.62
2:J:145:MET:HE3	2:J:164:CYS:SG	2.41	0.61
2:K:148:SER:HB3	2:K:149:PRO:HD3	1.82	0.60
1:D:615:VAL:HG11	1:D:692:LEU:HD12	1.82	0.60
1:C:615:VAL:HG11	1:C:692:LEU:CD1	2.33	0.58
1:B:615:VAL:HG11	1:B:692:LEU:HD13	1.87	0.57
2:I:148:SER:OG	2:I:163:ASN:HB2	2.04	0.56
2:I:227:ALA:HA	2:I:253:ILE:HD11	1.87	0.56
2:H:180:LEU:HD22	2:H:218:PHE:CG	2.40	0.56
2:H:158:GLU:HG3	2:H:159:ARG:N	2.20	0.55
2:I:180:LEU:HD12	2:I:236:GLN:O	2.06	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:158:GLU:HG3	2:H:159:ARG:H	1.71	0.55
2:I:180:LEU:HD22	2:I:218:PHE:CG	2.41	0.55
2:H:195:ILE:HD13	2:H:201:ARG:HA	1.88	0.54
2:K:232:VAL:HG22	2:K:250:ARG:HG2	1.90	0.54
1:D:648:TYR:CE2	1:D:662:LEU:HD22	2.42	0.53
2:K:184:GLN:HB2	2:K:194:LEU:HD11	1.90	0.53
2:H:158:GLU:CG	2:H:159:ARG:H	2.21	0.53
2:J:154:VAL:HG21	2:J:160:ALA:HB2	1.92	0.51
1:B:632:VAL:HG11	1:B:635:LEU:HD12	1.95	0.49
2:I:180:LEU:HD22	2:I:218:PHE:CD1	2.49	0.48
2:J:159:ARG:HA	2:J:222:ILE:O	2.13	0.48
2:H:232:VAL:HG22	2:H:250:ARG:HG2	1.95	0.48
2:H:145:MET:HE3	2:H:164:CYS:SG	2.53	0.47
2:H:158:GLU:CG	2:H:159:ARG:N	2.77	0.47
2:J:148:SER:OG	2:J:163:ASN:HB3	2.15	0.47
2:H:11:VAL:HG12	2:H:118:THR:HB	1.97	0.46
1:A:563:SER:HB2	1:B:563:SER:HB2	1.97	0.46
2:H:148:SER:OG	2:H:163:ASN:HB2	2.15	0.46
2:I:158:GLU:HG3	2:I:159:ARG:N	2.30	0.46
1:C:648:TYR:CE2	1:C:662:LEU:HD22	2.50	0.46
1:B:603:GLU:N	1:B:716:LYS:HG2	2.31	0.46
2:I:227:ALA:HA	2:I:253:ILE:CD1	2.45	0.45
2:J:159:ARG:HD3	2:J:221:THR:HG23	1.96	0.45
2:H:236:GLN:HE21	2:H:243:ILE:HG23	1.82	0.45
1:B:603:GLU:HG3	1:B:716:LYS:HZ3	1.81	0.45
2:I:154:VAL:O	2:I:254:LYS:N	2.49	0.45
2:K:208:ARG:NH2	2:K:228:GLU:OE2	2.50	0.45
1:D:618:PRO:HB3	1:D:694:PRO:HA	1.99	0.44
2:J:147:GLN:O	2:J:247:GLN:NE2	2.46	0.44
1:B:654:GLU:OE2	1:B:661:LYS:HE3	2.17	0.44
2:J:236:GLN:HE21	2:J:243:ILE:HG23	1.82	0.44
1:A:615:VAL:HG11	1:A:692:LEU:HD13	1.99	0.44
1:A:699:ALA:HB2	1:A:729:PRO:HD3	1.98	0.44
2:H:36:TRP:CE2	2:H:81:MET:HB2	2.53	0.44
2:I:184:GLN:HB2	2:I:194:LEU:HD11	1.99	0.44
1:C:615:VAL:HG13	1:C:620:ILE:HG13	1.99	0.44
2:K:49:GLY:CA	2:K:70:MET:HE1	2.48	0.44
1:A:569:SER:HA	1:A:720:ARG:HH12	1.82	0.43
1:B:615:VAL:HG11	1:B:692:LEU:HD12	2.00	0.43
2:I:230:VAL:CG2	2:I:253:ILE:HG13	2.48	0.43
1:B:715:ASN:OD1	2:H:212:SER:HB3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:165:LYS:HD2	2:H:216:THR:O	2.19	0.42
2:H:223:SER:O	2:H:224:SER:C	2.62	0.42
2:H:39:GLN:HB2	2:H:45:LEU:HD23	2.02	0.42
2:J:208:ARG:NH1	2:J:229:ASP:OD2	2.52	0.42
2:K:49:GLY:C	2:K:70:MET:HE1	2.45	0.41
2:J:39:GLN:HB2	2:J:45:LEU:HD23	2.02	0.41
1:A:568:GLU:OE1	1:A:571:ASP:HB3	2.21	0.41
1:B:567:THR:O	1:B:568:GLU:HB2	2.21	0.41
2:K:145:MET:HE3	2:K:164:CYS:SG	2.60	0.41
1:A:655:GLU:HG3	1:A:657:LYS:HE2	2.01	0.41
2:I:154:VAL:HG12	2:I:155:SER:N	2.36	0.41
2:H:184:GLN:HB2	2:H:194:LEU:HD11	2.02	0.40
2:K:149:PRO:O	2:K:249:THR:HG23	2.22	0.40
2:I:149:PRO:O	2:I:249:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/177 (90%)	160 (100%)	0	100	100
1	B	161/177 (91%)	161 (100%)	0	100	100
1	C	137/177 (77%)	137 (100%)	0	100	100
1	D	139/177 (78%)	139 (100%)	0	100	100
2	H	196/210 (93%)	196 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	I	197/210 (94%)	197 (100%)	0	100	100
2	J	197/210 (94%)	197 (100%)	0	100	100
2	K	197/210 (94%)	197 (100%)	0	100	100
All	All	1384/1548 (89%)	1384 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	598	GLN
2	H	184	GLN
2	J	39	GLN
2	J	57	ASN
2	J	185	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/191 (90%)	0.68	21 (12%) 8 7	20, 37, 72, 92	0
1	B	173/191 (90%)	0.56	19 (10%) 10 8	12, 33, 73, 92	0
1	C	145/191 (75%)	0.61	16 (11%) 10 8	14, 36, 78, 103	0
1	D	147/191 (76%)	0.40	10 (6%) 23 20	14, 31, 71, 83	0
2	H	232/264 (87%)	0.13	3 (1%) 75 71	13, 29, 54, 73	0
2	I	233/264 (88%)	0.09	4 (1%) 69 65	12, 24, 54, 75	0
2	J	233/264 (88%)	-0.09	5 (2%) 63 59	8, 20, 44, 62	0
2	K	233/264 (88%)	-0.12	3 (1%) 75 71	8, 19, 40, 61	0
All	All	1568/1820 (86%)	0.23	81 (5%) 33 29	8, 27, 64, 103	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	585	GLN	5.9
1	A	585	GLN	5.8
2	J	148	SER	5.6
1	B	570	GLY	5.0
1	A	582	TYR	5.0
1	C	656	THR	4.6
2	I	148	SER	4.4
1	C	714	ASP	4.4
1	B	582	TYR	4.4
1	A	570	GLY	4.3
1	B	571	ASP	4.1
1	A	586	ASN	4.1
2	H	148	SER	4.1
1	B	586	ASN	3.9
1	C	713	ASP	3.8
1	B	573	ALA	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	654	GLU	3.7
1	A	715	ASN	3.5
1	B	572	THR	3.5
1	B	584	SER	3.5
2	K	66	GLY	3.4
1	A	563	SER	3.4
1	C	732	ASN	3.3
1	B	583	ALA	3.3
1	B	677	ASN	3.3
2	I	253	ILE	3.3
2	K	148	SER	3.2
1	D	655	GLU	3.1
1	D	656	THR	3.0
1	C	654	GLU	3.0
1	A	657	LYS	2.9
1	C	715	ASN	2.9
1	A	567	THR	2.9
1	A	571	ASP	2.9
1	C	603	GLU	2.8
2	J	250	ARG	2.8
1	B	563	SER	2.8
1	C	653	LYS	2.8
1	A	569	SER	2.8
1	B	561	ASP	2.8
1	C	657	LYS	2.7
2	H	250	ARG	2.7
2	J	65	GLN	2.7
1	A	561	ASP	2.7
1	D	657	LYS	2.7
1	D	653	LYS	2.7
2	I	87	ARG	2.6
1	C	634	LYS	2.6
1	A	572	THR	2.6
1	B	653	LYS	2.6
1	A	573	ALA	2.6
2	I	64	LEU	2.5
1	B	569	SER	2.5
1	C	717	LYS	2.5
1	A	587	THR	2.5
2	J	66	GLY	2.5
1	D	586	ASN	2.5
1	B	560	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	600	LYS	2.4
1	A	562	LEU	2.4
1	D	658	LEU	2.4
1	C	658	LEU	2.3
2	J	63	LYS	2.3
1	B	562	LEU	2.3
1	C	611	CYS	2.3
1	C	684	LYS	2.3
1	B	715	ASN	2.2
2	H	89	ASP	2.2
1	A	677	ASN	2.2
1	B	568	GLU	2.2
1	A	714	ASP	2.1
1	B	634	LYS	2.1
1	D	600	LYS	2.1
1	A	597	ASP	2.1
1	D	714	ASP	2.1
1	A	581	LYS	2.1
2	K	217	ASP	2.1
1	A	588	ASN	2.0
1	D	602	THR	2.0
1	C	655	GLU	2.0
1	A	638	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.