



wwPDB EM Validation Summary Report ⓘ

Mar 14, 2026 – 01:26 AM UTC

PDB ID : 6UC4 / pdb_00006uc4
EMDB ID : EMD-20726
Title : Barbed end side of a cofilactin cluster
Authors : Huehn, A.R.; Bibeau, J.P.; Schramm, A.C.; Cao, W.; De La Cruz, E.M.; Sindelar, C.V.
Deposited on : 2019-09-13
Resolution : 9.20 Å (reported)
Based on initial models : 5YU8, 6DJO

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 9.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	239 (8.70 - 9.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	21% (red), 73% (green), 25% (yellow), 6% (grey)
1	B	377	45% (red), 73% (green), 24% (yellow), 6% (grey)
1	C	377	34% (red), 74% (green), 24% (yellow), 6% (grey)
1	D	377	43% (red), 79% (green), 15% (yellow), 6% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	377	
1	F	377	
1	G	377	
1	H	377	
1	J	377	
1	K	377	
1	L	377	
2	I	166	
2	M	166	
2	N	166	
2	O	166	
2	P	166	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 32038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	370	Total 2887	C 1827	N 488	O 551	S 21	0	0
1	B	370	Total 2887	C 1827	N 488	O 551	S 21	0	0
1	C	370	Total 2887	C 1827	N 488	O 551	S 21	0	0
1	D	356	Total 2785	C 1767	N 466	O 533	S 19	0	0
1	E	190	Total 1481	C 931	N 248	O 290	S 12	0	0
1	F	190	Total 1481	C 931	N 248	O 290	S 12	0	0
1	G	166	Total 1304	C 836	N 218	O 243	S 7	0	0
1	H	166	Total 1304	C 836	N 218	O 243	S 7	0	0
1	J	356	Total 2785	C 1767	N 466	O 533	S 19	0	0
1	K	356	Total 2785	C 1767	N 466	O 533	S 19	0	0
1	L	356	Total 2785	C 1767	N 466	O 533	S 19	0	0

- Molecule 2 is a protein called Cofilin-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	I	164	Total 1283	C 818	N 209	O 249	S 7	0	0
2	M	164	Total 1283	C 818	N 209	O 249	S 7	0	0
2	N	164	Total 1283	C 818	N 209	O 249	S 7	0	0
2	O	164	Total 1283	C 818	N 209	O 249	S 7	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	P	164	1283	818	209	249	7	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
3	A	1	Total	Mg	0
			1	1	
3	B	1	Total	Mg	0
			1	1	
3	C	1	Total	Mg	0
			1	1	
3	D	1	Total	Mg	0
			1	1	
3	E	1	Total	Mg	0
			1	1	
3	F	1	Total	Mg	0
			1	1	
3	J	1	Total	Mg	0
			1	1	
3	K	1	Total	Mg	0
			1	1	
3	L	1	Total	Mg	0
			1	1	

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total 27	C 10	N 5	O 10	P 2	0
4	B	1	Total 27	C 10	N 5	O 10	P 2	0
4	C	1	Total 27	C 10	N 5	O 10	P 2	0
4	D	1	Total 27	C 10	N 5	O 10	P 2	0
4	E	1	Total 27	C 10	N 5	O 10	P 2	0
4	F	1	Total 27	C 10	N 5	O 10	P 2	0
4	J	1	Total 27	C 10	N 5	O 10	P 2	0
4	K	1	Total 27	C 10	N 5	O 10	P 2	0
4	L	1	Total 27	C 10	N 5	O 10	P 2	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Actin, alpha skeletal muscle



- Molecule 1: Actin, alpha skeletal muscle




LEU ASP
GLU SER
LYS GLY
LEU ASP
CYS ASP
GLY VAL
THR THR
HIS HIS
ASN ASN
VAL VAL
PRO PRO
LYS LYS
ILE ILE
TYR TYR

LYS
GLU
LYS
LEU
CYS
TYR
VAL
ALA
LEU
ASP
PHE
GLU
ASN
GLY
MET
GLU
GLY
MET
GLU
GLY
PHE
THR
PHE
GLN
PRO
ALA
GLU
SER
PHE
GLY
ILE
MET
VAL
ARG
ASP
ALA

GLY
ILE
HIS
GLU
LYS
THR
THR
TYR
ASN
SER
ILE
MET
LYS
CYS
ASP
ILE
ALA
THR
ASP
ALA
ALA
ALA
SER
LEU
SER
SER
SER
ALA
ALA
ASN
ASN
VAL
MET
GLY
GLY
THR
THR
MET
TYR
PRO
ILE
ILE
ALA
ASP
ARG
MET
GLN
LYS
GLY
ILE
PHE
THR
THR
VAL
LEU
PHE
SER
GLN
PRO
ALA
GLU
SER
PHE
LYS
ILE
ILE
ILE
ALA
ALA
PRO

PRO
GLU
ARG
LYS
Y337
T341
G342
Q353
Q354
K355
W356
D363
E364
A365
V370
H371
R372
K373
C374
PHE

• Molecule 1: Actin, alpha skeletal muscle



MET
CYS
ASP
GLU
ASP
GLU
T5
L8
V9
S14
G15
L16
V17
K18
A19
G23
D24
D25
A26
P27
R28
A29
V30
S33
I34
V35
P38
R39
HIS
GLN
GLY
VAL
MET
VAL
GLY
MET
GLY
LYS
ASP
S52
Y53
V54
G55
D56
E57
A58
Q59
K61
ARG
GLY
I64
L65
T66

L67
K68
Y69
P70
I71
E72
H73
G74
I75
I76
T77
N78
W79
D80
D81
M82
E83
Y91
N92
E93
L94
R95
V96
E99
E100
T103
L104
E107
A108
P109
L110
N111
P112
M115
R116
M123
A135
V139
L140
S141
L142
Y143
A144
S145
G146
THR
THR
GLY
ILE
VAL
LEU


ASP
SER
GLY
ASP
VAL
GLY
THR
HIS
ASN
VAL
PRO
ILE
TYR
GLY
GLY
TYR
ALA
ALA
LEU
LEU
PRO
HIS
SER
ALA
ILE
MET
ARG
GLU
LEU
ASP
SER
TYR
LEU
ALA
GLY
ARG
ASP
LEU
THR
TYR
PRO
ILE
ILE
MET
LYS
LEU
THR
ARG
GLY
CYS
PRO
THR
PHE
VAL
LEU
THR
GLN
ALA
GLU
SER
PHE
ARG
ILE
ILE
VAL
ARG
ASP
ILE
LYS

GLU
LYS
CYS
TYR
VAL
ALA
LEU
ASP
PHE
GLU
ASN
GLY
MET
ASP
ALA
THR
ALA
ALA
SER
SER
TYR
SER
LEU
LEU
PRO
ASP
GLY
GLN
VAL
ILE
THR
LEU
MET
LYS
ILE
LEU
THR
ARG
GLY
CYS
PRO
THR
PHE
VAL
LEU
PHE
GLN
PRO
SER
PHE
ILE
LYS
GLY
MET
GLU
SER
ALA
GLY

ILE
HIS
GLU
THR
TYR
THR
ASN
SER
ILE
MET
LYS
CYS
ASP
ILE
ASP
ILE
ILE
ARG
LYS
VAL
LEU
SER
LEU
TYR
ALA
ALA
ASN
VAL
MET
SER
GLY
THR
THR
MET
TYR
PRO
ILE
ILE
MET
ARG
GLY
CYS
PRO
THR
PHE
VAL
LEU
THR
GLN
ALA
GLU
SER
PHE
ARG
ILE
ILE
VAL
ARG
ASP
ILE
LYS

GLU
ARG
LYS
Y337
T341
G342
G343
S344
I345
Q353
W356
D363
E364
V370
H371
R372
K373
C374
PHE


• Molecule 1: Actin, alpha skeletal muscle





MET
CYS
ASP
GLU
ASP
GLU
T6
T6
A7
L8
V9
M12
K18
A19
G20
F21
A22
G23
D24
D25
A26
P27
R28
A29
V30
F31
P32
S33
I34
V35
G36
R37
P38
R39
HIS
GLN
GLY
VAL
MET
VAL
GLY
MET
GLY
GLN
LYS
ASP
S52
Y53
V54
Q59
S60
K61
ARG
GLY
I64
L65
T66

L67
E72
H73
I76
T77
D80
D81
W86
N92
R95
V96
A97
E100
H101
P102
T103
T106
E107
A108
P109
L110
M111
M115
R116
V129
P130
A131
I136
Q137
A138
V139
A144
S145
G146
R147
T148
T149
G150
I151
V152
L153
D154
D157
M162
N163
P164



I166
Y166
E167
G168
H173
R177
L178
D179
L180
R183
Y188
L189
M190
R196
G197
Y198
S199
F200
V201
T202
T203
A204
E205
R206
V209
K215
L216
C217
Y218
V219
A228
T229
A230
A231
S232
S233
S234
S235
L236
E237
K238
S239
Y240
E241
L242
P243
D244
G245
Q246
G251
W252




• Molecule 1: Actin, alpha skeletal muscle




• Molecule 1: Actin, alpha skeletal muscle





• Molecule 2: Cofilin-1









• Molecule 2: Cofilin-1





• Molecule 2: Cofilin-1



• Molecule 2: Cofilin-1



• Molecule 2: Cofilin-1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	671	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.052	Depositor
Minimum map value	-0.021	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.021	Depositor
Map size (Å)	293.04, 293.04, 293.04	wwPDB
Map dimensions	220, 220, 220	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.332, 1.332, 1.332	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	1/2949 (0.0%)	0.53	3/3996 (0.1%)
1	B	0.38	1/2949 (0.0%)	0.53	3/3996 (0.1%)
1	C	0.38	1/2949 (0.0%)	0.53	3/3996 (0.1%)
1	D	0.15	0/2844	0.28	0/3855
1	E	0.39	0/1506	0.55	2/2035 (0.1%)
1	F	0.40	0/1506	0.55	2/2035 (0.1%)
1	G	0.34	1/1336 (0.1%)	0.44	0/1814
1	H	0.34	1/1336 (0.1%)	0.44	0/1814
1	J	0.15	0/2844	0.28	0/3855
1	K	0.15	0/2844	0.28	0/3855
1	L	0.15	0/2844	0.28	0/3855
2	I	0.27	0/1302	0.43	0/1745
2	M	0.27	0/1302	0.43	0/1745
2	N	0.27	0/1302	0.43	0/1745
2	O	0.27	0/1302	0.43	0/1745
2	P	0.27	0/1302	0.43	0/1745
All	All	0.29	5/32417 (0.0%)	0.43	13/43831 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	HIS	CE1-NE2	7.98	1.40	1.32
1	H	73	HIS	CE1-NE2	7.95	1.40	1.32
1	B	73	HIS	CE1-NE2	7.89	1.40	1.32
1	C	73	HIS	CE1-NE2	7.89	1.40	1.32
1	G	73	HIS	CE1-NE2	7.88	1.40	1.32

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39	ARG	N-CA-C	7.04	119.04	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	39	ARG	N-CA-C	7.01	119.00	111.36
1	B	39	ARG	N-CA-C	7.01	119.00	111.36
1	C	223	PHE	N-CA-C	5.95	117.85	111.36
1	F	223	PHE	N-CA-C	5.94	117.84	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2887	0	2862	103	0
1	B	2887	0	2862	79	0
1	C	2887	0	2862	65	0
1	D	2785	0	2759	65	0
1	E	1481	0	1479	59	0
1	F	1481	0	1481	75	0
1	G	1304	0	1276	137	0
1	H	1304	0	1276	61	0
1	J	2785	0	2759	90	0
1	K	2785	0	2759	57	0
1	L	2785	0	2759	49	0
2	I	1283	0	1328	47	0
2	M	1283	0	1328	27	0
2	N	1283	0	1328	34	0
2	O	1283	0	1328	38	0
2	P	1283	0	1327	94	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	27	0	12	2	0
4	B	27	0	12	1	0
4	C	27	0	12	1	0
4	D	27	0	12	2	0
4	E	27	0	12	4	0
4	F	27	0	12	4	0
4	J	27	0	12	2	0
4	K	27	0	12	2	0
4	L	27	0	12	2	0
All	All	32038	0	31881	742	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 742 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:177:ARG:NH1	1:G:73:HIS:CA	1.73	1.50
1:F:177:ARG:CZ	1:G:73:HIS:CB	1.89	1.48
1:G:95:ARG:HH22	2:P:99:LEU:CD2	1.27	1.46
1:H:95:ARG:NH2	2:O:99:LEU:HD22	1.27	1.46
1:G:95:ARG:NH2	2:P:99:LEU:CD1	1.88	1.37

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	368/377 (98%)	356 (97%)	11 (3%)	1 (0%)	36 72
1	B	368/377 (98%)	356 (97%)	11 (3%)	1 (0%)	36 72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	368/377 (98%)	356 (97%)	11 (3%)	1 (0%)	36	72
1	D	350/377 (93%)	346 (99%)	4 (1%)	0	100	100
1	E	188/377 (50%)	181 (96%)	7 (4%)	0	100	100
1	F	188/377 (50%)	181 (96%)	7 (4%)	0	100	100
1	G	158/377 (42%)	155 (98%)	2 (1%)	1 (1%)	21	59
1	H	158/377 (42%)	155 (98%)	2 (1%)	1 (1%)	21	59
1	J	350/377 (93%)	346 (99%)	4 (1%)	0	100	100
1	K	350/377 (93%)	346 (99%)	4 (1%)	0	100	100
1	L	350/377 (93%)	346 (99%)	4 (1%)	0	100	100
2	I	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
2	M	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
2	N	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
2	O	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
2	P	162/166 (98%)	160 (99%)	2 (1%)	0	100	100
All	All	4006/4977 (80%)	3924 (98%)	77 (2%)	5 (0%)	49	83

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	GLU
1	B	72	GLU
1	C	72	GLU
1	G	72	GLU
1	H	72	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/320 (98%)	311 (99%)	2 (1%)	78	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	313/320 (98%)	311 (99%)	2 (1%)	78	83
1	C	313/320 (98%)	311 (99%)	2 (1%)	78	83
1	D	303/320 (95%)	302 (100%)	1 (0%)	86	86
1	E	163/320 (51%)	162 (99%)	1 (1%)	78	83
1	F	163/320 (51%)	162 (99%)	1 (1%)	78	83
1	G	140/320 (44%)	139 (99%)	1 (1%)	76	81
1	H	140/320 (44%)	139 (99%)	1 (1%)	76	81
1	J	303/320 (95%)	302 (100%)	1 (0%)	86	86
1	K	303/320 (95%)	302 (100%)	1 (0%)	86	86
1	L	303/320 (95%)	302 (100%)	1 (0%)	86	86
2	I	143/144 (99%)	141 (99%)	2 (1%)	59	72
2	M	143/144 (99%)	141 (99%)	2 (1%)	59	72
2	N	143/144 (99%)	141 (99%)	2 (1%)	59	72
2	O	143/144 (99%)	141 (99%)	2 (1%)	59	72
2	P	143/144 (99%)	141 (99%)	2 (1%)	59	72
All	All	3472/4240 (82%)	3448 (99%)	24 (1%)	73	81

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	129	VAL
2	M	153	LEU
2	M	91	THR
2	N	91	THR
1	D	129	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	136	GLN
1	K	246	GLN
1	K	12	ASN
1	K	115	ASN
1	K	297	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	B	802	3	28,29,29	1.41	4 (14%)	43,45,45	1.82	9 (20%)
4	ADP	D	402	3	28,29,29	1.41	4 (14%)	43,45,45	1.82	9 (20%)
4	ADP	K	402	3	28,29,29	1.41	4 (14%)	43,45,45	1.83	9 (20%)
4	ADP	J	402	3	28,29,29	1.42	4 (14%)	43,45,45	1.83	10 (23%)
4	ADP	E	802	3	28,29,29	1.42	4 (14%)	43,45,45	1.83	9 (20%)
4	ADP	A	802	3	28,29,29	1.41	4 (14%)	43,45,45	1.82	8 (18%)
4	ADP	L	402	3	28,29,29	1.41	4 (14%)	43,45,45	1.82	9 (20%)
4	ADP	C	802	3	28,29,29	1.42	4 (14%)	43,45,45	1.82	8 (18%)
4	ADP	F	802	3	28,29,29	1.42	4 (14%)	43,45,45	1.82	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	B	802	3	-	4/16/32/32	0/3/3/3
4	ADP	D	402	3	-	0/16/32/32	0/3/3/3
4	ADP	K	402	3	-	0/16/32/32	0/3/3/3
4	ADP	J	402	3	-	0/16/32/32	0/3/3/3
4	ADP	E	802	3	-	4/16/32/32	0/3/3/3
4	ADP	A	802	3	-	4/16/32/32	0/3/3/3
4	ADP	L	402	3	-	0/16/32/32	0/3/3/3
4	ADP	C	802	3	-	4/16/32/32	0/3/3/3
4	ADP	F	802	3	-	4/16/32/32	0/3/3/3

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	802	ADP	C5-C4	4.80	1.47	1.39
4	J	402	ADP	C5-C4	4.75	1.47	1.39
4	B	802	ADP	C5-C4	4.75	1.47	1.39
4	C	802	ADP	C5-C4	4.74	1.47	1.39
4	K	402	ADP	C5-C4	4.72	1.47	1.39

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	K	402	ADP	C5-C4-N3	-5.88	118.62	126.72
4	E	802	ADP	C5-C4-N3	-5.87	118.64	126.72
4	J	402	ADP	C5-C4-N3	-5.86	118.65	126.72
4	A	802	ADP	C5-C4-N3	-5.85	118.66	126.72
4	F	802	ADP	C5-C4-N3	-5.84	118.67	126.72

There are no chirality outliers.

5 of 20 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	802	ADP	C5'-O5'-PA-O2A
4	A	802	ADP	C5'-O5'-PA-O3A
4	B	802	ADP	C5'-O5'-PA-O2A
4	B	802	ADP	C5'-O5'-PA-O3A
4	C	802	ADP	C5'-O5'-PA-O2A



There are no ring outliers.



9 monomers are involved in 20 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	802	ADP	1	0
4	D	402	ADP	2	0
4	K	402	ADP	2	0
4	J	402	ADP	2	0
4	E	802	ADP	4	0
4	A	802	ADP	2	0
4	L	402	ADP	2	0
4	C	802	ADP	1	0
4	F	802	ADP	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20726. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map




6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 110




Y Index: 110




Z Index: 110


6.2.2 Raw map



X Index: 110



Y Index: 110




Z Index: 110


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 116




Y Index: 103




Z Index: 97


6.3.2 Raw map



X Index: 111



Y Index: 107




Z Index: 108


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.021. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_20726_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 312 nm³; this corresponds to an approximate mass of 281 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.109 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.109 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	9.20	-	-
Author-provided FSC curve	9.07	16.56	9.51
Unmasked-calculated*	17.42	25.84	22.88

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 17.42 differs from the reported value 9.2 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20726 and PDB model 6UC4. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.021 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.021).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 73% of all backbone atoms, 54% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.021) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5430	 0.1150
A	 0.6170	 0.1240
B	 0.4550	 0.1000
C	 0.5240	 0.1150
D	 0.4610	 0.0950
E	 0.6480	 0.1310
F	 0.5960	 0.1240
G	 0.5640	 0.1190
H	 0.6020	 0.1130
I	 0.4230	 0.1000
J	 0.5600	 0.1120
K	 0.6050	 0.1210
L	 0.5950	 0.1200
M	 0.4710	 0.1150
N	 0.5880	 0.1310
O	 0.4990	 0.1190
P	 0.4410	 0.1200

