



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:00 AM UTC

PDB ID : 8UCE / pdb\_00008uce  
Title : Thermophilic RNA Ligase from *Palaeococcus pacificus* + AMP  
Authors : Rousseau, M.D.; Hicks, J.L.; Oulavallickal, T.; Williamson, A.; Arcus, V.L.;  
Patrick, M.W.  
Deposited on : 2023-09-26  
Resolution : 2.12 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	 5% 87% 7% 5%
1	B	402	 6% 87% 7% 6%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP dependent DNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	380	3013	1962	506	534	11	0	0	0
1	B	379	2992	1945	503	534	10	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP A0A075LQ94
A	-19	GLY	-	expression tag	UNP A0A075LQ94
A	-18	SER	-	expression tag	UNP A0A075LQ94
A	-17	SER	-	expression tag	UNP A0A075LQ94
A	-16	HIS	-	expression tag	UNP A0A075LQ94
A	-15	HIS	-	expression tag	UNP A0A075LQ94
A	-14	HIS	-	expression tag	UNP A0A075LQ94
A	-13	HIS	-	expression tag	UNP A0A075LQ94
A	-12	HIS	-	expression tag	UNP A0A075LQ94
A	-11	HIS	-	expression tag	UNP A0A075LQ94
A	-10	SER	-	expression tag	UNP A0A075LQ94
A	-9	SER	-	expression tag	UNP A0A075LQ94
A	-8	GLY	-	expression tag	UNP A0A075LQ94
A	-7	LEU	-	expression tag	UNP A0A075LQ94
A	-6	VAL	-	expression tag	UNP A0A075LQ94
A	-5	PRO	-	expression tag	UNP A0A075LQ94
A	-4	ARG	-	expression tag	UNP A0A075LQ94
A	-3	GLY	-	expression tag	UNP A0A075LQ94
A	-2	SER	-	expression tag	UNP A0A075LQ94
A	-1	HIS	-	expression tag	UNP A0A075LQ94
A	0	MET	-	expression tag	UNP A0A075LQ94
B	-20	MET	-	initiating methionine	UNP A0A075LQ94
B	-19	GLY	-	expression tag	UNP A0A075LQ94
B	-18	SER	-	expression tag	UNP A0A075LQ94
B	-17	SER	-	expression tag	UNP A0A075LQ94

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP A0A075LQ94
B	-15	HIS	-	expression tag	UNP A0A075LQ94
B	-14	HIS	-	expression tag	UNP A0A075LQ94
B	-13	HIS	-	expression tag	UNP A0A075LQ94
B	-12	HIS	-	expression tag	UNP A0A075LQ94
B	-11	HIS	-	expression tag	UNP A0A075LQ94
B	-10	SER	-	expression tag	UNP A0A075LQ94
B	-9	SER	-	expression tag	UNP A0A075LQ94
B	-8	GLY	-	expression tag	UNP A0A075LQ94
B	-7	LEU	-	expression tag	UNP A0A075LQ94
B	-6	VAL	-	expression tag	UNP A0A075LQ94
B	-5	PRO	-	expression tag	UNP A0A075LQ94
B	-4	ARG	-	expression tag	UNP A0A075LQ94
B	-3	GLY	-	expression tag	UNP A0A075LQ94
B	-2	SER	-	expression tag	UNP A0A075LQ94
B	-1	HIS	-	expression tag	UNP A0A075LQ94
B	0	MET	-	expression tag	UNP A0A075LQ94

- Molecule 2 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula:  $C_{10}H_{14}N_5O_7P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		
2	B	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Mg 2 2	0	0
4	B	2	Total Mg 2 2	0	0


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	241	Total 241	O 241	0	0
5	B	235	Total 235	O 235	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP dependent DNA ligase



- Molecule 1: ATP dependent DNA ligase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.22Å 98.80Å 142.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.93 – 2.12 42.93 – 2.12	Depositor EDS
% Data completeness (in resolution range)	92.2 (42.93-2.12) 92.2 (42.93-2.12)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.53 (at 2.12Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.189 , 0.237 0.190 , 0.237	Depositor DCC
$R_{free}$ test set	1838 reflections (2.79%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.2	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6571	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.26 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0174e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/3083	0.48	0/4164
1	B	0.33	0/3060	0.48	0/4134
All	All	0.33	0/6143	0.48	0/8298

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3013	0	2936	22	0
1	B	2992	0	2896	17	0
2	A	22	0	12	0	0
2	B	22	0	12	0	0
3	A	30	0	40	4	0
3	B	12	0	16	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	241	0	0	3	2
5	B	235	0	0	2	1
All	All	6571	0	5912	39	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:MET:HE1	1:A:173:GLU:HG3	1.73	0.70
1:A:342:ILE:HG12	1:A:361:ARG:HG3	1.77	0.67
1:A:312:ASP:OD1	5:A:501:HOH:O	2.12	0.67
1:A:318:PHE:CE1	3:A:402:GOL:H2	2.30	0.65
1:B:342:ILE:HG12	1:B:361:ARG:HG3	1.81	0.63
1:A:129:THR:HG22	1:A:131:ASP:H	1.68	0.58
1:B:310:GLY:O	5:B:501:HOH:O	2.17	0.58
1:A:312:ASP:OD2	1:A:312:ASP:N	2.41	0.54
1:B:265:MET:HE2	1:B:268:ILE:HD12	1.89	0.54
1:A:154:VAL:HG13	3:A:402:GOL:H11	1.92	0.52
1:A:345:ASP:HA	3:A:404:GOL:H31	1.92	0.51
1:B:129:THR:HG22	1:B:131:ASP:H	1.76	0.51
1:B:5:TYR:O	1:B:9:ILE:HG13	2.12	0.49
1:A:139:MET:HE3	5:A:720:HOH:O	2.12	0.49
1:B:138:ASN:HA	1:B:174:LYS:HD2	1.95	0.48
1:A:318:PHE:CD1	3:A:402:GOL:H2	2.49	0.47
1:B:97:ASN:HB3	1:B:112:ARG:HG3	1.96	0.47
1:A:158:PRO:HG3	1:A:221:GLU:O	2.15	0.47
1:A:325:LEU:HD13	1:A:355:TRP:CD2	2.50	0.47
1:B:158:PRO:HG3	1:B:221:GLU:O	2.16	0.46
1:B:343:HIS:HE1	5:B:502:HOH:O	1.97	0.46
1:B:96:TYR:HA	1:B:112:ARG:HH22	1.81	0.46
1:B:298:GLU:HB2	1:B:299:PRO:HD3	1.98	0.45
1:B:31:GLU:OE1	1:B:38:ARG:HD3	2.17	0.44
1:A:332:VAL:O	1:A:336:GLU:HG3	2.19	0.43
1:A:327:THR:O	1:A:331:MET:HG3	2.19	0.43
1:B:222:LYS:HZ2	1:B:222:LYS:HG2	1.78	0.42
1:B:246:ILE:HG13	1:B:271:LEU:HD11	2.00	0.42
1:A:138:ASN:O	1:A:139:MET:HE2	2.19	0.42
1:A:38:ARG:HD3	1:A:38:ARG:HA	1.83	0.42
1:B:348:GLU:HG2	1:B:356:ARG:HB3	2.02	0.41
1:B:254:ARG:HG2	1:B:299:PRO:HB3	2.01	0.41
1:A:32:ASP:HB3	1:A:39:TYR:CZ	2.56	0.41
1:A:307:ILE:HD13	1:A:371:ARG:HA	2.02	0.41
1:A:142:CYS:HB2	1:A:170:ASP:HB3	2.03	0.40
1:B:211:LEU:HA	1:B:211:LEU:HD23	1.86	0.40
1:A:16:ASP:O	1:A:20:ILE:HG13	2.22	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:LYS:HE2	5:A:719:HOH:O	2.20	0.40
1:A:233:MET:HE3	1:A:233:MET:HB3	1.91	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:571:HOH:O	5:A:741:HOH:O[2_566]	1.98	0.22
5:A:688:HOH:O	5:B:729:HOH:O[3_645]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/402 (94%)	372 (98%)	6 (2%)	0	100	100
1	B	375/402 (93%)	369 (98%)	6 (2%)	0	100	100
All	All	753/804 (94%)	741 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/357 (86%)	305 (100%)	1 (0%)	86	91

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	301/357 (84%)	299 (99%)	2 (1%)	76	83
All	All	607/714 (85%)	604 (100%)	3 (0%)	81	87

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	THR
1	B	3	SER
1	B	15	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	HIS
1	A	266	GLN
1	A	329	HIS
1	B	138	ASN
1	B	247	ASN
1	B	343	HIS
1	B	377	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	402	-	5,5,5	1.15	0	5,5,5	1.13	1 (20%)
3	GOL	A	403	-	5,5,5	0.89	0	5,5,5	0.84	0
2	AMP	B	401	4	21,24,25	1.46	3 (14%)	30,35,38	2.06	9 (30%)
3	GOL	A	405	-	5,5,5	0.87	0	5,5,5	1.14	1 (20%)
2	AMP	A	401	4	21,24,25	1.45	5 (23%)	30,35,38	1.92	8 (26%)
3	GOL	B	403	-	5,5,5	1.39	0	5,5,5	0.93	0
3	GOL	A	404	-	5,5,5	1.05	0	5,5,5	1.02	0
3	GOL	A	402	-	5,5,5	0.66	0	5,5,5	1.35	0
3	GOL	A	406	-	5,5,5	0.92	0	5,5,5	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	402	-	-	2/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-
2	AMP	B	401	4	-	5/7/25/26	0/3/3/3
3	GOL	A	405	-	-	2/4/4/4	-
2	AMP	A	401	4	-	5/7/25/26	0/3/3/3
3	GOL	B	403	-	-	0/4/4/4	-
3	GOL	A	404	-	-	2/4/4/4	-
3	GOL	A	402	-	-	3/4/4/4	-
3	GOL	A	406	-	-	0/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	AMP	C5-C4	4.64	1.47	1.39
2	B	401	AMP	C5-C4	4.50	1.47	1.39
2	B	401	AMP	C5-N7	-2.33	1.34	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	AMP	C5-C6	2.27	1.47	1.41
2	A	401	AMP	C5-C6	2.18	1.47	1.41
2	A	401	AMP	C5-N7	-2.18	1.35	1.39
2	A	401	AMP	C8-N7	2.06	1.35	1.31
2	A	401	AMP	C4-N9	-2.04	1.33	1.37

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	AMP	N3-C4-N9	5.32	136.21	127.17
2	B	401	AMP	C5-C4-N3	-5.17	119.60	126.72
2	A	401	AMP	C5-C4-N3	-5.06	119.75	126.72
2	A	401	AMP	N3-C4-N9	4.50	134.83	127.17
2	B	401	AMP	C4-N9-C8	3.98	109.91	105.74
2	A	401	AMP	C2-N3-C4	3.00	119.17	111.83
2	A	401	AMP	C4-N9-C8	3.00	108.89	105.74
2	B	401	AMP	C2-N3-C4	2.97	119.08	111.83
2	A	401	AMP	N3-C2-N1	-2.90	124.19	128.58
2	B	401	AMP	N3-C2-N1	-2.87	124.24	128.58
2	A	401	AMP	C4-C5-N7	-2.37	107.87	110.58
2	B	401	AMP	C1'-N9-C8	-2.34	121.91	127.09
2	B	401	AMP	N6-C6-N1	2.32	123.54	118.38
2	B	401	AMP	N9-C8-N7	-2.29	110.68	113.94
2	B	401	AMP	C5-N7-C8	2.18	106.88	103.45
3	B	402	GOL	C3-C2-C1	-2.16	103.89	111.80
2	A	401	AMP	C2-N1-C6	2.14	122.24	118.73
3	A	405	GOL	C3-C2-C1	-2.05	104.29	111.80
2	A	401	AMP	C5-N7-C8	2.03	106.64	103.45

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	AMP	C4'-C5'-O5'-P
2	B	401	AMP	C4'-C5'-O5'-P
3	A	402	GOL	O1-C1-C2-C3
3	A	402	GOL	C1-C2-C3-O3
3	A	404	GOL	O1-C1-C2-C3
3	A	405	GOL	C1-C2-C3-O3
2	B	401	AMP	C3'-C4'-C5'-O5'
3	A	402	GOL	O2-C2-C3-O3
2	A	401	AMP	C3'-C4'-C5'-O5'

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	403	GOL	C1-C2-C3-O3
3	B	402	GOL	C1-C2-C3-O3
3	A	403	GOL	O2-C2-C3-O3
3	A	404	GOL	O1-C1-C2-O2
3	A	405	GOL	O2-C2-C3-O3
2	A	401	AMP	O4'-C4'-C5'-O5'
2	B	401	AMP	O4'-C4'-C5'-O5'
3	B	402	GOL	O2-C2-C3-O3
2	A	401	AMP	C2'-C1'-N9-C8
2	B	401	AMP	C2'-C1'-N9-C8
2	B	401	AMP	O4'-C1'-N9-C8
2	A	401	AMP	O4'-C1'-N9-C8

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	GOL	1	0
3	A	402	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	380/402 (94%)	-0.03	22 (5%) 29 31	15, 24, 52, 77	0
1	B	379/402 (94%)	-0.01	24 (6%) 26 28	14, 22, 57, 75	0
All	All	759/804 (94%)	-0.02	46 (6%) 27 29	14, 23, 53, 77	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	GLY	5.0
1	A	46	ALA	3.9
1	A	14	GLY	3.9
1	B	11	LEU	3.9
1	A	48	SER	3.7
1	A	311	ASP	3.6
1	B	309	SER	3.5
1	B	48	SER	3.3
1	B	1	MET	3.3
1	B	12	ASN	3.2
1	B	46	ALA	3.0
1	A	1	MET	2.9
1	B	30	VAL	2.8
1	A	11	LEU	2.6
1	B	112	ARG	2.6
1	A	15	LEU	2.6
1	B	25	ASN	2.5
1	B	23	LEU	2.5
1	B	379	PHE	2.5
1	B	19	ARG	2.5
1	A	12	ASN	2.5
1	B	311	ASP	2.5
1	A	310	GLY	2.4
1	A	13	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	13	LEU	2.4
1	A	16	ASP	2.4
1	A	44	ASP	2.4
1	A	337	ARG	2.4
1	A	17	GLU	2.3
1	B	2	VAL	2.3
1	B	20	ILE	2.3
1	A	22	VAL	2.3
1	A	45	SER	2.2
1	B	27	GLY	2.2
1	B	5	TYR	2.2
1	B	44	ASP	2.2
1	A	18	GLU	2.1
1	A	47	ARG	2.1
1	A	338	LEU	2.1
1	A	21	GLU	2.1
1	B	351	ASP	2.1
1	B	340	LEU	2.1
1	A	312	ASP	2.1
1	B	129	THR	2.1
1	A	20	ILE	2.0
1	B	15	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	402	6/6	0.76	0.17	42,60,67,73	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	403	6/6	0.79	0.18	52,54,72,75	0
3	GOL	B	403	6/6	0.81	0.21	37,47,55,58	0
3	GOL	A	402	6/6	0.83	0.22	43,48,51,59	0
3	GOL	A	404	6/6	0.85	0.14	46,53,58,69	0
3	GOL	A	405	6/6	0.86	0.15	52,56,60,66	0
3	GOL	A	406	6/6	0.87	0.13	44,48,56,60	0
4	MG	A	408	1/1	0.95	0.03	24,24,24,24	0
2	AMP	A	401	22/23	0.98	0.05	15,15,17,19	0
2	AMP	B	401	22/23	0.98	0.05	14,14,16,19	0
4	MG	A	407	1/1	0.99	0.01	16,16,16,16	0
4	MG	B	405	1/1	0.99	0.03	21,21,21,21	0
4	MG	B	404	1/1	1.00	0.02	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around MG A 408:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around MG A 407:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MG B 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.