



Full wwPDB EM Validation Report ⓘ

Mar 7, 2026 – 02:21 AM UTC

PDB ID : 8UCR / pdb_00008ucr
EMDB ID : EMD-42136
Title : PhiCb5 maturation protein with Caulobacter crescentus bNY30a pili
Authors : Wang, Y.; Zhang, J.
Deposited on : 2023-09-27
Resolution : 6.45 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY




The reported resolution of this entry is 6.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.















Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	502 (5.95 - 6.95)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	MT	372	
2	A	45	
2	X	45	
2	Z	45	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	a	45	 93%7%
2	b	45	 100%
2	c	45	 93%7%
2	d	45	 98%.
2	e	45	 100%
2	f	45	 100%
2	g	45	 100%
2	h	45	 100%
2	i	45	 100%
2	j	45	 98%.
2	k	45	 100%
2	l	45	 100%
2	m	45	 100%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maturation protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	MT	372	Total	C	N	O	S	0	0
			2866	1805	522	528	11		

- Molecule 2 is a protein called Flp family type IVb pilin.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	A	45	Total	C	N	O	0	0
			308	200	48	60		
2	X	45	Total	C	N	O	0	0
			308	200	48	60		
2	Z	45	Total	C	N	O	0	0
			308	200	48	60		
2	a	45	Total	C	N	O	0	0
			308	200	48	60		
2	b	45	Total	C	N	O	0	0
			308	200	48	60		
2	c	45	Total	C	N	O	0	0
			308	200	48	60		
2	d	45	Total	C	N	O	0	0
			308	200	48	60		
2	e	45	Total	C	N	O	0	0
			308	200	48	60		
2	f	45	Total	C	N	O	0	0
			308	200	48	60		
2	g	45	Total	C	N	O	0	0
			308	200	48	60		
2	h	45	Total	C	N	O	0	0
			308	200	48	60		
2	i	45	Total	C	N	O	0	0
			308	200	48	60		
2	j	45	Total	C	N	O	0	0
			308	200	48	60		
2	k	45	Total	C	N	O	0	0
			308	200	48	60		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf	Trace
2	l	45	Total	C	N	O	0	0
			308	200	48	60		
2	m	45	Total	C	N	O	0	0
			308	200	48	60		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maturation protein



- Molecule 2: Flp family type IVb pilin



There are no outlier residues recorded for this chain.


- Molecule 2: Flp family type IVb pilin



- Molecule 2: Flp family type IVb pilin



- Molecule 2: Flp family type IVb pilin





- Molecule 2: Flp family type IVb pilin

Chain b:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: Flp family type IVb pilin

Chain c:  93% 7%




- Molecule 2: Flp family type IVb pilin

Chain d:  98%



- Molecule 2: Flp family type IVb pilin

Chain e:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: Flp family type IVb pilin

Chain f:  100%



- Molecule 2: Flp family type IVb pilin

Chain g:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: Flp family type IVb pilin

Chain h:  100%





- Molecule 2: Flp family type IVb pilin

Chain i:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: Flp family type IVb pilin

Chain j:  98%




- Molecule 2: Flp family type IVb pilin

Chain k:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: Flp family type IVb pilin

Chain l:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Flp family type IVb pilin

Chain m:  100%

There are no outlier residues recorded for this chain.

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	56330	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	5000	Depositor
Maximum defocus (nm)	25000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.030	Depositor
Minimum map value	-0.009	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.00127	Depositor
Map size ($\text{\AA}$)	516.0, 516.0, 516.0	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.86, 0.86, 0.86	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	MT	1.62	39/2933 (1.3%)	1.82	63/3980 (1.6%)
2	A	0.69	0/309	1.31	0/423
2	X	0.69	0/309	1.29	0/423
2	Z	0.69	0/309	1.32	0/423
2	a	0.99	1/309 (0.3%)	1.75	3/423 (0.7%)
2	b	0.69	0/309	1.29	0/423
2	c	0.68	0/309	1.26	0/423
2	d	0.71	0/309	1.31	0/423
2	e	0.70	0/309	1.29	0/423
2	f	0.69	0/309	1.28	0/423
2	g	0.70	0/309	1.27	0/423
2	h	0.69	0/309	1.30	0/423
2	i	0.69	0/309	1.35	0/423
2	j	0.72	0/309	1.34	0/423
2	k	0.69	0/309	1.29	0/423
2	l	0.69	0/309	1.31	0/423
2	m	0.70	0/309	1.34	0/423
All	All	1.14	40/7877 (0.5%)	1.53	66/10748 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	MT	0	1

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	a	38	LEU	C-N	12.08	1.51	1.33
1	MT	177	HIS	ND1-CE1	-8.74	1.23	1.32
1	MT	369	HIS	ND1-CE1	-8.54	1.24	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	MT	250	ARG	CZ-NH2	-8.03	1.23	1.33
1	MT	124	ARG	CZ-NH2	-7.91	1.23	1.33
1	MT	372	ARG	CZ-NH2	-7.91	1.23	1.33
1	MT	128	ARG	CZ-NH2	-7.90	1.23	1.33
1	MT	178	ARG	CZ-NH2	-7.89	1.23	1.33
1	MT	139	ARG	CZ-NH2	-7.87	1.23	1.33
1	MT	170	ARG	CZ-NH2	-7.87	1.23	1.33
1	MT	151	GLY	N-CA	-6.59	1.38	1.45
1	MT	249	ALA	CA-CB	-6.57	1.43	1.53
1	MT	167	ALA	CA-CB	-6.50	1.43	1.53
1	MT	250	ARG	CZ-NH1	-6.49	1.23	1.32
1	MT	139	ARG	CZ-NH1	-6.45	1.23	1.32
1	MT	278	GLY	N-CA	-6.45	1.37	1.45
1	MT	170	ARG	CZ-NH1	-6.41	1.23	1.32
1	MT	128	ARG	CZ-NH1	-6.39	1.23	1.32
1	MT	144	ALA	CA-CB	-6.37	1.43	1.53
1	MT	372	ARG	CZ-NH1	-6.35	1.23	1.32
1	MT	134	GLY	N-CA	-6.35	1.38	1.45
1	MT	124	ARG	CZ-NH1	-6.34	1.23	1.32
1	MT	100	GLY	N-CA	-6.30	1.38	1.45
1	MT	105	ALA	CA-CB	-6.26	1.43	1.53
1	MT	178	ARG	CZ-NH1	-6.21	1.24	1.32
1	MT	118	ALA	CA-CB	-6.16	1.43	1.53
1	MT	142	ALA	CA-CB	-6.04	1.43	1.53
1	MT	234	ALA	CA-CB	-5.83	1.43	1.53
1	MT	269	PRO	CA-CB	-5.64	1.46	1.53
1	MT	139	ARG	CD-NE	-5.51	1.38	1.46
1	MT	98	GLY	N-CA	-5.47	1.38	1.45
1	MT	170	ARG	CD-NE	-5.46	1.38	1.46
1	MT	128	ARG	CD-NE	-5.43	1.38	1.46
1	MT	178	ARG	CD-NE	-5.42	1.38	1.46
1	MT	124	ARG	CD-NE	-5.37	1.38	1.46
1	MT	372	ARG	CD-NE	-5.29	1.38	1.46
1	MT	323	GLY	N-CA	-5.28	1.37	1.45
1	MT	153	LYS	N-CA	-5.26	1.41	1.46
1	MT	250	ARG	CD-NE	-5.24	1.39	1.46
1	MT	153	LYS	CA-CB	-5.18	1.47	1.53

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	a	38	LEU	O-C-N	-20.39	100.51	122.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	MT	22	GLY	N-CA-C	-7.80	104.66	113.79
2	a	38	LEU	CA-C-N	7.77	136.65	121.41
2	a	38	LEU	C-N-CA	7.77	136.65	121.41
1	MT	135	ASP	CA-CB-CG	7.36	119.96	112.60
1	MT	174	ASP	CA-CB-CG	7.30	119.90	112.60
1	MT	307	ASN	CA-CB-CG	-7.16	105.44	112.60
1	MT	177	HIS	CA-CB-CG	7.05	120.85	113.80
1	MT	111	ARG	CA-C-N	6.85	126.89	119.90
1	MT	111	ARG	C-N-CA	6.85	126.89	119.90
1	MT	125	ASP	CA-CB-CG	6.84	119.44	112.60
1	MT	279	ASN	CA-CB-CG	-6.52	106.08	112.60
1	MT	151	GLY	N-CA-C	6.49	119.32	110.69
1	MT	275	ASP	CA-CB-CG	6.31	118.91	112.60
1	MT	372	ARG	CD-NE-CZ	6.11	132.95	124.40
1	MT	250	ARG	CD-NE-CZ	6.11	132.95	124.40
1	MT	145	PHE	CA-CB-CG	5.81	119.61	113.80
1	MT	259	GLY	CA-C-N	5.81	128.07	120.28
1	MT	259	GLY	C-N-CA	5.81	128.07	120.28
1	MT	143	ASN	CA-CB-CG	5.80	118.40	112.60
1	MT	246	GLU	CA-CB-CG	5.75	125.61	114.10
1	MT	244	SER	CA-C-N	5.69	127.83	120.44
1	MT	244	SER	C-N-CA	5.69	127.83	120.44
1	MT	314	ARG	CA-C-N	5.67	132.38	121.54
1	MT	314	ARG	C-N-CA	5.67	132.38	121.54
1	MT	180	TYR	N-CA-CB	5.50	118.14	109.94
1	MT	178	ARG	CA-C-N	5.47	127.61	120.28
1	MT	178	ARG	C-N-CA	5.47	127.61	120.28
1	MT	233	ASP	CA-CB-CG	5.46	118.06	112.60
1	MT	369	HIS	CE1-NE2-CD2	-5.39	103.61	109.00
1	MT	60	LYS	CB-CA-C	5.35	115.26	110.44
1	MT	166	SER	CA-C-N	5.35	127.39	120.44
1	MT	166	SER	C-N-CA	5.35	127.39	120.44
1	MT	177	HIS	CE1-NE2-CD2	-5.34	103.66	109.00
1	MT	138	LEU	CA-C-N	5.33	127.42	120.28
1	MT	138	LEU	C-N-CA	5.33	127.42	120.28
1	MT	274	VAL	N-CA-CB	5.31	116.40	110.51
1	MT	369	HIS	CG-CD2-NE2	5.28	112.48	107.20
1	MT	276	TRP	N-CA-CB	5.27	117.71	110.07
1	MT	177	HIS	ND1-CG-CD2	-5.25	100.85	106.10
1	MT	179	LEU	N-CA-CB	5.25	117.84	110.12
1	MT	175	GLU	N-CA-CB	5.25	117.62	110.01
1	MT	369	HIS	ND1-CG-CD2	-5.23	100.87	106.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	MT	271	SER	CA-C-N	5.21	127.22	120.44
1	MT	271	SER	C-N-CA	5.21	127.22	120.44
1	MT	170	ARG	CA-C-N	5.21	127.21	120.44
1	MT	170	ARG	C-N-CA	5.21	127.21	120.44
1	MT	131	LYS	CA-C-N	5.19	127.10	120.56
1	MT	131	LYS	C-N-CA	5.19	127.10	120.56
1	MT	177	HIS	CG-CD2-NE2	5.19	112.39	107.20
1	MT	131	LYS	N-CA-CB	5.16	117.49	110.01
1	MT	146	LEU	N-CA-CB	5.16	117.80	110.06
1	MT	127	PRO	CA-C-N	5.15	127.14	120.44
1	MT	127	PRO	C-N-CA	5.15	127.14	120.44
1	MT	249	ALA	CA-C-N	5.14	127.12	120.44
1	MT	249	ALA	C-N-CA	5.14	127.12	120.44
1	MT	132	ILE	N-CA-CB	5.10	116.52	110.55
1	MT	178	ARG	CA-CB-CG	5.07	124.25	114.10
1	MT	173	SER	CA-C-N	5.06	127.02	120.44
1	MT	173	SER	C-N-CA	5.06	127.02	120.44
1	MT	27	ASN	CA-CB-CG	-5.05	107.55	112.60
1	MT	137	MET	N-CA-CB	5.04	117.31	110.01
1	MT	128	ARG	N-CA-CB	5.02	117.29	110.01
1	MT	335	ARG	NE-CZ-NH2	5.02	123.72	119.20
1	MT	151	GLY	CA-C-N	5.01	127.26	120.65
1	MT	151	GLY	C-N-CA	5.01	127.26	120.65

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	MT	165	PHE	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	MT	2866	0	2877	27	0
2	A	308	0	338	0	0
2	X	308	0	338	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Z	308	0	338	0	0
2	a	308	0	338	1	0
2	b	308	0	338	0	0
2	c	308	0	338	16	0
2	d	308	0	338	1	0
2	e	308	0	338	0	0
2	f	308	0	338	0	0
2	g	308	0	338	0	0
2	h	308	0	338	0	0
2	i	308	0	338	0	0
2	j	308	0	338	1	0
2	k	308	0	338	0	0
2	l	308	0	338	0	0
2	m	308	0	338	0	0
All	All	7794	0	8285	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:MT:87:SER:H	2:c:41:LYS:NZ	1.39	1.18
1:MT:87:SER:HB3	2:c:41:LYS:HZ3	1.15	1.12
1:MT:87:SER:CB	2:c:41:LYS:HZ3	1.65	1.07
1:MT:87:SER:OG	2:c:44:LEU:CD2	2.05	1.04
1:MT:87:SER:OG	2:c:44:LEU:HD22	1.59	1.02
1:MT:87:SER:CB	2:c:41:LYS:NZ	2.27	0.97
1:MT:87:SER:N	2:c:41:LYS:NZ	2.15	0.94
1:MT:87:SER:H	2:c:41:LYS:HZ1	0.89	0.87
1:MT:243:SER:OG	1:MT:246:GLU:OE2	1.95	0.83
1:MT:87:SER:N	2:c:41:LYS:HZ1	1.74	0.80
1:MT:87:SER:HB3	2:c:41:LYS:NZ	1.95	0.73
1:MT:87:SER:HB2	2:c:41:LYS:NZ	2.05	0.69
1:MT:87:SER:HB2	2:c:41:LYS:HZ2	1.59	0.67
1:MT:216:HIS:CD2	1:MT:217:THR:H	2.18	0.61
1:MT:216:HIS:CD2	1:MT:310:PHE:CE2	2.92	0.57
1:MT:170:ARG:NH1	1:MT:246:GLU:OE2	2.38	0.57
1:MT:87:SER:CA	2:c:41:LYS:NZ	2.68	0.56
1:MT:87:SER:CA	2:c:41:LYS:HZ3	2.19	0.56
1:MT:358:LEU:H	1:MT:358:LEU:HD23	1.75	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:MT:242:LYS:NZ	1:MT:242:LYS:HB3	2.26	0.50
1:MT:270:TRP:O	1:MT:274:VAL:HG23	2.16	0.46
1:MT:87:SER:OG	2:c:44:LEU:HD21	2.11	0.43
1:MT:163:LEU:HD12	1:MT:246:GLU:HB3	2.01	0.42
1:MT:240:ILE:HG13	1:MT:240:ILE:O	2.20	0.41
1:MT:287:SER:HB2	1:MT:290:THR:H	1.84	0.41
1:MT:219:ILE:H	1:MT:219:ILE:HD12	1.85	0.41
2:c:18:ILE:HD13	2:j:42:LEU:HD22	2.03	0.40
1:MT:219:ILE:HD12	1:MT:219:ILE:N	2.36	0.40
2:a:34:ALA:HB1	2:d:46:PHE:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	MT	370/372 (100%)	351 (95%)	19 (5%)	0	100	100
2	A	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	X	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	Z	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	a	43/45 (96%)	39 (91%)	3 (7%)	1 (2%)	5	28
2	b	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	c	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	d	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	e	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	f	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	g	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	h	43/45 (96%)	42 (98%)	1 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	i	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	j	43/45 (96%)	41 (95%)	2 (5%)	0	100	100
2	k	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	l	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
2	m	43/45 (96%)	42 (98%)	1 (2%)	0	100	100
All	All	1058/1092 (97%)	1019 (96%)	38 (4%)	1 (0%)	49	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	a	39	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	MT	313/313 (100%)	304 (97%)	9 (3%)	37	58
2	A	31/31 (100%)	31 (100%)	0	100	100
2	X	31/31 (100%)	31 (100%)	0	100	100
2	Z	31/31 (100%)	31 (100%)	0	100	100
2	a	31/31 (100%)	31 (100%)	0	100	100
2	b	31/31 (100%)	31 (100%)	0	100	100
2	c	31/31 (100%)	31 (100%)	0	100	100
2	d	31/31 (100%)	31 (100%)	0	100	100
2	e	31/31 (100%)	31 (100%)	0	100	100
2	f	31/31 (100%)	31 (100%)	0	100	100
2	g	31/31 (100%)	31 (100%)	0	100	100
2	h	31/31 (100%)	31 (100%)	0	100	100
2	i	31/31 (100%)	31 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	j	31/31 (100%)	31 (100%)	0	100	100
2	k	31/31 (100%)	31 (100%)	0	100	100
2	l	31/31 (100%)	31 (100%)	0	100	100
2	m	31/31 (100%)	31 (100%)	0	100	100
All	All	809/809 (100%)	800 (99%)	9 (1%)	63	76

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	MT	74	ARG
1	MT	82	TYR
1	MT	88	LEU
1	MT	200	GLU
1	MT	203	VAL
1	MT	211	PHE
1	MT	246	GLU
1	MT	290	THR
1	MT	322	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	MT	206	HIS
1	MT	209	ASN
1	MT	307	ASN
2	m	43	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-42136. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300




Y Index: 300




Z Index: 300


6.2.2 Raw map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 268




Y Index: 323




Z Index: 327


6.3.2 Raw map



X Index: 267



Y Index: 322




Z Index: 328


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X




Y




Z

6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00127. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 7724 nm^3 ; this corresponds to an approximate mass of 6977 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.155 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.155 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.45	-	-
Author-provided FSC curve	6.35	8.45	6.64
Unmasked-calculated*	6.31	8.37	6.62

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-42136 and PDB model 8UCR. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.00127) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9760	 0.1580
A	 0.9710	 0.1100
MT	 0.9860	 0.2390
X	 0.9610	 0.0930
Z	 0.9150	 0.1330
a	 0.9580	 0.1520
b	 0.9940	 0.1070
c	 0.8990	 0.1170
d	 0.9740	 0.0860
e	 0.9970	 0.1240
f	 0.9480	 0.1300
g	 0.9870	 0.1260
h	 0.9710	 0.1390
i	 0.9970	 0.1320
j	 0.9770	 0.1350
k	 0.9940	 0.0500
l	 0.9900	 0.0590
m	 0.9840	 0.0870

