



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 29, 2026 – 12:05 PM UTC

PDB ID : 4UD2 / pdb_00004ud2
Title : Structure of anaerobically purified *D. fructosovorans* NiFe- hydrogenase
Authors : Volbeda, A.; Martin, L.; Liebgott, P.-P.; Fontecilla-Camps, J.C.
Deposited on : 2014-12-05
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	91% (Green), 9% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
1	B	264	92% (Green), 8% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
1	C	264	88% (Green), 10% (Yellow), 0% (Orange), 0% (Red), 2% (Grey)
2	Q	549	93% (Green), 6% (Yellow), 0% (Orange), 0% (Red), 1% (Grey)
2	R	549	91% (Green), 8% (Yellow), 0% (Orange), 0% (Red), 1% (Grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	S	549	 <p>A horizontal bar chart representing the quality of chain. The bar is 89% green and 11% yellow. A small red square is at the start, and a small grey square is at the end. The percentage values '89%' and '11%' are printed below the bar.</p>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 19572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYDROGENASE (NIFE) SMALL SUBUNIT HYDA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	Total 1973	C 1256	N 330	O 372	S 15	0	0	0
1	B	262	Total 1983	C 1262	N 331	O 375	S 15	0	1	0
1	C	260	Total 1960	C 1248	N 327	O 370	S 15	0	0	0

- Molecule 2 is a protein called NICKEL-DEPENDENT HYDROGENASE LARGE SUB-UNIT.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	Q	544	Total 4163	C 2652	N 723	O 766	S 22	0	0	0
2	R	544	Total 4168	C 2655	N 724	O 767	S 22	0	1	0
2	S	544	Total 4163	C 2652	N 723	O 766	S 22	0	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	Q	1	Total C O 6 3 3	0	0
3	Q	1	Total C O 6 3 3	0	0
3	Q	1	Total C O 6 3 3	0	0
3	Q	1	Total C O 6 3 3	0	0
3	R	1	Total C O 6 3 3	0	0
3	R	1	Total C O 6 3 3	0	0
3	R	1	Total C O 6 3 3	0	0
3	R	1	Total C O 6 3 3	0	0
3	S	1	Total C O 6 3 3	0	0
3	S	1	Total C O 6 3 3	0	0

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe S	0	0
			8	4 4		
4	A	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		
4	C	1	Total	Fe S	0	0
			8	4 4		
4	C	1	Total	Fe S	0	0
			8	4 4		

- Molecule 5 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Fe	S	0	0
			7	3	4		
5	B	1	Total	Fe	S	0	0
			7	3	4		
5	C	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 6 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: C_3FeN_2O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	Q	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	R	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	S	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 7 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	Q	1	Total	Ni	0	0
			1	1		
7	R	1	Total	Ni	0	0
			1	1		
7	S	1	Total	Ni	0	0
			1	1		

- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	Q	1	Total	Mg	0	0
			1	1		
8	R	1	Total	Mg	0	0
			1	1		
8	S	1	Total	Mg	0	0
			1	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	184	Total	O	0	0
			184	184		
9	B	115	Total	O	0	0
			115	115		
9	C	122	Total	O	0	0
			122	122		
9	Q	239	Total	O	0	0
			239	239		
9	R	206	Total	O	0	0
			206	206		
9	S	116	Total	O	0	0
			116	116		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HYDROGENASE (NIFE) SMALL SUBUNIT HYDA

Chain A: 




- Molecule 1: HYDROGENASE (NIFE) SMALL SUBUNIT HYDA

Chain B: 




- Molecule 1: HYDROGENASE (NIFE) SMALL SUBUNIT HYDA

Chain C: 



- Molecule 2: NICKEL-DEPENDENT HYDROGENASE LARGE SUBUNIT

Chain Q: 





- Molecule 2: NICKEL-DEPENDENT HYDROGENASE LARGE SUBUNIT

Chain R: 





● Molecule 2: NICKEL-DEPENDENT HYDROGENASE LARGE SUBUNIT



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.01Å 100.17Å 183.98Å 90.00° 91.42° 90.00°	Depositor
Resolution (Å)	25.00 – 2.30 25.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	88.7 (25.00-2.30) 88.7 (25.00-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.26 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.173 , 0.207 0.176 , 0.211	Depositor DCC
R_{free} test set	4631 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtrriage
Anisotropy	0.702	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.046 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19572	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SF4, F3S, FCO, CSO, NI, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.72	1/2027 (0.0%)	0.86	1/2759 (0.0%)
1	B	0.61	0/2037	0.83	0/2773
1	C	0.64	0/2014	0.84	2/2743 (0.1%)
2	Q	0.67	0/4260	0.85	3/5781 (0.1%)
2	R	0.64	0/4268	0.87	4/5792 (0.1%)
2	S	0.61	0/4260	0.86	4/5781 (0.1%)
All	All	0.65	1/18866 (0.0%)	0.85	14/25629 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	TYR	CG-CD1	-5.04	1.28	1.39

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	502	ASN	N-CA-C	6.99	118.69	111.14
2	S	502	ASN	N-CA-C	6.42	118.08	111.14
1	C	235	VAL	N-CA-C	5.85	119.27	113.71
2	R	502	ASN	N-CA-C	5.70	117.57	111.36
2	S	397	GLN	CA-C-N	5.54	125.00	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1911	15	0
1	B	1983	0	1920	13	0
1	C	1960	0	1897	17	0
2	Q	4163	0	4136	23	0
2	R	4168	0	4142	25	0
2	S	4163	0	4136	34	0
3	A	24	0	32	1	0
3	Q	24	0	32	3	0
3	R	24	0	32	2	0
3	S	12	0	16	2	0
4	A	16	0	0	0	0
4	B	16	0	0	0	0
4	C	16	0	0	0	0
5	A	7	0	0	0	0
5	B	7	0	0	0	0
5	C	7	0	0	0	0
6	Q	7	0	0	0	0
6	R	7	0	0	0	0
6	S	7	0	0	0	0
7	Q	1	0	0	0	0
7	R	1	0	0	0	0
7	S	1	0	0	0	0
8	Q	1	0	0	0	0
8	R	1	0	0	0	0
8	S	1	0	0	0	0
9	A	184	0	0	5	0
9	B	115	0	0	1	0
9	C	122	0	0	4	0
9	Q	239	0	0	3	0
9	R	206	0	0	2	0
9	S	116	0	0	1	0
All	All	19572	0	18254	126	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:Q:503:LEU:HG	2:Q:515:VAL:HG21	1.66	0.77
2:S:134:LEU:HD22	2:S:168:LYS:HG2	1.71	0.71
2:S:361:LYS:HG2	2:S:362:LEU:O	1.93	0.68
2:R:410:LYS:HG2	3:R:605:GOL:H12	1.78	0.65
2:Q:319:ARG:HH22	3:Q:606:GOL:H2	1.65	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/264 (98%)	252 (97%)	8 (3%)	0	100	100
1	B	261/264 (99%)	251 (96%)	10 (4%)	0	100	100
1	C	258/264 (98%)	251 (97%)	7 (3%)	0	100	100
2	Q	541/549 (98%)	527 (97%)	14 (3%)	0	100	100
2	R	542/549 (99%)	527 (97%)	15 (3%)	0	100	100
2	S	541/549 (98%)	523 (97%)	18 (3%)	0	100	100
All	All	2403/2439 (98%)	2331 (97%)	72 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/210 (99%)	208 (100%)	0	100	100
1	B	209/210 (100%)	209 (100%)	0	100	100
1	C	207/210 (99%)	205 (99%)	2 (1%)	68	82
2	Q	433/438 (99%)	432 (100%)	1 (0%)	87	94
2	R	434/438 (99%)	432 (100%)	2 (0%)	81	90
2	S	433/438 (99%)	430 (99%)	3 (1%)	76	87
All	All	1924/1944 (99%)	1916 (100%)	8 (0%)	84	92

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	S	473	ASP
2	S	410	LYS
2	R	473	ASP
2	R	402	LYS
2	S	245	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	S	9	GLN
2	S	241	GLN
1	C	240	GLN
2	Q	47	GLN
2	Q	509	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSO	Q	543	2,7	3,6,7	0.82	0	1,6,8	0.05	0
2	CSO	R	543	2,7	3,6,7	1.02	0	1,6,8	0.32	0
2	CSO	S	543	2,7	3,6,7	0.75	0	1,6,8	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	Q	543	2,7	-	1/1/5/7	-
2	CSO	R	543	2,7	-	1/1/5/7	-
2	CSO	S	543	2,7	-	1/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Q	543	CSO	N-CA-CB-SG
2	R	543	CSO	N-CA-CB-SG
2	S	543	CSO	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 6 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SF4	A	1267	1	0,12,12	-	-	-		
6	FCO	Q	1550	2,7	0,6,6	-	-	-		
4	SF4	B	1267	1	0,12,12	-	-	-		
4	SF4	C	1265	1	0,12,12	-	-	-		
4	SF4	A	1265	1	0,12,12	-	-	-		
3	GOL	R	607	-	5,5,5	0.31	0	5,5,5	0.27	0
3	GOL	S	604	-	5,5,5	0.39	0	5,5,5	0.22	0
6	FCO	R	1550	2,7	0,6,6	-	-	-		
3	GOL	R	605	-	5,5,5	0.27	0	5,5,5	0.19	0
5	F3S	A	1266	1	0,9,9	-	-	-		
5	F3S	B	1266	1	0,9,9	-	-	-		
4	SF4	C	1267	1	0,12,12	-	-	-		
3	GOL	A	306	-	5,5,5	0.37	0	5,5,5	0.32	0
3	GOL	A	305	-	5,5,5	0.41	0	5,5,5	0.70	0
3	GOL	Q	604	-	5,5,5	0.25	0	5,5,5	0.15	0
6	FCO	S	1550	2,7	0,6,6	-	-	-		
3	GOL	Q	606	-	5,5,5	0.48	0	5,5,5	0.71	0
3	GOL	Q	605	-	5,5,5	0.19	0	5,5,5	0.26	0
5	F3S	C	1266	1	0,9,9	-	-	-		
4	SF4	B	1265	1	0,12,12	-	-	-		
3	GOL	R	606	-	5,5,5	0.44	0	5,5,5	0.36	0
3	GOL	Q	607	-	5,5,5	0.34	0	5,5,5	0.21	0
3	GOL	A	304	-	5,5,5	0.25	0	5,5,5	0.42	0
3	GOL	A	307	-	5,5,5	0.45	0	5,5,5	0.26	0
3	GOL	R	604	-	5,5,5	0.36	0	5,5,5	0.34	0
3	GOL	S	605	-	5,5,5	0.24	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	1267	1	-	-	0/6/5/5
4	SF4	B	1267	1	-	-	0/6/5/5
4	SF4	C	1265	1	-	-	0/6/5/5
4	SF4	A	1265	1	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	R	607	-	-	2/4/4/4	-
3	GOL	S	604	-	-	1/4/4/4	-
3	GOL	R	605	-	-	4/4/4/4	-
5	F3S	A	1266	1	-	-	0/3/3/3
5	F3S	B	1266	1	-	-	0/3/3/3
4	SF4	C	1267	1	-	-	0/6/5/5
3	GOL	A	306	-	-	4/4/4/4	-
3	GOL	A	305	-	-	2/4/4/4	-
3	GOL	Q	604	-	-	0/4/4/4	-
3	GOL	Q	606	-	-	3/4/4/4	-
3	GOL	Q	605	-	-	2/4/4/4	-
5	F3S	C	1266	1	-	-	0/3/3/3
4	SF4	B	1265	1	-	-	0/6/5/5
3	GOL	R	606	-	-	0/4/4/4	-
3	GOL	Q	607	-	-	2/4/4/4	-
3	GOL	A	304	-	-	2/4/4/4	-
3	GOL	A	307	-	-	4/4/4/4	-
3	GOL	R	604	-	-	0/4/4/4	-
3	GOL	S	605	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	304	GOL	C1-C2-C3-O3
3	A	304	GOL	O2-C2-C3-O3
3	A	305	GOL	C1-C2-C3-O3
3	A	306	GOL	C1-C2-C3-O3
3	A	307	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	R	605	GOL	2	0
3	A	305	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Q	606	GOL	1	0
3	Q	607	GOL	2	0
3	S	605	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/264 (99%)	-0.49	0 100 100	17, 26, 40, 69	9 (3%)
1	B	262/264 (99%)	-0.08	1 (0%) 88 89	22, 39, 59, 73	11 (4%)
1	C	260/264 (98%)	-0.10	0 100 100	21, 37, 57, 81	8 (3%)
2	Q	543/549 (98%)	-0.43	0 100 100	16, 29, 46, 57	15 (2%)
2	R	543/549 (98%)	-0.27	0 100 100	18, 34, 50, 67	12 (2%)
2	S	543/549 (98%)	0.14	4 (0%) 84 85	22, 47, 69, 87	21 (3%)
All	All	2413/2439 (98%)	-0.20	5 (0%) 91 91	16, 35, 61, 87	76 (3%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	S	174	GLY	2.6
1	B	264	GLY	2.4
2	S	323	LYS	2.2
2	S	363	ASP	2.1
2	S	336	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	S	543	7/8	0.97	0.07	30,31,36,36	1
2	CSO	R	543	7/8	0.98	0.05	20,22,26,27	1
2	CSO	Q	543	7/8	0.98	0.07	20,21,22,25	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	306	6/6	0.79	0.12	58,61,68,75	0
3	GOL	A	307	6/6	0.79	0.15	57,66,68,77	0
3	GOL	Q	606	6/6	0.79	0.14	56,62,65,69	0
3	GOL	R	607	6/6	0.82	0.14	51,57,58,64	0
3	GOL	R	605	6/6	0.83	0.12	58,60,69,72	0
3	GOL	Q	607	6/6	0.85	0.14	62,67,70,73	0
3	GOL	S	605	6/6	0.85	0.12	56,59,64,72	0
3	GOL	A	305	6/6	0.87	0.12	54,55,61,65	0
3	GOL	A	304	6/6	0.89	0.12	47,56,71,76	0
3	GOL	S	604	6/6	0.90	0.08	36,38,39,39	0
3	GOL	R	606	6/6	0.90	0.10	38,41,42,42	0
3	GOL	R	604	6/6	0.96	0.05	29,30,32,32	0
3	GOL	Q	604	6/6	0.96	0.07	32,35,38,41	0
3	GOL	Q	605	6/6	0.96	0.07	31,32,36,38	0
8	MG	R	1553	1/1	0.96	0.04	23,23,23,23	0
4	SF4	B	1265	8/8	0.97	0.05	40,42,48,52	0
4	SF4	B	1267	8/8	0.98	0.03	25,26,28,28	0
5	F3S	B	1266	7/7	0.98	0.04	35,37,39,39	0
8	MG	Q	1553	1/1	0.98	0.03	21,21,21,21	0
4	SF4	A	1265	8/8	0.98	0.04	18,20,22,23	0
5	F3S	A	1266	7/7	0.99	0.03	16,19,20,22	0
4	SF4	A	1267	8/8	0.99	0.02	15,16,17,18	0
5	F3S	C	1266	7/7	0.99	0.03	23,23,25,27	0
6	FCO	Q	1550	7/7	0.99	0.04	15,17,19,20	0
6	FCO	R	1550	7/7	0.99	0.04	17,19,21,23	0
6	FCO	S	1550	7/7	0.99	0.06	25,29,32,33	0
7	NI	Q	1551	1/1	0.99	0.03	18,18,18,18	0
7	NI	S	1551	1/1	0.99	0.01	28,28,28,28	0
4	SF4	C	1265	8/8	0.99	0.03	22,24,25,26	0
4	SF4	C	1267	8/8	0.99	0.02	22,23,24,26	0
8	MG	S	1553	1/1	0.99	0.02	32,32,32,32	0
7	NI	R	1551	1/1	1.00	0.01	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.