



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:47 PM UTC

PDB ID : 4UDD / pdb_00004udd
Title : GR in complex with desisobutyrylciclesonide
Authors : Edman, K.; Hogner, A.; Hussein, A.; Bjursell, M.; Aagaard, A.; Backstrom, S.; Bodin, C.; Wissler, L.; Jellesmark-Jensen, T.; Cavallin, A.; Karlsson, U.; Nilsson, E.; Lecina, D.; Takahashi, R.; Grebner, C.; Lepisto, M.; Guallar, V.
Deposited on : 2014-12-09
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	
2	B	14	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOCORTICOID RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	4	0
			2062	1322	341	380	19			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	498	GLY	-	expression tag	UNP P04150
A	499	SER	-	expression tag	UNP P04150
A	517	ASP	ASN	engineered mutation	UNP P04150
A	571	MET	VAL	engineered mutation	UNP P04150
A	602	SER	PHE	engineered mutation	UNP P04150
A	638	ASP	CYS	engineered mutation	UNP P04150

- Molecule 2 is a protein called NUCLEAR RECEPTOR COACTIVATOR 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	13	Total	C	N	O	0	1	0
			122	75	22	25			


- Molecule 3 is 3-[(3-CHOLAMIDOPROPYL)DIMETHYLAMMONIO]-1-PROPANESULFONATE (CCD ID: CPS) (formula: C₃₂H₅₈N₂O₇S).



-
- EDO
- Chemical structure of EDO (Ethane-1,2-diol) showing a zigzag conformation. The carbon atoms are labeled C1 and C2 in green. The hydroxyl groups are labeled HO (O1) and OH (O2) in red. The oxygen atoms are labeled O1 and O2 in green.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DESISOBUTYRYL CICLESONIDE (CCD ID: CV7) (formula: C₂₈H₃₈O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			34	28	6		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	246	Total	O	0	0
			246	246		
6	B	4	Total	O	0	0
			4	4		

i

- Molecule 1: GLUCOCORTICOID RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.20Å 87.20Å 102.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.14 – 1.80 40.14 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.14-1.80) 92.1 (40.14-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.24 (at 1.79Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.213 , 0.224 0.192 , 0.206	Depositor DCC
R_{free} test set	2138 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.392	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2580	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CV7, CPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	2/2106 (0.1%)	1.10	3/2844 (0.1%)
2	B	0.64	0/122	1.39	2/161 (1.2%)
All	All	0.80	2/2228 (0.1%)	1.12	5/3005 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	560	MET	SD-CE	-8.23	1.58	1.79
1	A	761	ILE	CA-CB	5.45	1.56	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	704	ARG	CA-C-N	7.47	132.97	122.36
1	A	704	ARG	C-N-CA	7.47	132.97	122.36
2	B	752	ASP	CA-C-N	5.32	131.27	121.70
2	B	752	ASP	C-N-CA	5.32	131.27	121.70
1	A	619	ASN	CA-CB-CG	5.22	117.82	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2062	0	2071	8	0
2	B	122	0	119	2	0
3	A	108	0	156	1	0
4	A	4	0	6	0	0
5	A	34	0	38	2	0
6	A	246	0	0	0	0
6	B	4	0	0	0	0
All	All	2580	0	2390	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:646:MET:HE1	5:A:1782:CV7:H212	1.84	0.60
1:A:557:TRP:HA	1:A:560:MET:CE	2.37	0.55
1:A:602[B]:SER:OG	1:A:729:VAL:HG21	2.14	0.47
1:A:759:ASN:HB2	2:B:742:ASN:HD21	1.80	0.46
1:A:560:MET:HG2	5:A:1782:CV7:H251	2.01	0.43
3:A:1779:CPS:H29A	3:A:1779:CPS:H262	1.91	0.41
1:A:579:LYS:HD2	2:B:749:LEU:HA	2.02	0.40
1:A:557:TRP:HA	1:A:560:MET:HE3	2.02	0.40
1:A:610:TRP:HB2	1:A:663:TYR:CE1	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/280 (90%)	246 (98%)	5 (2%)	0	100	100
2	B	12/14 (86%)	12 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	263/294 (90%)	258 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/255 (91%)	232 (100%)	1 (0%)	84	83
2	B	13/13 (100%)	13 (100%)	0	100	100
All	All	246/268 (92%)	245 (100%)	1 (0%)	84	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	620	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	570	GLN
1	A	586	ASN
1	A	597	GLN
1	A	711	ASN
1	A	759	ASN
1	A	768	ASN
2	B	742	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CPS	A	1778	-	39,39,45	0.73	0	61,61,70	1.18	7 (11%)
4	EDO	A	1781	-	3,3,3	0.46	0	2,2,2	0.50	0
5	CV7	A	1782	-	38,39,39	1.11	1 (2%)	57,63,63	1.35	7 (12%)
3	CPS	A	1780	-	39,39,45	0.58	0	61,61,70	1.29	6 (9%)
3	CPS	A	1779	-	39,39,45	0.66	0	61,61,70	1.26	6 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CPS	A	1778	-	-	9/17/82/90	0/4/4/4
4	EDO	A	1781	-	-	0/1/1/1	-
5	CV7	A	1782	-	-	0/12/98/98	0/6/6/6
3	CPS	A	1780	-	-	7/17/82/90	0/4/4/4
3	CPS	A	1779	-	-	5/17/82/90	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1782	CV7	C8-C6	2.56	1.58	1.54

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1782	CV7	C31-C32-C38	3.80	118.85	115.86
3	A	1780	CPS	C14-C15-C2	3.48	116.36	112.66
3	A	1778	CPS	C21-C20-C22	-3.35	105.15	110.34
5	A	1782	CV7	C3-C2-C15	-3.23	113.14	115.86
3	A	1779	CPS	C11-C2-C1	-3.13	103.33	108.31
5	A	1782	CV7	C32-C33-C34	-3.10	119.90	122.72
3	A	1778	CPS	C11-C2-C1	-3.04	103.47	108.31
3	A	1779	CPS	C8-C9-C20	3.01	116.74	112.18
3	A	1780	CPS	C16-C15-C14	-3.01	107.79	111.23
3	A	1779	CPS	C10-C5-C6	3.00	115.90	111.20
3	A	1780	CPS	C11-C2-C1	-2.75	103.93	108.31
3	A	1779	CPS	C22-C23-C24	-2.63	107.20	113.06
3	A	1779	CPS	C14-C15-C2	2.63	115.46	112.66
3	A	1778	CPS	C7-C6-C5	2.50	105.96	103.54
3	A	1780	CPS	C7-C6-C5	2.48	105.95	103.54
5	A	1782	CV7	C36-C34-C33	2.33	119.49	117.12
5	A	1782	CV7	C39-C38-C6	2.32	116.54	113.66
3	A	1778	CPS	C9-C5-C6	-2.29	97.82	100.11
3	A	1780	CPS	C12-C1-C2	2.23	116.51	112.74
3	A	1780	CPS	C16-C15-C2	2.23	115.03	112.66
3	A	1779	CPS	C16-C17-C18	2.23	113.93	111.50
5	A	1782	CV7	C22-C21-C20	-2.09	108.17	111.89
3	A	1778	CPS	C8-C7-C6	-2.08	101.07	105.14
3	A	1778	CPS	C11-C2-C19	2.05	113.93	111.18
3	A	1778	CPS	C21-C20-C9	2.01	115.91	112.88
5	A	1782	CV7	C31-C32-C33	-2.01	117.64	120.86

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1778	CPS	C23-C24-N1-C25
3	A	1778	CPS	O1-C24-N1-C25
3	A	1779	CPS	C23-C24-N1-C25
3	A	1779	CPS	O1-C24-N1-C25
3	A	1780	CPS	C23-C24-N1-C25
3	A	1780	CPS	O1-C24-N1-C25

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1778	CPS	C26-C25-N1-C24
3	A	1778	CPS	C9-C20-C22-C23
3	A	1778	CPS	C21-C20-C22-C23
3	A	1780	CPS	N1-C25-C26-C27
3	A	1778	CPS	C25-C26-C27-N2
3	A	1779	CPS	C25-C26-C27-N2
3	A	1780	CPS	C25-C26-C27-N2
3	A	1780	CPS	C26-C27-N2-C28
3	A	1780	CPS	C26-C27-N2-C29
3	A	1780	CPS	C26-C27-N2-C30
3	A	1779	CPS	C9-C20-C22-C23
3	A	1779	CPS	C22-C20-C9-C8
3	A	1778	CPS	N1-C25-C26-C27
3	A	1778	CPS	C26-C27-N2-C29
3	A	1778	CPS	C26-C27-N2-C28


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1782	CV7	2	0
3	A	1779	CPS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/280 (88%)	0.16	12 (4%) 35 34	10, 30, 56, 68	4 (1%)
2	B	13/14 (92%)	1.23	3 (23%) 2 1	19, 40, 61, 73	1 (7%)
All	All	262/294 (89%)	0.22	15 (5%) 29 28	10, 31, 57, 73	5 (1%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	618	ALA	5.8
1	A	529	THR	4.4
1	A	712	TRP	4.3
1	A	617	SER	4.1
1	A	619	ASN	3.8
2	B	742	ASN	3.6
1	A	616	SER	3.3
1	A	709	SER	3.2
1	A	716	TYR	2.7
1	A	678	ASP	2.6
2	B	746[A]	ARG	2.3
1	A	705	GLU	2.2
2	B	743	ALA	2.2
1	A	632	GLN	2.1
1	A	638	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CPS	A	1780	36/42	0.78	0.29	83,85,98,98	0
4	EDO	A	1781	4/4	0.85	0.17	54,55,57,58	0
3	CPS	A	1779	36/42	0.89	0.14	34,38,74,75	0
3	CPS	A	1778	36/42	0.91	0.13	26,30,72,72	0
5	CV7	A	1782	34/34	0.97	0.06	18,21,31,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around CPS A 1779:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CPS A 1778:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.