



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:30 AM UTC

PDB ID : 7UDI / pdb_00007udi
Title : Full-length dimer of DNA-Damage Response Protein C from *Deinococcus radiodurans*
Authors : Szabla, R.; Li, M.C.; Junop, M.S.
Deposited on : 2022-03-19
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	6% 77% 12% 10%
1	B	231	20% 80% 16% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA damage response protein DdrC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	Se	0	0	0
			1563	967	295	298	3			
1	B	223	Total	C	N	O	Se	0	0	0
			1670	1027	322	318	3			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	29	Total 29	O 29	0	0
3	B	11	Total 11	O 11	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA damage response protein DdrC



- Molecule 1: DNA damage response protein DdrC



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	66.70Å 66.70Å 129.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.30 – 2.24 59.30 – 2.24	Depositor EDS
% Data completeness (in resolution range)	96.3 (59.30-2.24) 91.3 (59.30-2.24)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.25Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.231 , 0.250 0.231 , 0.250	Depositor DCC
R_{free} test set	1284 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	51.0	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.049 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3293	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	6/1586 (0.4%)	1.25	12/2149 (0.6%)
1	B	1.09	0/1694	1.23	8/2294 (0.3%)
All	All	1.18	6/3280 (0.2%)	1.24	20/4443 (0.5%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	VAL	CA-C	5.81	1.58	1.52
1	A	201	ARG	NE-CZ	5.67	1.39	1.33
1	A	47	LEU	CB-CG	-5.58	1.42	1.53
1	A	48	PRO	N-CA	-5.44	1.40	1.47
1	A	153	ARG	NE-CZ	5.39	1.39	1.33
1	A	34	LEU	CB-CG	-5.04	1.43	1.53

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	141	GLY	N-CA-C	-8.58	102.89	112.33
1	A	4	ALA	CA-C-N	6.34	126.29	119.76
1	A	4	ALA	C-N-CA	6.34	126.29	119.76
1	A	198	SER	N-CA-C	-6.12	102.45	110.53
1	A	222	GLU	CA-C-N	5.96	128.55	120.44
1	A	222	GLU	C-N-CA	5.96	128.55	120.44
1	B	226	GLN	N-CA-C	-5.72	106.34	113.55
1	A	20	ASP	CA-CB-CG	5.70	118.30	112.60
1	A	198	SER	CA-C-N	-5.54	114.15	122.24
1	A	198	SER	C-N-CA	-5.54	114.15	122.24
1	A	99	ARG	N-CA-C	-5.35	105.53	111.36
1	B	64	VAL	CA-C-N	5.17	125.21	119.32
1	B	64	VAL	C-N-CA	5.17	125.21	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	131	MSE	CG-SE-CE	5.12	110.18	98.92
1	B	206	ILE	N-CA-C	-5.10	104.67	111.05
1	A	199	GLU	N-CA-C	-5.05	104.29	110.65
1	B	47	LEU	CA-C-N	5.04	124.94	119.85
1	B	47	LEU	C-N-CA	5.04	124.94	119.85
1	A	108	ASN	CA-C-N	5.01	124.62	119.56
1	A	108	ASN	C-N-CA	5.01	124.62	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1563	0	1531	11	0
1	B	1670	0	1626	27	0
2	A	5	0	0	0	0
2	B	15	0	0	0	0
3	A	29	0	0	0	0
3	B	11	0	0	0	0
All	All	3293	0	3157	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:ALA:HA	1:B:208:THR:HB	1.49	0.94
1:B:205:ALA:HA	1:B:208:THR:CB	2.08	0.82
1:B:143:VAL:HB	1:B:203:ASN:ND2	2.05	0.71
1:B:160:SER:O	1:B:164:ARG:HG3	1.96	0.66
1:B:164:ARG:NH1	1:B:171:ALA:HA	2.13	0.63
1:B:24:HIS:CE1	1:B:26:PRO:HB2	2.34	0.63
1:A:87:ARG:HG2	1:B:72:PHE:CE2	2.36	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:PHE:HA	1:A:107:ARG:HH21	1.68	0.58
1:B:75:ASP:CG	1:B:80:ARG:HH21	2.13	0.56
1:A:127:ARG:O	1:A:131:MSE:HG2	2.07	0.55
1:B:143:VAL:HB	1:B:203:ASN:HD21	1.69	0.54
1:B:202:GLY:O	1:B:206:ILE:HD12	2.07	0.54
1:B:10:PHE:O	1:B:13:VAL:HG22	2.10	0.52
1:B:205:ALA:CA	1:B:208:THR:HB	2.32	0.51
1:B:140:GLU:CB	1:B:203:ASN:ND2	2.75	0.49
1:B:193:ARG:O	1:B:194:ALA:C	2.57	0.48
1:A:150:ILE:HD13	1:A:211:GLU:HG3	1.96	0.47
1:B:156:LEU:HD11	1:B:184:MSE:HE1	1.96	0.47
1:A:195:ILE:O	1:A:198:SER:O	2.33	0.47
1:B:77:PRO:HA	1:B:80:ARG:HD2	1.96	0.46
1:B:179:ALA:O	1:B:183:ARG:HG3	2.16	0.45
1:B:194:ALA:O	1:B:198:SER:N	2.34	0.45
1:B:205:ALA:HA	1:B:208:THR:OG1	2.17	0.45
1:B:198:SER:O	1:B:199:GLU:C	2.60	0.44
1:B:76:THR:O	1:B:80:ARG:HG2	2.18	0.44
1:A:179:ALA:O	1:A:183:ARG:HG3	2.17	0.44
1:B:34:LEU:HD12	1:B:34:LEU:HA	1.78	0.43
1:B:140:GLU:C	1:B:141:GLY:O	2.58	0.43
1:A:24:HIS:CE1	1:A:26:PRO:HB2	2.54	0.43
1:B:203:ASN:O	1:B:207:LEU:HD12	2.18	0.42
1:B:164:ARG:CZ	1:B:171:ALA:HA	2.50	0.42
1:A:181:LEU:C	1:A:181:LEU:HD23	2.45	0.42
1:B:205:ALA:C	1:B:208:THR:H	2.28	0.42
1:A:193:ARG:NH1	1:A:197:GLU:OE2	2.52	0.42
1:A:193:ARG:O	1:A:197:GLU:HG2	2.20	0.41
1:B:142:ARG:O	1:B:146:GLN:HG3	2.21	0.41
1:A:25:ALA:HB3	1:A:26:PRO:HD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/231 (88%)	202 (100%)	1 (0%)	0	100	100
1	B	221/231 (96%)	214 (97%)	6 (3%)	1 (0%)	24	23
All	All	424/462 (92%)	416 (98%)	7 (2%)	1 (0%)	43	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	142	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/172 (88%)	152 (100%)	0	100	100
1	B	160/172 (93%)	157 (98%)	3 (2%)	50	59
All	All	312/344 (91%)	309 (99%)	3 (1%)	68	76

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	81	ARG
1	B	120	ARG
1	B	196	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	HIS
1	B	137	HIS
1	B	196	GLN
1	B	203	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	303	-	4,4,4	0.39	0	6,6,6	0.09	0
2	SO4	B	301	-	4,4,4	0.37	0	6,6,6	0.07	0
2	SO4	A	301	-	4,4,4	0.38	0	6,6,6	0.09	0
2	SO4	B	302	-	4,4,4	0.39	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	204/231 (88%)	0.51	14 (6%)	23 21	39, 56, 84, 98	0
1	B	220/231 (95%)	1.17	46 (20%)	2 2	43, 73, 103, 106	0
All	All	424/462 (91%)	0.86	60 (14%)	6 5	39, 61, 100, 106	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	139	GLY	6.2
1	B	200	ALA	5.3
1	B	227	VAL	4.8
1	A	225	GLY	4.3
1	B	135	ALA	4.2
1	B	164	ARG	3.7
1	B	204	ALA	3.7
1	B	5	PRO	3.7
1	B	206	ILE	3.6
1	A	226	GLN	3.6
1	B	202	GLY	3.6
1	B	144	TYR	3.6
1	B	225	GLY	3.5
1	B	201	ARG	3.4
1	B	195	ILE	3.3
1	A	224	ALA	3.3
1	B	188	ASP	3.0
1	B	224	ALA	3.0
1	B	141	GLY	3.0
1	B	42	LEU	2.9
1	B	207	LEU	2.9
1	B	150	ILE	2.9
1	A	157	GLY	2.9
1	B	214	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	143	VAL	2.9
1	B	198	SER	2.8
1	B	193	ARG	2.8
1	B	137	HIS	2.6
1	B	203	ASN	2.6
1	B	33	GLY	2.6
1	A	109	PRO	2.5
1	B	72	PHE	2.5
1	B	163	VAL	2.4
1	B	210	HIS	2.4
1	B	140	GLU	2.4
1	A	118	ALA	2.3
1	B	192	ALA	2.3
1	A	174	ASP	2.3
1	B	43	VAL	2.3
1	B	147	LEU	2.3
1	B	208	THR	2.3
1	B	142	ARG	2.2
1	A	211	GLU	2.2
1	B	130	LEU	2.2
1	B	205	ALA	2.2
1	A	111	PRO	2.2
1	B	226	GLN	2.2
1	B	196	GLN	2.1
1	A	19	ALA	2.1
1	B	194	ALA	2.1
1	A	201	ARG	2.1
1	B	172	THR	2.1
1	B	171	ALA	2.1
1	B	159	ASP	2.1
1	B	76	THR	2.1
1	A	182	LEU	2.1
1	A	43	VAL	2.0
1	B	138	GLY	2.0
1	A	198	SER	2.0
1	B	81	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	303	5/5	0.64	0.17	71,73,81,93	0
2	SO4	B	302	5/5	0.73	0.11	108,110,114,117	0
2	SO4	B	301	5/5	0.73	0.11	80,87,88,97	0
2	SO4	A	301	5/5	0.80	0.13	81,82,90,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.