



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:41 AM UTC

PDB ID : 5UEC / pdb_00005uec
Title : Crystal Structure of CYP2B6 (Y226H/K262R) in complex with myrtenyl bromide.
Authors : Shah, M.B.; Halpert, J.R.
Deposited on : 2016-12-30
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytochrome P450 2B6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	0	0
			3716	2410	635	655	16			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP P20813
A	21	ALA	GLN	engineered mutation	UNP P20813
A	22	LYS	ARG	engineered mutation	UNP P20813
A	23	LYS	HIS	engineered mutation	UNP P20813
A	24	THR	PRO	engineered mutation	UNP P20813
A	25	SER	ASN	engineered mutation	UNP P20813
A	26	SER	THR	engineered mutation	UNP P20813
A	27	LYS	HIS	engineered mutation	UNP P20813
A	28	GLY	ASP	engineered mutation	UNP P20813
A	29	LYS	ARG	engineered mutation	UNP P20813
A	226	HIS	TYR	engineered mutation	UNP P20813
A	262	ARG	LYS	engineered mutation	UNP P20813
A	492	HIS	-	expression tag	UNP P20813
A	493	HIS	-	expression tag	UNP P20813
A	494	HIS	-	expression tag	UNP P20813
A	495	HIS	-	expression tag	UNP P20813

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5-CYCLOHEXYL-1-PENTYL-BETA-D-MALTOSIDE (CCD ID: CM5) (formula: C₂₃H₄₂O₁₁).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			34	23	11		
3	A	1	Total	C	O	0	0
			12	11	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	11	1		

- Molecule 4 is (1S,5R)-2-(bromomethyl)-6,6-dimethylbicyclo[3.1.1]hept-2-ene (CCD ID: 85D) (formula: C₁₀H₁₅Br).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Br	C	0	1
			22	2	20		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome P450 2B6



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.25Å 77.25Å 202.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.55 – 2.27 47.55 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.55-2.27) 99.7 (47.55-2.27)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.18 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.174 , 0.229 0.184 , 0.226	Depositor DCC
R_{free} test set	1678 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	47.5	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 26.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3964	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, 85D, CM5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	1/3817 (0.0%)	1.09	7/5168 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	268	TYR	CA-C	5.38	1.59	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	461	SER	CA-C-N	5.52	125.61	119.87
1	A	461	SER	C-N-CA	5.52	125.61	119.87
1	A	358	ARG	N-CA-C	-5.41	105.30	111.14
1	A	179	ILE	CB-CA-C	-5.34	104.86	112.22
1	A	375	THR	N-CA-C	5.25	117.23	108.99
1	A	481	PRO	CA-C-N	-5.22	115.20	120.21
1	A	481	PRO	C-N-CA	-5.22	115.20	120.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3716	0	3669	34	0
2	A	43	0	30	3	0
3	A	58	0	82	11	0
4	A	22	0	0	2	0
5	A	125	0	0	1	0
All	All	3964	0	3781	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:502:CM5:O22	3:A:502:CM5:C18	1.64	1.42
2:A:501:HEM:HBB2	2:A:501:HEM:HMB2	1.58	0.86
3:A:502:CM5:O22	3:A:502:CM5:C13	2.28	0.81
3:A:502:CM5:O34	3:A:502:CM5:H17	1.83	0.78
3:A:502:CM5:O34	3:A:502:CM5:C16	2.25	0.74
3:A:502:CM5:O34	3:A:502:CM5:C17	2.38	0.71
1:A:293:LEU:C	1:A:293:LEU:HD13	2.16	0.71
1:A:293:LEU:HD13	1:A:293:LEU:O	1.98	0.62
1:A:374:HIS:C	1:A:374:HIS:CD2	2.78	0.62
4:A:505[C]:85D:C8	4:A:505[C]:85D:C3	2.75	0.62
4:A:505[A]:85D:C3	4:A:505[A]:85D:C8	2.75	0.60
1:A:140:ARG:HH11	1:A:148:GLU:CD	2.10	0.59
1:A:73:ARG:NH1	1:A:387:GLU:OE1	2.35	0.59
1:A:198:MET:HG3	3:A:502:CM5:H52	1.87	0.57
1:A:111:TYR:HB2	1:A:290:LEU:HD12	1.87	0.57
3:A:502:CM5:C18	3:A:502:CM5:H22	2.08	0.57
1:A:252:HIS:C	1:A:254:GLU:H	2.13	0.56
1:A:28:GLY:O	1:A:381:ILE:HD13	2.05	0.55
1:A:304:SER:O	1:A:308:ARG:HG3	2.08	0.54
2:A:501:HEM:HBB2	2:A:501:HEM:CMB	2.36	0.54
1:A:293:LEU:C	1:A:293:LEU:CD1	2.81	0.53
1:A:296:PHE:CE1	3:A:502:CM5:H101	2.44	0.52
1:A:360:SER:O	1:A:361:ASP:C	2.52	0.51
1:A:299:GLY:HA2	2:A:501:HEM:HMC2	1.92	0.51
1:A:297:PHE:HE1	1:A:301:GLU:HG3	1.76	0.51
1:A:135:PHE:O	1:A:139:LYS:HG3	2.11	0.50
1:A:48:ARG:HA	3:A:504:CM5:H51	1.94	0.50
1:A:252:HIS:C	1:A:254:GLU:N	2.72	0.48
1:A:212:VAL:HG23	5:A:652:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ARG:NH2	1:A:218:GLU:HG3	2.30	0.47
3:A:502:CM5:C13	3:A:502:CM5:H22	2.25	0.46
1:A:144:GLU:O	1:A:148:GLU:HG3	2.16	0.46
1:A:441:ILE:O	1:A:445:GLU:HG3	2.16	0.46
1:A:370:ILE:O	1:A:370:ILE:HG23	2.16	0.45
1:A:398:ASP:OD1	1:A:399:PRO:HD2	2.17	0.44
1:A:180:CYS:SG	1:A:296:PHE:HE1	2.41	0.44
1:A:459:MET:HG2	1:A:486:ILE:HD11	1.99	0.43
1:A:367:VAL:HA	1:A:368:PRO:HD3	1.84	0.43
3:A:502:CM5:O22	3:A:502:CM5:C17	2.53	0.43
1:A:175:THR:HG21	1:A:303:THR:HB	2.01	0.42
1:A:314:MET:HA	1:A:314:MET:HE2	2.01	0.42
1:A:51:LEU:HG	1:A:215:GLN:HB3	2.01	0.42
1:A:169:PHE:CZ	1:A:170:LEU:HG	2.56	0.41
1:A:435:ILE:O	1:A:436:CYS:C	2.64	0.41
1:A:153:LEU:HD22	1:A:174:ILE:HD13	2.03	0.41
1:A:461:SER:HB2	1:A:462:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/476 (97%)	449 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	394/418 (94%)	389 (99%)	5 (1%)	61 76

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	293	LEU
1	A	309	TYR
1	A	371	VAL
1	A	374	HIS
1	A	381	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	193	GLN
1	A	233	GLN
1	A	237	ASN
1	A	247	HIS
1	A	374	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	501	1	50,50,50	2.41	24 (48%)	67,82,82	1.56	13 (19%)
4	85D	A	505[C]	-	11,12,12	2.87	4 (36%)	14,19,19	1.11	1 (7%)
3	CM5	A	503	-	12,12,36	0.64	0	13,13,49	0.61	0
3	CM5	A	502	-	36,36,36	3.10	16 (44%)	49,49,49	4.37	24 (48%)
4	85D	A	505[A]	-	11,12,12	2.88	4 (36%)	14,19,19	1.11	1 (7%)
3	CM5	A	504	-	12,12,36	0.66	0	13,13,49	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1	-	4/14/54/54	-
4	85D	A	505[C]	-	-	0/1/27/27	0/3/2/2
3	CM5	A	503	-	-	4/6/14/65	0/1/1/3
3	CM5	A	502	-	-	8/17/65/65	0/3/3/3
4	85D	A	505[A]	-	-	0/1/27/27	0/3/2/2
3	CM5	A	504	-	-	5/6/14/65	0/1/1/3

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	CM5	O22-C18	8.62	1.64	1.43
2	A	501	HEM	FE-NB	-8.20	1.69	1.94
3	A	502	CM5	O12-C13	7.77	1.53	1.40
4	A	505[A]	85D	C1-C2	-7.13	1.38	1.51
4	A	505[C]	85D	C1-C2	-7.11	1.38	1.51
3	A	502	CM5	O34-C29	-6.36	1.27	1.43
2	A	501	HEM	C3D-C2D	5.65	1.49	1.36
4	A	505[A]	85D	C4-C3	-4.97	1.40	1.50
4	A	505[C]	85D	C4-C3	-4.97	1.40	1.50
3	A	502	CM5	O14-C13	4.91	1.54	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	CM5	C17-C16	4.34	1.64	1.52
2	A	501	HEM	C3C-C2C	-4.20	1.28	1.37
2	A	501	HEM	C3B-C2B	-4.10	1.28	1.37
3	A	502	CM5	C27-C26	4.05	1.61	1.53
2	A	501	HEM	C2A-C3A	-3.69	1.29	1.38
3	A	502	CM5	O25-C24	3.49	1.50	1.41
3	A	502	CM5	C16-C15	3.48	1.62	1.52
3	A	502	CM5	O31-C30	3.30	1.56	1.42
3	A	502	CM5	C30-C26	3.27	1.62	1.51
2	A	501	HEM	CHC-C1C	-3.18	1.32	1.38
3	A	502	CM5	O14-C15	3.04	1.51	1.44
3	A	502	CM5	C28-C27	3.03	1.60	1.52
2	A	501	HEM	O2D-CGD	-2.91	1.21	1.30
3	A	502	CM5	O25-C26	2.85	1.51	1.44
2	A	501	HEM	CHB-C4A	-2.83	1.33	1.39
3	A	502	CM5	C13-C18	-2.79	1.44	1.52
3	A	502	CM5	O23-C16	-2.75	1.36	1.43
2	A	501	HEM	CHC-C4B	-2.66	1.33	1.39
2	A	501	HEM	O2A-CGA	-2.66	1.22	1.30
4	A	505[A]	85D	C3-C2	2.59	1.39	1.33
2	A	501	HEM	C4D-C3D	-2.57	1.40	1.45
4	A	505[C]	85D	C3-C2	2.57	1.39	1.33
2	A	501	HEM	C1C-NC	-2.50	1.34	1.39
2	A	501	HEM	C3C-C4C	-2.47	1.41	1.46
2	A	501	HEM	C1C-C2C	-2.44	1.40	1.45
3	A	502	CM5	C19-C15	2.44	1.60	1.51
2	A	501	HEM	C3B-C4B	-2.43	1.40	1.44
2	A	501	HEM	CHA-C4D	-2.34	1.33	1.38
4	A	505[A]	85D	C6-C1	-2.27	1.51	1.57
4	A	505[C]	85D	C6-C1	-2.24	1.52	1.57
2	A	501	HEM	FE-ND	2.24	2.01	1.94
2	A	501	HEM	CHD-C4C	-2.22	1.34	1.38
2	A	501	HEM	C4A-NA	-2.19	1.35	1.39
2	A	501	HEM	CAB-C3B	2.18	1.53	1.47
2	A	501	HEM	CHA-C1A	-2.14	1.34	1.39
2	A	501	HEM	CHD-C1D	-2.13	1.34	1.39
2	A	501	HEM	CAC-C3C	2.07	1.52	1.47
2	A	501	HEM	CHB-C1B	-2.00	1.34	1.38

All (39) bond angle outliers are listed below:

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	CM5	O23-C24-C29	-15.41	70.18	108.09
3	A	502	CM5	O21-C17-C18	-12.13	81.78	110.38
3	A	502	CM5	C24-O23-C16	11.84	146.04	117.98
3	A	502	CM5	C24-O25-C26	8.44	130.20	113.72
3	A	502	CM5	O22-C18-C13	-6.20	95.30	110.08
3	A	502	CM5	O23-C16-C17	5.57	121.38	107.23
3	A	502	CM5	O34-C29-C24	5.31	122.74	110.08
3	A	502	CM5	C30-C26-C27	-5.11	100.47	113.02
3	A	502	CM5	O33-C28-C29	4.88	121.87	110.38
3	A	502	CM5	C24-C29-C28	4.51	119.50	110.01
2	A	501	HEM	C3B-C4B-NB	4.29	112.55	109.47
3	A	502	CM5	O21-C17-C16	-4.23	99.09	109.94
3	A	502	CM5	O12-C13-C18	4.17	114.61	108.27
3	A	502	CM5	C13-C18-C17	4.04	118.52	110.01
3	A	502	CM5	O14-C15-C16	3.94	117.86	109.72
3	A	502	CM5	C9-C10-C11	3.73	119.10	111.42
3	A	502	CM5	C10-C11-C6	3.61	119.67	112.08
2	A	501	HEM	C1B-NB-C4B	-3.46	101.11	105.21
2	A	501	HEM	CHC-C4B-NB	3.33	128.00	124.42
3	A	502	CM5	C13-O14-C15	3.19	119.95	113.72
3	A	502	CM5	O25-C26-C27	-2.95	104.38	109.70
2	A	501	HEM	CHC-C4B-C3B	-2.83	119.42	125.07
3	A	502	CM5	C19-C15-C16	-2.79	105.55	113.38
3	A	502	CM5	O14-C13-C18	2.67	115.86	110.37
4	A	505[A]	85D	C9-C6-C8	2.65	114.54	108.67
4	A	505[C]	85D	C9-C6-C8	2.65	114.53	108.67
3	A	502	CM5	C29-C28-C27	2.62	115.42	110.83
2	A	501	HEM	C1C-CHC-C4B	-2.54	120.63	126.02
3	A	502	CM5	C28-C27-C26	2.47	114.72	110.23
2	A	501	HEM	O2D-CGD-CBD	2.47	121.79	114.00
2	A	501	HEM	C4A-CHB-C1B	-2.46	120.47	126.25
2	A	501	HEM	O2A-CGA-CBA	2.35	121.43	114.00
2	A	501	HEM	C4A-NA-C1A	-2.30	102.07	105.82
3	A	502	CM5	O14-C15-C19	2.25	112.03	106.44
2	A	501	HEM	CHA-C4D-C3D	-2.16	121.24	125.23
2	A	501	HEM	CMD-C2D-C1D	2.16	128.41	125.03
3	A	502	CM5	C1-O12-C13	-2.16	109.99	113.68
2	A	501	HEM	CHD-C1D-ND	2.02	126.60	124.42
2	A	501	HEM	C1D-C2D-C3D	-2.02	104.86	106.98

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	CM5	O25-C26-C30-O31
3	A	502	CM5	C27-C26-C30-O31
3	A	502	CM5	C17-C16-O23-C24
3	A	502	CM5	C15-C16-O23-C24
3	A	504	CM5	C2-C3-C4-C5
3	A	504	CM5	C4-C5-C6-C7
3	A	503	CM5	C2-C3-C4-C5
3	A	503	CM5	C1-C2-C3-C4
3	A	502	CM5	O25-C24-O23-C16
3	A	504	CM5	O12-C1-C2-C3
3	A	504	CM5	C3-C4-C5-C6
3	A	504	CM5	C4-C5-C6-C11
3	A	502	CM5	C1-C2-C3-C4
3	A	502	CM5	C4-C5-C6-C7
3	A	503	CM5	C4-C5-C6-C11
2	A	501	HEM	C2C-C3C-CAC-CBC
3	A	503	CM5	C4-C5-C6-C7
3	A	502	CM5	C4-C5-C6-C11
2	A	501	HEM	C4C-C3C-CAC-CBC
2	A	501	HEM	CAA-CBA-CGA-O2A
2	A	501	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.


5 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	3	0
4	A	505[C]	85D	1	0
3	A	502	CM5	10	0
4	A	505[A]	85D	1	0
3	A	504	CM5	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	465/476 (97%)	-0.09	15 (3%)	50 52	31, 45, 66, 85	8 (1%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	136	GLY	5.0
1	A	492	HIS	3.4
1	A	138	GLY	3.2
1	A	137	MET	3.2
1	A	247	HIS	2.7
1	A	139	LYS	2.7
1	A	253	ARG	2.6
1	A	28	GLY	2.4
1	A	400	HIS	2.3
1	A	39	LEU	2.2
1	A	134	ASP	2.2
1	A	29	LYS	2.2
1	A	246	GLY	2.2
1	A	135	PHE	2.1
1	A	140	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CM5	A	502	34/34	0.72	0.20	59,96,114,115	0
4	85D	A	505[A]	11/11	0.85	0.21	44,50,91,96	11
4	85D	A	505[C]	11/11	0.85	0.21	37,51,87,100	11
3	CM5	A	504	12/34	0.86	0.26	83,85,91,92	0
3	CM5	A	503	12/34	0.94	0.16	83,84,86,87	0
2	HEM	A	501	43/43	0.98	0.06	27,34,42,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around CM5 A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CM5 A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.