



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:17 PM UTC

PDB ID : 5UEM / pdb_00005uem
Title : Crystal structure of 354NC37 Fab in complex with HIV-1 clade AE strain 93TH057 gp120
Authors : Sievers, S.A.; Gristick, H.B.; Bjorkman, P.J.
Deposited on : 2017-01-02
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	361	
2	H	230	
3	L	213	
4	A	2	
5	B	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	C	7	 57% 43%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	342	Total	C	N	O	S	0	0	1
			2637	1658	456	499	24			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	65	CYS	VAL	conflict	UNP A0A0M3KKW9
G	115	CYS	SER	conflict	UNP A0A0M3KKW9
G	493	GLY	-	expression tag	UNP A0A0M3KKW9
G	494	SER	-	expression tag	UNP A0A0M3KKW9
G	495	HIS	-	expression tag	UNP A0A0M3KKW9
G	496	HIS	-	expression tag	UNP A0A0M3KKW9
G	497	HIS	-	expression tag	UNP A0A0M3KKW9
G	498	HIS	-	expression tag	UNP A0A0M3KKW9
G	499	HIS	-	expression tag	UNP A0A0M3KKW9
G	500	HIS	-	expression tag	UNP A0A0M3KKW9

- Molecule 2 is a protein called 354NC37 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	230	Total	C	N	O	S	0	0	0
			1719	1084	296	331	8			

- Molecule 3 is a protein called 354NC37 Fab Light Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1599	1007	274	314	4			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	A	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	B	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	C	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	G	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	G	31	Total	O	0	0
			31	31		
9	H	22	Total	O	0	0
			22	22		
9	L	17	Total	O	0	0
			17	17		

- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  80% 20%



- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  57% 43%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.63Å 67.51Å 210.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.74 – 2.70 35.74 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (35.74-2.70) 99.6 (35.74-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.68Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.220 , 0.266 0.220 , 0.264	Depositor DCC
R_{free} test set	1432 reflections (3.97%)	wwPDB-VP
Wilson B-factor (Å ²)	42.3	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6296	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, NAG, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.11	0/2693	0.30	0/3660
2	H	0.13	0/1765	0.33	0/2409
3	L	0.11	0/1638	0.33	0/2233
All	All	0.12	0/6096	0.32	0/8302

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2637	0	2535	22	0
2	H	1719	0	1661	22	0
3	L	1599	0	1541	13	0
4	A	28	0	25	0	0
5	B	61	0	52	0	0
6	C	83	0	70	0	0
7	G	84	0	78	1	0
8	G	15	0	17	0	0
9	G	31	0	0	0	0
9	H	22	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	L	17	0	0	0	0
All	All	6296	0	5979	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.71	0.73
2:H:50:LEU:HD21	2:H:58:MET:HE3	1.75	0.69
1:G:124:GLY:H	2:H:74:HIS:HA	1.59	0.67
2:H:127:SER:H	2:H:130:SER:HB3	1.59	0.66
3:L:145:LYS:HB3	3:L:197:THR:HB	1.78	0.65
1:G:275:GLU:OE2	1:G:282:LYS:NZ	2.34	0.60
2:H:168:ALA:HB2	2:H:178:LEU:HD23	1.84	0.57
1:G:219:THR:HG21	1:G:224:VAL:HA	1.89	0.55
2:H:47:TRP:HZ2	2:H:50:LEU:HD22	1.72	0.54
1:G:270:ILE:HD13	1:G:345:VAL:HG12	1.90	0.53
1:G:379:ARG:HG3	1:G:443:ILE:HG23	1.89	0.53
1:G:381:GLU:HG3	1:G:443:ILE:HD13	1.92	0.52
1:G:207:LYS:HE2	1:G:436:ALA:HB3	1.93	0.52
1:G:367:GLY:HA3	1:G:371:ILE:HD11	1.92	0.52
1:G:66:HIS:CD2	1:G:111:LEU:HD21	2.45	0.51
1:G:446:VAL:HG21	7:G:617:NAG:H82	1.93	0.51
2:H:52(A):VAL:HA	2:H:71:ARG:HD3	1.93	0.50
2:H:130:SER:OG	2:H:137:ALA:O	2.28	0.49
3:L:90:ASN:HD21	3:L:93:GLY:H	1.60	0.49
3:L:90:ASN:HD21	3:L:93:GLY:N	2.11	0.49
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.94	0.48
3:L:54:ARG:HD3	3:L:62:PHE:O	2.14	0.47
3:L:149:LYS:HG2	3:L:154:LEU:HD23	1.96	0.47
1:G:258:GLN:HA	1:G:471:GLY:O	2.15	0.47
2:H:100(B):VAL:HG22	2:H:100(E):ARG:HB2	1.97	0.46
3:L:142:ARG:NH1	3:L:163:VAL:HG21	2.30	0.46
1:G:297:THR:OG1	1:G:444:ASN:ND2	2.45	0.46
1:G:368:ASP:CG	2:H:71:ARG:HH22	2.24	0.46
1:G:93:PHE:HB2	1:G:233:PHE:HZ	1.81	0.46
3:L:35:TRP:HB2	3:L:48:ILE:HB	1.97	0.46
2:H:18:VAL:HG21	2:H:109:ILE:HD12	1.99	0.44
2:H:50:LEU:N	2:H:69:MET:HE1	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.00	0.44
1:G:273:ARG:HB2	1:G:285:ILE:HB	1.99	0.44
3:L:94:GLY:HA3	3:L:95:THR:HA	1.77	0.44
1:G:476:LYS:O	1:G:480:ARG:HG3	2.17	0.44
2:H:2:VAL:HG12	2:H:102:VAL:HG11	1.99	0.44
2:H:4:LEU:HD23	2:H:24:ALA:HA	1.98	0.44
2:H:196:CYS:SG	2:H:209:LYS:HB3	2.58	0.44
2:H:72:ASP:HB3	2:H:75:THR:HG22	1.99	0.43
2:H:51:ILE:HD11	2:H:69:MET:HB3	2.01	0.43
2:H:100(A):PRO:HB3	2:H:100(G):TYR:CE1	2.53	0.43
1:G:477:ASP:OD1	1:G:480:ARG:NH1	2.51	0.43
1:G:100:MET:HE2	1:G:483:LEU:HD22	2.00	0.43
1:G:257:THR:O	1:G:259:LEU:N	2.48	0.42
1:G:212:PRO:HB3	1:G:253:PRO:HD2	2.01	0.42
2:H:33:TYR:HA	2:H:52(A):VAL:HG13	2.02	0.42
2:H:66:ARG:HD2	2:H:82(A):GLN:O	2.20	0.41
2:H:146:PHE:HA	2:H:147:PRO:HA	1.83	0.41
3:L:158:ASN:OD1	3:L:158:ASN:N	2.54	0.41
3:L:19:ALA:HB3	3:L:75:ILE:HB	2.02	0.41
2:H:154:TRP:CH2	2:H:196:CYS:HB3	2.57	0.40
1:G:390:LEU:HD13	1:G:416:LEU:HD21	2.03	0.40
3:L:142:ARG:CZ	3:L:163:VAL:HG21	2.52	0.40
3:L:163:VAL:HG22	3:L:175:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	336/361 (93%)	324 (96%)	12 (4%)	0	100	100
2	H	228/230 (99%)	217 (95%)	11 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	L	211/213 (99%)	201 (95%)	10 (5%)	0	100	100
All	All	775/804 (96%)	742 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	293/318 (92%)	292 (100%)	1 (0%)	86	94
2	H	186/190 (98%)	181 (97%)	5 (3%)	39	69
3	L	175/184 (95%)	172 (98%)	3 (2%)	53	79
All	All	654/692 (94%)	645 (99%)	9 (1%)	59	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	390	LEU
2	H	64	GLN
2	H	71	ARG
2	H	169	VAL
2	H	183	THR
2	H	193	THR
3	L	5	THR
3	L	24	ARG
3	L	28	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	94	ASN
1	G	425	ASN
1	G	444	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	37	GLN
3	L	90	ASN
3	L	124	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1	4,1	14,14,15	0.31	0	17,19,21	0.73	0
4	NAG	A	2	4	14,14,15	0.28	0	17,19,21	0.61	0
5	NAG	B	1	1,5	14,14,15	0.32	0	17,19,21	0.79	0
5	NAG	B	2	5	14,14,15	0.30	0	17,19,21	0.66	0
5	BMA	B	3	5	11,11,12	0.29	0	15,15,17	1.01	1 (6%)
5	MAN	B	4	5	11,11,12	0.26	0	15,15,17	0.73	0
5	MAN	B	5	5	11,11,12	0.27	0	15,15,17	0.59	0
6	NAG	C	1	6,1	14,14,15	0.31	0	17,19,21	0.81	1 (5%)
6	NAG	C	2	6	14,14,15	0.32	0	17,19,21	0.94	2 (11%)
6	BMA	C	3	6	11,11,12	0.23	0	15,15,17	0.75	0
6	MAN	C	4	6	11,11,12	0.26	0	15,15,17	0.68	0
6	MAN	C	5	6	11,11,12	0.26	0	15,15,17	0.77	0
6	MAN	C	6	6	11,11,12	0.29	0	15,15,17	1.10	1 (6%)
6	MAN	C	7	6	11,11,12	0.33	0	15,15,17	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	A	2	4	-	2/6/23/26	0/1/1/1
5	NAG	B	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	B	2	5	-	0/6/23/26	0/1/1/1
5	BMA	B	3	5	-	1/2/19/22	0/1/1/1
5	MAN	B	4	5	-	2/2/19/22	0/1/1/1
5	MAN	B	5	5	-	1/2/19/22	0/1/1/1
6	NAG	C	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	C	2	6	-	0/6/23/26	0/1/1/1
6	BMA	C	3	6	-	1/2/19/22	0/1/1/1
6	MAN	C	4	6	-	0/2/19/22	0/1/1/1
6	MAN	C	5	6	-	2/2/19/22	0/1/1/1
6	MAN	C	6	6	-	1/2/19/22	0/1/1/1
6	MAN	C	7	6	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	6	MAN	O3-C3-C2	-3.07	103.78	110.05
5	B	3	BMA	O3-C3-C2	-2.36	105.24	110.05
6	C	1	NAG	C1-O5-C5	2.26	115.21	112.19
6	C	2	NAG	O5-C1-C2	-2.24	107.82	111.29
6	C	2	NAG	O4-C4-C3	-2.06	105.52	110.38

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2	NAG	C8-C7-N2-C2
4	A	2	NAG	O7-C7-N2-C2
5	B	4	MAN	O5-C5-C6-O6
5	B	4	MAN	C4-C5-C6-O6
6	C	5	MAN	O5-C5-C6-O6
6	C	3	BMA	O5-C5-C6-O6
6	C	6	MAN	O5-C5-C6-O6
5	B	5	MAN	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	C	7	MAN	O5-C5-C6-O6
5	B	1	NAG	O5-C5-C6-O6
6	C	5	MAN	C4-C5-C6-O6
5	B	3	BMA	C4-C5-C6-O6
5	B	1	NAG	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EPE	G	621	-	15,15,15	0.75	1 (6%)	19,20,20	1.81	3 (15%)
7	NAG	G	619	1	14,14,15	0.31	0	17,19,21	0.73	0
7	NAG	G	603	1	14,14,15	0.30	0	17,19,21	0.61	0
7	NAG	G	609	1	14,14,15	0.31	0	17,19,21	0.80	0
7	NAG	G	620	1	14,14,15	0.26	0	17,19,21	0.81	0
7	NAG	G	618	1	14,14,15	0.31	0	17,19,21	0.80	1 (5%)
7	NAG	G	617	1	14,14,15	0.33	0	17,19,21	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EPE	G	621	-	-	6/9/19/19	0/1/1/1
7	NAG	G	619	1	-	0/6/23/26	0/1/1/1
7	NAG	G	603	1	-	0/6/23/26	0/1/1/1
7	NAG	G	609	1	-	2/6/23/26	0/1/1/1
7	NAG	G	620	1	-	0/6/23/26	0/1/1/1
7	NAG	G	618	1	-	1/6/23/26	0/1/1/1
7	NAG	G	617	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	G	621	EPE	C10-S	2.41	1.81	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	G	621	EPE	C5-N4-C3	4.44	118.41	108.84
8	G	621	EPE	C7-N4-C5	3.64	120.94	111.24
8	G	621	EPE	C7-N4-C3	3.23	119.84	111.24
7	G	618	NAG	C1-O5-C5	2.06	114.95	112.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	G	621	EPE	C8-C7-N4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	G	621	EPE	C9-C10-S-O1S
8	G	621	EPE	C9-C10-S-O2S
8	G	621	EPE	C9-C10-S-O3S
7	G	609	NAG	O5-C5-C6-O6
7	G	609	NAG	C4-C5-C6-O6
8	G	621	EPE	N4-C7-C8-O8
8	G	621	EPE	C8-C7-N4-C3
7	G	618	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	G	617	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	342/361 (94%)	0.33	16 (4%) 36 33	28, 38, 59, 76	0
2	H	230/230 (100%)	0.24	5 (2%) 62 59	23, 36, 59, 71	0
3	L	213/213 (100%)	0.21	5 (2%) 61 58	22, 36, 53, 60	0
All	All	785/804 (97%)	0.27	26 (3%) 49 45	22, 37, 58, 76	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	460	ALA	5.0
1	G	124	GLY	4.5
1	G	462	ASN	3.8
1	G	301	ASN	3.5
1	G	411	ASN	3.1
1	G	463	THR	3.0
1	G	198	GLY	3.0
1	G	44	VAL	2.9
2	H	129	LYS	2.6
2	H	212	GLU	2.5
1	G	461	ASN	2.4
1	G	409	GLY	2.4
3	L	94	GLY	2.4
2	H	128	SER	2.4
2	H	133	GLY	2.3
1	G	465	ASN	2.3
1	G	59	LYS	2.3
1	G	241	ASN	2.3
3	L	156	SER	2.2
3	L	12	SER	2.2
1	G	85	HIS	2.2
1	G	82	GLN	2.1
1	G	199	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	69	THR	2.1
3	L	165	GLU	2.0
2	H	132	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	1	14/15	-	-	23,50,61,71	0
4	NAG	A	2	14/15	-	-	49,59,68,72	14
5	NAG	B	1	14/15	-	-	32,42,60,65	0
5	NAG	B	2	14/15	-	-	42,58,79,84	0
5	BMA	B	3	11/12	-	-	53,70,85,87	0
5	MAN	B	4	11/12	-	-	66,66,66,66	0
5	MAN	B	5	11/12	-	-	94,94,94,94	0
6	NAG	C	1	14/15	-	-	37,50,57,62	0
6	NAG	C	2	14/15	-	-	39,50,73,77	14
6	BMA	C	3	11/12	-	-	32,49,56,62	11
6	MAN	C	4	11/12	-	-	51,51,51,51	0
6	MAN	C	5	11/12	-	-	46,46,46,46	0
6	MAN	C	6	11/12	-	-	56,56,56,56	0
6	MAN	C	7	11/12	-	-	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain A:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain B:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NAG	G	603	14/15	0.21	0.20	66,66,66,66	0
7	NAG	G	619	14/15	0.72	0.16	54,60,65,67	14
7	NAG	G	620	14/15	0.72	0.19	40,45,50,52	14
7	NAG	G	617	14/15	0.78	0.19	43,59,74,75	0
7	NAG	G	609	14/15	0.86	0.11	32,37,49,50	0
7	NAG	G	618	14/15	0.87	0.11	35,54,75,82	0
8	EPE	G	621	15/15	0.96	0.10	25,30,45,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.