



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:48 PM UTC

PDB ID : 7UE1 / pdb_00007ue1
Title : HIV-1 Integrase Catalytic Core Domain Mutant (KGD) in Complex with Inhibitor GRL-142
Authors : Aoki, M.; Aoki-Ogata, H.; Bulut, H.; Hayashi, H.; Davis, D.; Hasegawa, K.; Yarchoan, R.; Ghosh, A.K.; Pau, A.K.; Mitsuya, H.
Deposited on : 2022-03-21
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	
1	B	166	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	302	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4545 atoms, of which 2268 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	140	Total	C	H	N	O	S	0	0	0
			2150	683	1076	189	198	4			
1	B	148	Total	C	H	N	O	S	0	0	0
			2302	732	1152	202	212	4			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLY	-	expression tag	UNP Q9YUI7
A	48	SER	-	expression tag	UNP Q9YUI7
A	49	HIS	-	expression tag	UNP Q9YUI7
A	74	ILE	LEU	conflict	UNP Q9YUI7
A	121	VAL	PHE	conflict	UNP Q9YUI7
A	125	ALA	THR	conflict	UNP Q9YUI7
A	148	ARG	GLN	conflict	UNP Q9YUI7
A	151	VAL	ILE	conflict	UNP Q9YUI7
A	185	HIS	PHE	conflict	UNP Q9YUI7
A	208	LEU	ILE	conflict	UNP Q9YUI7
B	47	GLY	-	expression tag	UNP Q9YUI7
B	48	SER	-	expression tag	UNP Q9YUI7
B	49	HIS	-	expression tag	UNP Q9YUI7
B	74	ILE	LEU	conflict	UNP Q9YUI7
B	121	VAL	PHE	conflict	UNP Q9YUI7
B	125	ALA	THR	conflict	UNP Q9YUI7
B	148	ARG	GLN	conflict	UNP Q9YUI7
B	151	VAL	ILE	conflict	UNP Q9YUI7
B	185	HIS	PHE	conflict	UNP Q9YUI7
B	208	LEU	ILE	conflict	UNP Q9YUI7

- Molecule 2 is (3S,3aR,5R,7aS,8S)-hexahydro-4H-3,5-methanofuro[2,3-b]pyran-8-yl [(2S,3R)-4-[[2-(cyclopropylamino)-1,3-benzothiazol-6-yl]sulfonyl](2-methylpropyl)amino]-1-(3,5-difluorophenyl)-3-hydroxybutan-2-yl]carbamate (CCD ID: 7OA) (formula: C₃₃H₄₀F₂N₄O₇S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	H	N	O		
2	B	1	88	33	2	40	4	7	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	B	1	5	4	1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Integrase



• Molecule 1: Integrase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	111.37Å 111.37Å 137.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	86.49 – 3.00 86.49 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (86.49-3.00) 100.0 (86.49-3.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.223 , 0.317 0.240 , 0.321	Depositor DCC
R_{free} test set	420 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	88.8	Xtriage
Anisotropy	0.598	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 90.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4545	wwPDB-VP
Average B, all atoms (Å ²)	112.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 7OA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/1094	0.94	0/1481
1	B	0.82	0/1174	1.23	0/1593
All	All	0.79	0/2268	1.10	0/3074

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1074	1076	1085	38	1
1	B	1150	1152	1152	30	3
2	B	48	40	0	4	1
3	B	5	0	0	0	11
All	All	2277	2268	2237	66	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:301:7OA:C7	2:B:301:7OA:O1	1.71	1.38
2:B:301:7OA:O1	2:B:301:7OA:C8	1.87	1.22
1:A:84:ILE:HD11	1:A:154:MET:HG3	1.65	0.78
1:A:195:SER:OG	1:A:198:GLU:HG3	1.84	0.77
1:A:169:ALA:HB2	1:A:178:MET:CE	2.15	0.76
1:A:118:GLY:O	1:A:122:THR:HG23	1.88	0.73
1:A:84:ILE:HD11	1:A:154:MET:CG	2.23	0.68
1:B:116:ASP:HB3	1:B:141:ILE:HD11	1.76	0.67
1:A:84:ILE:HD13	1:A:158:LEU:HD21	1.77	0.66
1:A:202:ASP:O	1:A:206:THR:HG23	1.96	0.65
1:B:173:LYS:O	1:B:176:VAL:HG22	1.96	0.65
1:A:169:ALA:HB2	1:A:178:MET:HE1	1.78	0.64
1:A:112:THR:HG23	1:A:136:LYS:O	1.99	0.63
1:A:57:SER:CB	1:A:60:ILE:HD13	2.28	0.62
1:A:57:SER:HB2	1:A:60:ILE:HD13	1.81	0.62
1:A:172:LEU:O	1:A:176:VAL:HG13	1.98	0.62
1:A:176:VAL:O	1:A:180:VAL:HG23	2.01	0.61
1:A:158:LEU:O	1:A:162:ILE:HG13	2.01	0.60
1:B:72:VAL:HG21	1:B:91:ALA:HA	1.84	0.59
1:B:101:LEU:HD11	1:B:113:VAL:HG21	1.83	0.58
1:A:78:HIS:CG	1:A:200:ILE:HD12	2.39	0.57
1:A:201:VAL:HG12	1:B:205:ALA:HB2	1.85	0.57
1:A:62:GLN:NE2	1:A:79:VAL:HG22	2.23	0.54
1:A:62:GLN:HE21	1:A:79:VAL:HG22	1.75	0.52
1:A:201:VAL:CG1	1:B:205:ALA:HB2	2.41	0.51
1:B:141:ILE:N	1:B:142:PRO:CD	2.74	0.50
1:A:64:ASP:OD1	1:A:65:CYS:N	2.45	0.49
1:B:128:ALA:HB1	2:B:301:7OA:C11	2.43	0.48
1:A:57:SER:HB3	1:A:60:ILE:HD13	1.95	0.48
1:A:68:LEU:HD13	1:A:159:LYS:HD3	1.96	0.48
1:B:63:LEU:HD22	1:B:101:LEU:HD13	1.94	0.48
1:B:115:THR:HB	1:B:121:VAL:HG11	1.95	0.48
1:B:116:ASP:O	1:B:117:ASN:CG	2.57	0.48
1:A:67:HIS:O	1:A:159:LYS:NZ	2.47	0.48
1:A:62:GLN:HE21	1:A:79:VAL:CG2	2.26	0.47
1:B:187:ARG:HD2	1:B:194:TYR:HA	1.95	0.47
1:A:63:LEU:HD12	1:A:101:LEU:HD13	1.95	0.47
1:B:117:ASN:OD1	1:B:117:ASN:C	2.57	0.46
1:B:141:ILE:N	1:B:142:PRO:HD3	2.29	0.46
1:A:74:ILE:O	1:A:86:ALA:HA	2.16	0.46
1:A:78:HIS:ND1	1:A:200:ILE:HD12	2.30	0.46
1:B:181:PHE:C	1:B:181:PHE:CD1	2.94	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:CD1	1:A:158:LEU:HD21	2.45	0.45
1:A:102:LEU:HD21	1:B:178:MET:HG3	1.98	0.45
1:B:77:VAL:HG22	1:B:84:ILE:HG12	1.99	0.45
1:A:121:VAL:O	1:A:137:GLN:NE2	2.51	0.44
1:B:154:MET:HA	1:B:154:MET:HE2	1.99	0.44
1:B:72:VAL:HG11	1:B:89:ILE:HD11	1.99	0.44
1:B:84:ILE:HD12	1:B:183:HIS:CD2	2.53	0.44
1:B:115:THR:HB	1:B:121:VAL:CG1	2.48	0.44
1:A:63:LEU:HD23	1:A:64:ASP:N	2.33	0.43
1:B:187:ARG:HD2	1:B:194:TYR:CA	2.48	0.43
1:B:63:LEU:HD11	1:B:74:ILE:HD11	2.00	0.43
1:A:63:LEU:HD12	1:A:101:LEU:CD1	2.49	0.43
2:B:301:7OA:C7	2:B:301:7OA:C8	2.94	0.43
1:B:182:ILE:O	1:B:186:LYS:HB2	2.19	0.42
1:B:187:ARG:NE	1:B:193:GLY:O	2.41	0.42
1:A:106:GLY:HA2	1:B:181:PHE:HA	2.00	0.42
1:B:108:TRP:CZ3	1:B:200:ILE:HG12	2.53	0.42
1:A:191:ILE:HD12	1:A:191:ILE:N	2.35	0.42
1:A:107:ARG:HB2	1:B:83:TYR:OH	2.19	0.42
1:B:200:ILE:HG23	1:B:201:VAL:N	2.34	0.42
1:A:152:GLU:O	1:A:155:ASN:HB2	2.19	0.41
1:A:84:ILE:HD13	1:A:158:LEU:CD2	2.47	0.41
1:A:200:ILE:HG23	1:A:201:VAL:N	2.35	0.40
1:B:66:THR:O	1:B:72:VAL:O	2.40	0.40

All (13) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:302:SO4:O1	3:B:302:SO4:O4[16_555]	1.06	1.14
3:B:302:SO4:O2	3:B:302:SO4:O4[16_555]	1.44	0.76
3:B:302:SO4:O2	3:B:302:SO4:O3[16_555]	1.44	0.76
3:B:302:SO4:S	3:B:302:SO4:O3[16_555]	1.45	0.75
3:B:302:SO4:S	3:B:302:SO4:O1[16_555]	1.45	0.75
3:B:302:SO4:S	3:B:302:SO4:O4[16_555]	1.46	0.74
3:B:302:SO4:S	3:B:302:SO4:O2[16_555]	1.46	0.74
3:B:302:SO4:O1	3:B:302:SO4:O1[16_555]	1.85	0.35
1:B:93:THR:OG1	3:B:302:SO4:O1[16_555]	1.89	0.31
3:B:302:SO4:O3	3:B:302:SO4:O4[16_555]	1.98	0.22
1:A:170:GLU:OE2	1:B:119:SER:OG[16_555]	2.02	0.18

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:302:SO4:O1	3:B:302:SO4:O3[16_555]	2.09	0.11
1:B:135:ILE:O	2:B:301:7OA:H26[15_455]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/166 (82%)	131 (96%)	5 (4%)	0	100	100
1	B	144/166 (87%)	138 (96%)	5 (4%)	1 (1%)	18	53
All	All	280/332 (84%)	269 (96%)	10 (4%)	1 (0%)	30	65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	73	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/135 (83%)	111 (99%)	1 (1%)	70	85
1	B	122/135 (90%)	119 (98%)	3 (2%)	42	72
All	All	234/270 (87%)	230 (98%)	4 (2%)	53	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	176	VAL
1	B	117	ASN
1	B	147	SER
1	B	171	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	120	ASN
1	A	177	GLN
1	B	78	HIS
1	B	183	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7OA	B	301	-	54,54,54	5.81	24 (44%)	73,81,81	4.37	28 (38%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	302	-	4,4,4	0.25	0	6,6,6	2.64	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7OA	B	301	-	-	11/40/72/72	1/9/7/7

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	7OA	O1-C8	25.78	1.87	1.43
2	B	301	7OA	C1-S1	-17.43	1.54	1.75
2	B	301	7OA	O1-C7	14.11	1.71	1.41
2	B	301	7OA	C39-S1	-10.51	1.53	1.74
2	B	301	7OA	C2-N1	9.16	1.58	1.39
2	B	301	7OA	S8-N11	8.06	1.74	1.63
2	B	301	7OA	C1-N2	7.96	1.46	1.34
2	B	301	7OA	C8-C9	-7.49	1.42	1.51
2	B	301	7OA	C1-N1	5.69	1.42	1.30
2	B	301	7OA	C10-C9	-5.38	1.39	1.53
2	B	301	7OA	C21-N20	5.24	1.47	1.34
2	B	301	7OA	C5-S8	5.19	1.83	1.76
2	B	301	7OA	C7-C69	-4.93	1.45	1.52
2	B	301	7OA	O23-C21	4.81	1.43	1.35
2	B	301	7OA	C59-C49	-4.01	1.43	1.53
2	B	301	7OA	C11-C59	-3.79	1.45	1.53
2	B	301	7OA	O2-C11	3.47	1.51	1.43
2	B	301	7OA	C10-C69	-3.03	1.45	1.53
2	B	301	7OA	O9-S8	2.89	1.46	1.43
2	B	301	7OA	O10-S8	2.79	1.46	1.43
2	B	301	7OA	C69-C59	-2.39	1.49	1.55
2	B	301	7OA	C36-C79	2.11	1.53	1.48
2	B	301	7OA	C9-C49	2.07	1.56	1.53
2	B	301	7OA	C29-C79	2.04	1.53	1.48

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	7OA	C39-S1-C1	20.24	107.06	88.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	7OA	S1-C1-N2	17.59	140.94	118.89
2	B	301	7OA	N2-C1-N1	-11.87	110.67	124.40
2	B	301	7OA	O9-S8-O10	-11.66	101.39	119.59
2	B	301	7OA	O23-C21-N20	7.67	120.04	110.26
2	B	301	7OA	S1-C1-N1	-7.22	108.39	117.06
2	B	301	7OA	O10-S8-N11	5.76	112.11	106.69
2	B	301	7OA	C4-C5-S8	-5.04	114.72	119.73
2	B	301	7OA	O9-S8-N11	4.90	111.30	106.69
3	B	302	SO4	O4-S-O2	4.81	134.72	109.56
2	B	301	7OA	C6-C5-S8	4.58	124.29	119.30
2	B	301	7OA	O1-C8-C9	-4.10	106.22	111.36
2	B	301	7OA	C39-C2-N1	-3.87	108.44	115.20
2	B	301	7OA	C10-C9-C49	3.48	107.26	101.97
2	B	301	7OA	O22-C21-N20	-3.47	119.17	124.86
2	B	301	7OA	O10-S8-C5	3.04	111.89	108.10
2	B	301	7OA	C35-C34-C08	-2.86	120.03	123.50
2	B	301	7OA	C29-C79-N2	-2.80	114.85	118.52
2	B	301	7OA	C32-C19-C17	-2.72	107.24	111.66
2	B	301	7OA	C79-N2-C1	-2.66	119.81	124.48
2	B	301	7OA	C35-C33-C06	-2.65	120.28	123.50
2	B	301	7OA	C34-C35-C33	2.61	119.88	116.08
2	B	301	7OA	O23-C21-O22	-2.56	120.79	124.55
2	B	301	7OA	C6-C39-C2	-2.55	120.09	121.99
3	B	302	SO4	O2-S-O1	-2.48	91.55	109.06
2	B	301	7OA	O9-S8-C5	2.47	111.17	108.10
2	B	301	7OA	C6-C39-S1	2.32	131.87	126.07
2	B	301	7OA	C07-C32-C19	-2.32	109.47	113.40
2	B	301	7OA	C16-N11-C12	2.26	122.35	115.05
3	B	302	SO4	O4-S-O3	-2.18	96.49	108.54
2	B	301	7OA	C36-C79-N2	-2.13	115.73	118.52

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	7OA	C13-C12-N11-C16
2	B	301	7OA	C36-C79-N2-C1
2	B	301	7OA	C12-N11-S8-O9
2	B	301	7OA	N1-C1-N2-C79
2	B	301	7OA	S1-C1-N2-C79
2	B	301	7OA	O22-C21-N20-C19
2	B	301	7OA	O23-C21-N20-C19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	301	7OA	C12-N11-S8-C5
2	B	301	7OA	C12-N11-S8-O10
2	B	301	7OA	C29-C79-N2-C1
2	B	301	7OA	C16-N11-S8-O9


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	7OA	C11-C49-C59-C7-C8-C9-O1-O2

2 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	7OA	4	1
3	B	302	SO4	0	11

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	140/166 (84%)	0.10	3 (2%) 63 40	80, 104, 143, 182	0
1	B	148/166 (89%)	0.19	0 100 100	75, 110, 167, 187	0
All	All	288/332 (86%)	0.15	3 (1%) 79 59	75, 108, 161, 187	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	151	VAL	2.2
1	A	66	THR	2.2
1	A	158	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	302	5/5	0.75	0.19	354,364,408,524	1
2	7OA	B	301	48/48	0.90	0.14	73,102,121,137	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7OA B 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.