



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:40 AM UTC

PDB ID : 1UFK / pdb\_00001ufk  
Title : Crystal structure of TT0836  
Authors : Kaminishi, T.; Sakai, H.; Takemoto-Hori, C.; Terada, T.; Nakagawa, N.;  
Maoka, N.; Kuramitsu, S.; Shirouzu, M.; Yokoyama, S.; RIKEN Structural  
Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2003-05-31  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TT0836 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	248	1919	1245	333	338	3	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	88	Total	O	0	0
			88	88		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TT0836 protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.86Å 76.00Å 62.13Å 90.00° 130.32° 90.00°	Depositor
Resolution (Å)	47.80 – 1.90 47.80 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.4 (47.80-1.90) 94.5 (47.80-1.90)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.74 (at 1.90Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.232 , 0.250 0.228 , 0.241	Depositor DCC
$R_{free}$ test set	1045 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtrriage
Anisotropy	0.761	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2007	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/1974	1.00	10/2691 (0.4%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	GLY	N-CA-C	-7.67	103.97	115.32
1	A	145	ALA	N-CA-C	6.58	120.41	109.95
1	A	124	VAL	N-CA-C	5.98	117.09	108.48
1	A	169	PRO	N-CA-C	-5.49	102.95	111.13
1	A	72	ALA	CA-C-N	5.48	126.03	120.38
1	A	72	ALA	C-N-CA	5.48	126.03	120.38
1	A	232	ALA	N-CA-C	-5.43	105.26	111.07
1	A	129	THR	N-CA-C	5.11	117.58	111.71
1	A	151	ASP	CA-C-N	5.05	124.66	119.05
1	A	151	ASP	C-N-CA	5.05	124.66	119.05

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1919	0	1918	49	0
2	A	88	0	0	4	0
All	All	2007	0	1918	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:GLU:H	1:A:85:GLU:CD	1.96	0.70
1:A:65:ARG:HH11	1:A:65:ARG:HG3	1.55	0.70
1:A:88:GLU:H	1:A:88:GLU:CD	1.97	0.68
1:A:220:LEU:HD13	1:A:247:TRP:CZ3	2.29	0.67
1:A:152:PRO:HG3	1:A:173:GLU:CD	2.22	0.65
1:A:168:ARG:HG3	2:A:308:HOH:O	1.97	0.64
1:A:201:ALA:HB3	1:A:202:PRO:HD3	1.82	0.61
1:A:112:LYS:HG2	2:A:271:HOH:O	2.02	0.59
1:A:176:LEU:O	1:A:180:LEU:HD22	2.04	0.58
1:A:81:TRP:CH2	1:A:95:PRO:HG2	2.39	0.57
1:A:254:ARG:HG3	1:A:254:ARG:O	2.05	0.56
1:A:226:LEU:N	1:A:226:LEU:HD12	2.21	0.56
1:A:177:GLU:HA	1:A:180:LEU:CD2	2.36	0.56
1:A:205:ARG:HD2	1:A:234:ALA:O	2.06	0.55
1:A:31:ARG:HD3	1:A:36:TRP:CZ2	2.43	0.52
1:A:14:ASP:HB3	1:A:15:PRO:HD3	1.91	0.52
1:A:175:SER:OG	1:A:176:LEU:N	2.38	0.52
1:A:123:LYS:NZ	1:A:123:LYS:HB3	2.26	0.51
1:A:172:LEU:HD23	1:A:172:LEU:C	2.36	0.49
1:A:151:ASP:O	1:A:154:VAL:HG22	2.11	0.49
1:A:226:LEU:CD1	1:A:226:LEU:H	2.25	0.49
1:A:61:GLU:O	1:A:65:ARG:HG2	2.12	0.49
1:A:153:MET:HE2	1:A:153:MET:HA	1.95	0.49
1:A:224:ALA:HB3	1:A:225:PRO:HD3	1.94	0.49
1:A:223:ARG:HG3	1:A:223:ARG:HH11	1.77	0.49
1:A:65:ARG:HG3	1:A:65:ARG:NH1	2.26	0.48
1:A:81:TRP:HH2	1:A:95:PRO:HG2	1.79	0.47
1:A:61:GLU:OE2	1:A:61:GLU:HA	2.15	0.47
1:A:226:LEU:N	1:A:226:LEU:CD1	2.79	0.46
1:A:87:ALA:N	1:A:88:GLU:OE1	2.48	0.46
1:A:222:ASP:OD2	1:A:223:ARG:NH1	2.48	0.46
1:A:254:ARG:O	1:A:254:ARG:CG	2.64	0.45
1:A:10:LEU:HG	1:A:30:GLU:HB2	1.98	0.45
1:A:237:ARG:HH11	1:A:253:GLY:HA3	1.82	0.45
1:A:159:GLU:O	1:A:163:LYS:HG3	2.16	0.45
1:A:187:LEU:HD23	1:A:187:LEU:C	2.41	0.44
1:A:85:GLU:OE2	1:A:86:GLY:N	2.50	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:THR:HG21	1:A:218:GLY:HA2	2.00	0.43
1:A:16:ILE:HB	1:A:47:TYR:CE2	2.54	0.43
1:A:205:ARG:HG3	1:A:254:ARG:NE	2.34	0.43
1:A:91:LEU:HD21	1:A:137:ALA:HB2	2.01	0.42
1:A:203:ARG:HD3	2:A:294:HOH:O	2.20	0.42
1:A:40:PRO:HD2	2:A:314:HOH:O	2.20	0.42
1:A:205:ARG:HG3	1:A:254:ARG:HD3	2.02	0.41
1:A:74:PRO:HD2	1:A:75:PHE:CD1	2.56	0.41
1:A:139:GLU:HG3	1:A:169:PRO:HG3	2.02	0.41
1:A:220:LEU:HD13	1:A:247:TRP:CE3	2.55	0.41
1:A:226:LEU:HD12	1:A:226:LEU:H	1.85	0.41
1:A:88:GLU:CD	1:A:88:GLU:N	2.72	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	244/254 (96%)	238 (98%)	6 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	186/189 (98%)	174 (94%)	12 (6%)	15 8

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	GLU
1	A	88	GLU
1	A	94	GLU
1	A	114	LEU
1	A	144	LYS
1	A	153	MET
1	A	180	LEU
1	A	200	LEU
1	A	216	LEU
1	A	220	LEU
1	A	237	ARG
1	A	248	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	248/254 (97%)	0.99	39 (15%) <b>5</b>   <b>5</b>	22, 36, 51, 66	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	103	HIS	4.6
1	A	83	THR	4.1
1	A	225	PRO	4.0
1	A	104	HIS	3.3
1	A	68	LYS	3.3
1	A	87	ALA	3.0
1	A	237	ARG	2.9
1	A	56	ASP	2.9
1	A	229	GLU	2.7
1	A	62	ALA	2.6
1	A	163	LYS	2.6
1	A	84	TRP	2.6
1	A	41	ALA	2.6
1	A	172	LEU	2.5
1	A	178	ALA	2.5
1	A	71	LEU	2.5
1	A	60	LEU	2.5
1	A	31	ARG	2.4
1	A	222	ASP	2.4
1	A	33	GLY	2.3
1	A	248	VAL	2.3
1	A	254	ARG	2.3
1	A	57	GLU	2.3
1	A	177	GLU	2.2
1	A	61	GLU	2.2
1	A	180	LEU	2.2
1	A	153	MET	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	67	LEU	2.2
1	A	226	LEU	2.2
1	A	86	GLY	2.2
1	A	45	LEU	2.1
1	A	29	TRP	2.1
1	A	102	GLY	2.1
1	A	48	GLU	2.1
1	A	232	ALA	2.1
1	A	22	ASP	2.1
1	A	65	ARG	2.1
1	A	200	LEU	2.0
1	A	164	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.