



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 01:37 AM UTC

PDB ID : 5UGL / pdb_00005ugl
Title : Crystal Structure of FGF Receptor 2 Tyrosine Kinase Domain Harboring the D650V Activating Mutation
Authors : Mohammadi, M.; Chen, H.
Deposited on : 2017-01-09
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	7% 79% 8% 12%
1	B	324	15% 77% 6% 16%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	271	Total	C	N	O	S	0	2	0
			2159	1381	363	394	21			
1	A	285	Total	C	N	O	S	0	2	0
			2292	1463	387	422	20			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	445	MET	-	initiating methionine	UNP P21802
B	446	GLY	-	expression tag	UNP P21802
B	447	SER	-	expression tag	UNP P21802
B	448	SER	-	expression tag	UNP P21802
B	449	HIS	-	expression tag	UNP P21802
B	450	HIS	-	expression tag	UNP P21802
B	451	HIS	-	expression tag	UNP P21802
B	452	HIS	-	expression tag	UNP P21802
B	453	HIS	-	expression tag	UNP P21802
B	454	HIS	-	expression tag	UNP P21802
B	455	SER	-	expression tag	UNP P21802
B	456	GLN	-	expression tag	UNP P21802
B	457	ASP	-	expression tag	UNP P21802
B	491	ALA	CYS	conflict	UNP P21802
B	650	VAL	ASP	engineered mutation	UNP P21802
A	445	MET	-	initiating methionine	UNP P21802
A	446	GLY	-	expression tag	UNP P21802
A	447	SER	-	expression tag	UNP P21802
A	448	SER	-	expression tag	UNP P21802
A	449	HIS	-	expression tag	UNP P21802
A	450	HIS	-	expression tag	UNP P21802
A	451	HIS	-	expression tag	UNP P21802
A	452	HIS	-	expression tag	UNP P21802
A	453	HIS	-	expression tag	UNP P21802
A	454	HIS	-	expression tag	UNP P21802

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	455	SER	-	expression tag	UNP P21802
A	456	GLN	-	expression tag	UNP P21802
A	457	ASP	-	expression tag	UNP P21802
A	491	ALA	CYS	conflict	UNP P21802
A	650	VAL	ASP	engineered mutation	UNP P21802

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
3	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	87	Total	O	0	0
			87	87		
4	A	95	Total	O	0	0
			95	95		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibroblast growth factor receptor 2



• Molecule 1: Fibroblast growth factor receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.22Å 78.79Å 116.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.84 – 1.86 34.84 – 1.86	Depositor EDS
% Data completeness (in resolution range)	87.9 (34.84-1.86) 87.9 (34.84-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	17.78 (at 1.87Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.219 , 0.258 0.220 , 0.259	Depositor DCC
R_{free} test set	2000 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.581	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4705	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7206e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2343	0.51	0/3166
1	B	0.30	0/2207	0.50	0/2982
All	All	0.31	0/4550	0.50	0/6148

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2292	0	2302	20	1
1	B	2159	0	2166	13	1
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	31	0	13	1	0
3	B	31	0	13	0	0
4	A	95	0	0	2	1
4	B	87	0	0	3	1
All	All	4705	0	4494	30	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:644[B]:ASP:OD2	3:A:802:ANP:O1G	2.07	0.72
1:A:644[B]:ASP:OD1	4:A:901:HOH:O	2.13	0.66
1:A:630:ARG:NH1	4:A:902:HOH:O	2.29	0.59
1:B:508:PRO:O	4:B:901:HOH:O	2.17	0.58
1:B:521:ASP:OD2	1:A:702:SER:OG	2.22	0.56
1:B:724:LYS:HD3	1:B:725:PRO:O	2.07	0.55
1:A:656:TYR:HD2	1:A:680[B]:TYR:HD2	1.56	0.52
1:A:651:ILE:HG21	1:A:656:TYR:HB2	1.92	0.51
1:A:573:ARG:HE	1:A:577:ARG:NH1	2.09	0.51
1:A:756:ASP:OD1	1:A:759:ARG:NH2	2.44	0.51
1:B:521:ASP:CG	1:A:702:SER:HG	2.21	0.47
1:A:680[A]:TYR:CE1	1:A:684:SER:HB2	2.50	0.47
1:B:561:TYR:OH	4:B:902:HOH:O	2.17	0.46
1:B:548:ILE:HG13	1:B:633:LEU:HD12	1.97	0.46
1:A:579:ARG:HE	1:A:596:GLU:CD	2.24	0.45
1:A:548:ILE:HD12	1:A:641:LYS:HD3	1.98	0.45
1:B:625:ARG:HA	1:B:680[A]:TYR:CZ	2.52	0.44
1:B:499:GLU:HG2	1:B:511:ALA:HB1	1.99	0.44
1:B:541:ILE:HD11	4:B:946:HOH:O	2.18	0.44
1:A:687:TRP:CE3	1:A:740:TRP:HA	2.53	0.44
1:B:724:LYS:HG3	1:B:733:TYR:HB2	2.00	0.43
1:A:520:LYS:HE3	1:A:520:LYS:HB2	1.72	0.42
1:A:759:ARG:O	1:A:763:LEU:HD23	2.18	0.42
1:B:519:LEU:HD11	1:B:528:LEU:HD13	2.00	0.42
1:A:630:ARG:HH21	1:A:666:PRO:HG2	1.84	0.42
1:B:698:THR:HG22	1:B:725:PRO:HB3	2.02	0.42
1:A:652:ASN:HB3	1:A:653:ASN:H	1.59	0.42
1:B:708:PRO:HB3	1:A:559:PRO:HB3	2.02	0.41
1:A:625:ARG:HA	1:A:680[A]:TYR:CZ	2.56	0.41
1:A:674:ALA:HB1	1:A:680[B]:TYR:CE1	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:506:ASP:OD1	1:A:612:ARG:NH1[1_655]	2.05	0.15
4:B:918:HOH:O	4:A:978:HOH:O[1_655]	2.07	0.13

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/324 (87%)	276 (98%)	7 (2%)	0	100	100
1	B	267/324 (82%)	262 (98%)	5 (2%)	0	100	100
All	All	550/648 (85%)	538 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/284 (88%)	245 (98%)	5 (2%)	48	36
1	B	234/284 (82%)	230 (98%)	4 (2%)	53	42
All	All	484/568 (85%)	475 (98%)	9 (2%)	50	38

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	699	LEU
1	B	724	LYS
1	B	727	ASN
1	B	734	MET
1	A	470	GLU
1	A	543	LYS
1	A	548	ILE
1	A	652	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	659	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	494	GLN
1	B	683	GLN
1	A	620	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ANP	B	802	-	33,33,33	2.43	10 (30%)	45,52,52	1.84	13 (28%)
2	SO4	A	801	-	4,4,4	0.29	0	6,6,6	0.14	0
2	SO4	B	801	-	4,4,4	0.25	0	6,6,6	0.15	0
3	ANP	A	802	-	33,33,33	2.50	8 (24%)	45,52,52	1.97	12 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	B	802	-	-	5/18/38/38	0/3/3/3
3	ANP	A	802	-	-	6/18/38/38	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	ANP	PB-O1B	8.02	1.58	1.46
3	B	802	ANP	PB-O1B	7.96	1.58	1.46
3	A	802	ANP	PG-O1G	7.07	1.56	1.46
3	B	802	ANP	PG-O1G	5.98	1.55	1.46
3	B	802	ANP	C6-N6	4.39	1.45	1.34
3	A	802	ANP	C6-N6	4.25	1.45	1.34
3	A	802	ANP	C2'-C3'	-3.79	1.43	1.53
3	B	802	ANP	C2'-C3'	-3.51	1.43	1.53
3	A	802	ANP	C2'-C1'	-2.96	1.44	1.53
3	A	802	ANP	C3'-C4'	-2.89	1.45	1.53
3	B	802	ANP	C2'-C1'	-2.64	1.45	1.53
3	B	802	ANP	C3'-C4'	-2.54	1.46	1.53
3	B	802	ANP	O4'-C1'	2.48	1.47	1.42
3	A	802	ANP	O4'-C1'	2.42	1.47	1.42
3	B	802	ANP	PG-O3G	-2.39	1.50	1.56
3	B	802	ANP	PB-O3A	2.36	1.62	1.59
3	B	802	ANP	PB-N3B	2.20	1.69	1.63
3	A	802	ANP	PG-O3G	-2.16	1.51	1.56

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	ANP	N3-C2-N1	-5.34	120.50	128.58
3	B	802	ANP	N3-C2-N1	-5.28	120.59	128.58
3	A	802	ANP	C5-C4-N3	-4.78	120.13	126.72
3	B	802	ANP	C5-C4-N3	-4.32	120.77	126.72
3	A	802	ANP	N3-C4-N9	4.10	134.13	127.17
3	B	802	ANP	N3-C4-N9	3.84	133.71	127.17
3	A	802	ANP	C5-N7-C8	3.38	108.76	103.45
3	A	802	ANP	C2-N3-C4	3.30	119.89	111.83
3	B	802	ANP	C5-N7-C8	3.10	108.33	103.45
3	A	802	ANP	C3'-C2'-C1'	3.02	107.17	101.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	ANP	C2-N3-C4	2.99	119.14	111.83
3	B	802	ANP	N9-C8-N7	-2.67	110.15	113.94
3	A	802	ANP	C4-C5-N7	-2.63	107.57	110.58
3	B	802	ANP	C2'-C1'-N9	-2.61	106.81	113.30
3	A	802	ANP	N9-C8-N7	-2.59	110.25	113.94
3	A	802	ANP	O4'-C1'-N9	2.57	113.03	108.09
3	A	802	ANP	O1B-PB-N3B	-2.53	108.04	111.77
3	B	802	ANP	O4'-C1'-N9	2.51	112.92	108.09
3	B	802	ANP	O2G-PG-O1G	-2.38	107.47	113.45
3	A	802	ANP	C2'-C1'-N9	-2.38	107.40	113.30
3	B	802	ANP	C4-C5-N7	-2.37	107.87	110.58
3	A	802	ANP	C2'-C3'-C4'	2.35	107.15	102.61
3	B	802	ANP	C4'-O4'-C1'	-2.31	104.37	109.47
3	B	802	ANP	C4-N9-C8	2.27	108.12	105.74
3	B	802	ANP	C2-N1-C6	2.22	122.37	118.73

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	802	ANP	PG-N3B-PB-O1B
3	B	802	ANP	PG-N3B-PB-O3A
3	B	802	ANP	PA-O3A-PB-O2B
3	B	802	ANP	C5'-O5'-PA-O2A
3	B	802	ANP	C5'-O5'-PA-O3A
3	A	802	ANP	PB-N3B-PG-O1G
3	A	802	ANP	PG-N3B-PB-O1B
3	A	802	ANP	C5'-O5'-PA-O3A
3	A	802	ANP	O4'-C4'-C5'-O5'
3	A	802	ANP	C3'-C4'-C5'-O5'
3	A	802	ANP	C5'-O5'-PA-O2A


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/324 (87%)	0.48	23 (8%) 18 19	7, 21, 51, 85	2 (0%)
1	B	271/324 (83%)	0.87	49 (18%) 3 3	9, 25, 57, 87	2 (0%)
All	All	556/648 (85%)	0.67	72 (12%) 7 7	7, 23, 56, 87	4 (0%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	490	GLY	6.5
1	B	581	PRO	6.0
1	B	491	ALA	5.7
1	B	763	LEU	5.5
1	A	660	THR	5.1
1	A	661	THR	4.9
1	A	491	ALA	4.8
1	A	492	PHE	4.8
1	B	490	GLY	4.3
1	B	720	HIS	4.3
1	B	764	THR	4.2
1	B	727	ASN	4.1
1	B	726	ALA	4.1
1	B	663	GLY	4.1
1	B	734	MET	4.1
1	B	729	THR	4.0
1	B	733	TYR	3.9
1	A	662	ASN	3.9
1	B	467	GLU	3.8
1	A	659	LYS	3.7
1	B	762	THR	3.6
1	B	492	PHE	3.6
1	B	598	MET	3.3
1	B	743	VAL	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	676	PHE	3.2
1	A	763	LEU	3.2
1	B	728	CYS	3.2
1	A	652	ASN	3.1
1	B	745	SER	3.1
1	B	741	HIS	3.1
1	B	760	ILE	3.0
1	A	651	ILE	3.0
1	B	715	LEU	3.0
1	A	468	LEU	3.0
1	B	725	PRO	3.0
1	B	730	ASN	3.0
1	A	596	GLU	2.9
1	A	762	THR	2.9
1	A	764	THR	2.9
1	B	680[A]	TYR	2.9
1	B	746	GLN	2.8
1	B	677	ASP	2.8
1	B	742	ALA	2.7
1	B	744	PRO	2.7
1	A	593	VAL	2.7
1	B	719	GLY	2.7
1	B	713	PHE	2.6
1	B	723	ASP	2.6
1	A	540	MET	2.6
1	B	724	LYS	2.6
1	B	575	TYR	2.5
1	B	542	GLY	2.4
1	B	732	LEU	2.4
1	B	578	ALA	2.4
1	A	542	GLY	2.4
1	B	600	PHE	2.4
1	B	738	ASP	2.4
1	B	716	LEU	2.3
1	B	651	ILE	2.3
1	B	521	ASP	2.3
1	B	541	ILE	2.3
1	A	488	GLY	2.3
1	A	541	ILE	2.3
1	B	755	GLU	2.2
1	B	714	LYS	2.2
1	A	637	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	653	ASN	2.2
1	B	540	MET	2.1
1	A	470	GLU	2.1
1	B	752	GLN	2.1
1	A	489	GLU	2.0
1	B	705	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	801	5/5	0.68	0.19	89,92,92,93	0
2	SO4	A	801	5/5	0.75	0.17	64,68,69,69	0
3	ANP	A	802	31/31	0.79	0.17	13,38,70,71	0
3	ANP	B	802	31/31	0.87	0.16	11,37,94,94	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP A 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP B 802:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.