



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 29, 2026 – 11:22 AM UTC

PDB ID : 5UGX / pdb_00005ugx
Title : Crystal Structure of the Tyrosine Kinase Domain of FGF Receptor 2 Harboring a E565A/D650V double Gain-of-Function Mutation
Authors : Mohammadi, M.; Chen, H.
Deposited on : 2017-01-10
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	7% 79% 12% 8%
1	B	324	6% 75% 12% 13%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	282	Total	C	N	O	S	0	1	0
			2210	1408	375	405	22			
1	A	297	Total	C	N	O	S	0	1	0
			2360	1502	400	437	21			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	445	MET	-	initiating methionine	UNP P21802
B	446	GLY	-	expression tag	UNP P21802
B	447	SER	-	expression tag	UNP P21802
B	448	SER	-	expression tag	UNP P21802
B	449	HIS	-	expression tag	UNP P21802
B	450	HIS	-	expression tag	UNP P21802
B	451	HIS	-	expression tag	UNP P21802
B	452	HIS	-	expression tag	UNP P21802
B	453	HIS	-	expression tag	UNP P21802
B	454	HIS	-	expression tag	UNP P21802
B	455	SER	-	expression tag	UNP P21802
B	456	GLN	-	expression tag	UNP P21802
B	457	ASP	-	expression tag	UNP P21802
B	491	ALA	CYS	engineered mutation	UNP P21802
B	565	ALA	GLU	engineered mutation	UNP P21802
B	650	VAL	ASP	conflict	UNP P21802
A	445	MET	-	initiating methionine	UNP P21802
A	446	GLY	-	expression tag	UNP P21802
A	447	SER	-	expression tag	UNP P21802
A	448	SER	-	expression tag	UNP P21802
A	449	HIS	-	expression tag	UNP P21802
A	450	HIS	-	expression tag	UNP P21802
A	451	HIS	-	expression tag	UNP P21802
A	452	HIS	-	expression tag	UNP P21802
A	453	HIS	-	expression tag	UNP P21802

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	454	HIS	-	expression tag	UNP P21802
A	455	SER	-	expression tag	UNP P21802
A	456	GLN	-	expression tag	UNP P21802
A	457	ASP	-	expression tag	UNP P21802
A	491	ALA	CYS	engineered mutation	UNP P21802
A	565	ALA	GLU	engineered mutation	UNP P21802
A	650	VAL	ASP	conflict	UNP P21802

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total 31	C 11	N 5	O 12	P 3	0	0
3	A	1	Total 31	C 11	N 5	O 12	P 3	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	22	Total O 22 22	0	0
5	A	16	Total O 16 16	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibroblast growth factor receptor 2



• Molecule 1: Fibroblast growth factor receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.97Å 77.86Å 115.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.18 – 2.35 38.18 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.4 (38.18-2.35) 99.3 (38.18-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.35 (at 2.34Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.215 , 0.275 0.216 , 0.275	Depositor DCC
R_{free} test set	2000 reflections (7.71%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.723	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4692	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3511e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2414	0.63	2/3267 (0.1%)
1	B	0.40	0/2258	0.63	3/3054 (0.1%)
All	All	0.40	0/4672	0.63	5/6321 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	499	GLU	CG-CD-OE2	-6.91	102.52	118.40
1	B	574	GLU	CB-CA-C	6.83	121.20	109.24
1	B	574	GLU	CB-CG-CD	-5.66	102.97	112.60
1	A	499	GLU	CB-CA-C	5.56	119.39	110.16
1	B	574	GLU	N-CA-C	-5.26	106.99	113.41

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	499	GLU	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2360	0	2352	24	0
1	B	2210	0	2209	26	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	31	0	14	1	0
3	B	31	0	14	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	16	0	0	1	0
5	B	22	0	0	0	0
All	All	4692	0	4589	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:568:SER:OG	1:A:636:GLU:HG2	1.82	0.79
1:A:665:LEU:HD12	1:A:670:MET:HE3	1.71	0.73
1:B:507:LYS:HB2	1:B:510:GLU:HG3	1.69	0.72
1:B:704:TYR:H	1:B:722:MET:HE3	1.54	0.72
1:A:520:LYS:NZ	5:A:901:HOH:O	2.21	0.71
1:B:731:GLU:OE2	1:B:759:ARG:NH2	2.25	0.69
1:B:547:ILE:HD11	1:B:617:LEU:HD12	1.76	0.67
1:B:604:VAL:HG22	1:B:761:LEU:HD13	1.75	0.67
1:A:491:ALA:N	3:A:803:ACP:O3G	2.31	0.62
1:B:704:TYR:H	1:B:722:MET:CE	2.16	0.57
1:A:468:LEU:HD12	1:A:469:PRO:HD2	1.86	0.57
1:B:491:ALA:HB3	1:A:491:ALA:HB3	1.86	0.56
1:B:703:PRO:HA	1:B:722:MET:HE1	1.87	0.56
1:B:704:TYR:N	1:B:722:MET:HE3	2.21	0.55
1:B:537:MET:HA	1:B:540:MET:HE3	1.91	0.53
1:B:573:ARG:O	1:B:577:ARG:HG3	2.09	0.53
1:A:708:PRO:HG2	1:A:711:GLU:HB2	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:651:ILE:HG21	1:A:656:TYR:HB2	1.92	0.52
1:B:734:MET:HA	1:B:734:MET:HE2	1.91	0.51
1:A:573:ARG:O	1:A:577:ARG:HG3	2.10	0.51
1:B:697:PHE:CZ	1:B:732:LEU:HD13	2.46	0.51
1:A:637:ASN:HB2	1:A:639:VAL:HG23	1.92	0.51
1:A:485:LYS:HE2	1:A:497:MET:HE1	1.95	0.49
1:B:544:HIS:HB3	1:B:547:ILE:HD12	1.95	0.48
1:A:497:MET:HE2	1:A:566:TYR:CE1	2.49	0.48
1:A:501:VAL:HA	1:A:511:ALA:HA	1.95	0.47
1:A:625:ARG:HA	1:A:680:TYR:OH	2.13	0.47
1:A:625:ARG:HG2	1:A:680:TYR:CE1	2.50	0.46
1:A:554:CYS:HB2	1:A:561:TYR:HB2	1.98	0.46
1:B:694:TRP:CE2	1:B:722:MET:HE2	2.50	0.46
1:A:587:SER:HB3	1:A:590:ILE:O	2.16	0.45
1:B:554:CYS:HB2	1:B:561:TYR:HB2	1.98	0.45
1:A:520:LYS:HE2	1:A:522:ASP:OD1	2.17	0.45
1:A:676:PHE:CE1	1:A:717:LYS:HE2	2.52	0.44
1:B:627:LEU:HD22	1:B:692:LEU:HD22	1.99	0.44
1:B:605:SER:O	1:B:609:GLN:HG3	2.18	0.44
1:B:687:TRP:C	1:B:687:TRP:CD1	2.96	0.43
1:B:492:PHE:N	3:B:803:ACP:O2G	2.28	0.43
1:B:760:ILE:O	1:B:764:THR:HG23	2.18	0.43
1:B:625:ARG:HG2	1:B:680:TYR:CE1	2.52	0.43
1:A:491:ALA:HA	1:A:520:LYS:HE3	2.01	0.43
1:B:717:LYS:HB3	1:B:717:LYS:NZ	2.35	0.41
1:A:626:ASP:OD2	1:A:666:PRO:HG2	2.21	0.41
1:A:625:ARG:HG2	1:A:680:TYR:CZ	2.56	0.41
1:B:625:ARG:HG2	1:B:680:TYR:CZ	2.56	0.41
1:A:519:LEU:HD11	1:A:528:LEU:HB2	2.03	0.41
1:A:605:SER:O	1:A:609:GLN:HG3	2.21	0.41
1:B:592:ARG:HE	1:B:594:PRO:HG3	1.87	0.40
1:B:714:LYS:NZ	1:B:718:GLU:OE2	2.48	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/324 (91%)	286 (97%)	6 (2%)	4 (1%)	9	6
1	B	277/324 (86%)	274 (99%)	3 (1%)	0	100	100
All	All	573/648 (88%)	560 (98%)	9 (2%)	4 (1%)	24	18

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	652	ASN
1	A	644[A]	ASP
1	A	644[B]	ASP
1	A	584	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/283 (90%)	255 (100%)	0	100	100
1	B	238/283 (84%)	237 (100%)	1 (0%)	84	89
All	All	493/566 (87%)	492 (100%)	1 (0%)	87	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	604	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	556	GLN
1	B	620	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	727	ASN
1	B	746	GLN
1	A	546	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	802	-	4,4,4	0.34	0	6,6,6	0.20	0
2	SO4	B	801	-	4,4,4	0.26	0	6,6,6	0.19	0
2	SO4	B	802	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	A	801	-	4,4,4	0.25	0	6,6,6	0.19	0
3	ACP	B	803	4	31,33,33	1.83	7 (22%)	47,52,52	1.83	11 (23%)
3	ACP	A	803	4	31,33,33	1.96	10 (32%)	47,52,52	1.95	11 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACP	B	803	4	-	5/19/38/38	0/3/3/3
3	ACP	A	803	4	-	3/19/38/38	0/3/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	803	ACP	C6-N6	4.57	1.45	1.34
3	A	803	ACP	C6-N6	4.33	1.45	1.34
3	A	803	ACP	PG-O1G	3.96	1.58	1.50
3	B	803	ACP	C2'-C3'	-3.58	1.43	1.53
3	A	803	ACP	C2'-C3'	-3.48	1.43	1.53
3	B	803	ACP	C5-N7	-3.34	1.33	1.39
3	B	803	ACP	O4'-C1'	3.32	1.49	1.42
3	B	803	ACP	PG-O1G	3.26	1.56	1.50
3	A	803	ACP	O4'-C1'	3.13	1.49	1.42
3	A	803	ACP	PB-O1B	2.92	1.58	1.51
3	A	803	ACP	C5-N7	-2.88	1.33	1.39
3	A	803	ACP	C3'-C4'	-2.48	1.46	1.53
3	A	803	ACP	PB-O3A	2.41	1.61	1.58
3	B	803	ACP	C3'-C4'	-2.39	1.46	1.53
3	A	803	ACP	C2'-C1'	-2.15	1.46	1.53
3	A	803	ACP	C1'-N9	-2.08	1.40	1.46
3	B	803	ACP	C2'-C1'	-2.03	1.47	1.53

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	803	ACP	C5-C4-N3	-5.60	119.01	126.72
3	B	803	ACP	C5-C4-N3	-5.47	119.18	126.72
3	A	803	ACP	N3-C2-N1	-4.96	121.07	128.58
3	B	803	ACP	N3-C2-N1	-4.86	121.22	128.58
3	A	803	ACP	N3-C4-N9	4.47	134.76	127.17
3	B	803	ACP	N3-C4-N9	3.99	133.96	127.17
3	B	803	ACP	C5-N7-C8	3.64	109.18	103.45
3	A	803	ACP	C2-N3-C4	3.64	120.71	111.83
3	A	803	ACP	C5-N7-C8	3.55	109.04	103.45
3	B	803	ACP	C2-N3-C4	3.46	120.29	111.83
3	B	803	ACP	C4-C5-N7	-3.27	106.84	110.58
3	A	803	ACP	C3'-C2'-C1'	3.22	107.56	101.46
3	A	803	ACP	C2'-C1'-N9	-3.11	105.57	113.30
3	A	803	ACP	O4'-C1'-N9	2.77	113.42	108.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	803	ACP	O4'-C1'-N9	2.75	113.36	108.09
3	B	803	ACP	C3'-C2'-C1'	2.68	106.52	101.46
3	A	803	ACP	C4-C5-N7	-2.63	107.57	110.58
3	A	803	ACP	N9-C8-N7	-2.62	110.22	113.94
3	A	803	ACP	PB-O3A-PA	-2.32	124.80	132.37
3	B	803	ACP	N9-C8-N7	-2.25	110.74	113.94
3	B	803	ACP	C2'-C1'-N9	-2.21	107.82	113.30
3	B	803	ACP	PB-O3A-PA	-2.13	125.42	132.37

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	803	ACP	PB-C3B-PG-O1G
3	B	803	ACP	PB-C3B-PG-O2G
3	B	803	ACP	C5'-O5'-PA-O1A
3	A	803	ACP	PG-C3B-PB-O1B
3	A	803	ACP	PG-C3B-PB-O2B
3	B	803	ACP	PB-O3A-PA-O2A
3	B	803	ACP	PB-O3A-PA-O1A
3	A	803	ACP	PB-O3A-PA-O1A

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	803	ACP	1	0
3	A	803	ACP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/324 (91%)	0.50	23 (7%) 19 23	13, 27, 60, 83	1 (0%)
1	B	282/324 (87%)	0.45	18 (6%) 25 30	14, 28, 50, 64	1 (0%)
All	All	579/648 (89%)	0.47	41 (7%) 22 26	13, 27, 56, 83	2 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	587	SER	5.2
1	A	588	TYR	5.2
1	A	590	ILE	5.0
1	B	590	ILE	4.9
1	A	661	THR	4.8
1	A	491	ALA	4.4
1	B	574	GLU	4.3
1	A	663	GLY	4.3
1	A	662	ASN	4.1
1	A	763	LEU	3.5
1	B	491	ALA	3.5
1	A	764	THR	3.4
1	B	468	LEU	3.3
1	A	726	ALA	3.2
1	B	469	PRO	3.2
1	A	660	THR	3.2
1	A	583	GLY	3.1
1	A	492	PHE	2.8
1	B	661	THR	2.8
1	A	606	CYS	2.7
1	B	764	THR	2.7
1	A	586	TYR	2.7
1	B	763	LEU	2.7
1	B	746	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	584	MET	2.6
1	A	499	GLU	2.5
1	A	582	PRO	2.5
1	B	762	THR	2.4
1	B	492	PHE	2.4
1	B	741	HIS	2.4
1	A	589	ASP	2.4
1	B	582	PRO	2.3
1	B	676	PHE	2.2
1	B	644	ASP	2.2
1	A	727	ASN	2.2
1	B	583	GLY	2.2
1	B	488	GLY	2.2
1	A	468	LEU	2.0
1	A	509	LYS	2.0
1	B	541	ILE	2.0
1	A	469	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	802	5/5	0.65	0.17	68,68,68,69	0
4	MG	A	804	1/1	0.69	0.12	40,40,40,40	0
3	ACP	A	803	31/31	0.74	0.15	34,47,57,57	0
4	MG	B	804	1/1	0.82	0.13	48,48,48,48	0
3	ACP	B	803	31/31	0.85	0.11	33,41,53,54	0
2	SO4	A	801	5/5	0.86	0.13	55,55,56,56	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	802	5/5	0.86	0.12	62,62,62,62	0
2	SO4	B	801	5/5	0.91	0.10	66,66,66,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.