



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 06:58 AM UTC

PDB ID : 6UGM / pdb_00006ugm
EMDB ID : EMD-20765
Title : Structural basis of COMPASS eCM recognition of an unmodified nucleosome
Authors : Hsu, P.L.; Shi, H.; Zheng, N.
Deposited on : 2019-09-26
Resolution : 3.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11569 (3.20 - 4.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	 64% 8% 28%
1	E	135	 56% 13% 29%
2	B	102	 70% 10% 20%
2	F	102	 72% 13% 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	107	
3	G	107	
4	D	125	
5	H	125	
6	I	147	
7	J	146	
8	K	327	
9	M	275	
10	N	439	
11	R	3	
12	X	342	
13	L	405	
14	O	134	
14	P	134	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 23609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	97	Total	C	N	O	S	0	0
			799	504	155	138	2		
1	E	96	Total	C	N	O	S	0	0
			793	499	154	137	3		

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	82	Total	C	N	O	S	0	0
			657	416	128	112	1		
2	F	86	Total	C	N	O	S	0	0
			694	436	140	117	1		

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	107	Total	C	N	O	0	0
			823	519	161	143		
3	G	107	Total	C	N	O	0	0
			823	519	161	143		

- Molecule 4 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	93	Total	C	N	O	S	0	0
			726	457	130	137	2		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	7	PRO	ALA	conflict	UNP A0A1L8FQ56

- Molecule 5 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	H	93	Total	C	N	O	S	0	0
			725	456	130	137	2		

- Molecule 6 is a DNA chain called DNA (146-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	I	146	Total	C	N	O	P	0	0
			2975	1413	540	876	146		

- Molecule 7 is a DNA chain called DNA (146-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	J	146	Total	C	N	O	P	0	0
			3011	1425	564	876	146		

- Molecule 8 is a protein called Swd3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	K	305	Total	C	N	O	S	0	0
			2365	1498	395	453	19		

- Molecule 9 is a protein called Histone-lysine N-methyltransferase, H3 lysine-4 specific.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	M	188	Total	C	N	O	S	0	0
			1490	944	261	277	8		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	761	GLU	ASP	conflict	UNP Q6CIT4

- Molecule 10 is a protein called Swd1.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	N	408	Total	C	N	O	S	0	0
			3273	2095	539	622	17		

- Molecule 11 is a protein called H3 N-terminus.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	R	3	Total	C	N	O	S	0	0
			24	14	4	5	1		

- Molecule 12 is a protein called Spp1.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	X	73	Total	C	N	O		0	0
			604	379	108	117			


- Molecule 13 is a protein called Bre2.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	L	382	Total	C	N	O	S	0	0
			3128	1985	519	609	15		

- Molecule 14 is a protein called Sdc1.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	O	42	Total	C	N	O	S	0	0
			342	216	63	62	1		
14	P	43	Total	C	N	O	S	0	0
			329	211	57	59	2		

- Molecule 15 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).



Mol	Chain	Residues	Atoms					AltConf
15	M	1	Total	C	N	O	S	0
			27	15	6	5	1	

- Molecule 16 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
16	M	1	Total	Zn	0
			1	1	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3

Chain A: 




- Molecule 1: Histone H3

Chain E: 




- Molecule 2: Histone H4

Chain B: 



- Molecule 2: Histone H4

Chain F: 




- Molecule 3: Histone H2A


Chain C: 




- Molecule 3: Histone H2A




- Molecule 4: Histone H2B




- Molecule 5: Histone H2B 1.1




- Molecule 6: DNA (146-MER)



- Molecule 7: DNA (146-MER)






- Molecule 8: Swd3







• Molecule 9: Histone-lysine N-methyltransferase, H3 lysine-4 specific




• Molecule 10: Swd1



• Molecule 11: H3 N-terminus



• Molecule 12: Spp1



ASN
GLN
LYS
PRO
SER
SER

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	100905	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	74	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.119	Depositor
Minimum map value	-0.063	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.008	Depositor
Map size (Å)	342.144, 342.144, 342.144	wwPDB
Map dimensions	324, 324, 324	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.056, 1.056, 1.056	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/811	0.76	0/1087
1	E	0.63	0/804	1.00	3/1077 (0.3%)
2	B	0.70	0/664	0.93	3/889 (0.3%)
2	F	0.65	0/702	0.82	0/937
3	C	0.58	0/833	0.79	0/1124
3	G	0.57	0/833	0.84	2/1124 (0.2%)
4	D	0.67	0/737	0.75	0/993
5	H	0.70	0/736	0.79	0/991
6	I	0.43	0/3333	0.55	0/5137
7	J	0.43	0/3381	0.54	0/5221
8	K	0.36	1/2410 (0.0%)	0.89	8/3262 (0.2%)
9	M	0.42	3/1514 (0.2%)	0.90	8/2031 (0.4%)
10	N	0.34	0/3360	0.82	15/4577 (0.3%)
11	R	0.43	0/23	1.21	0/29
12	X	0.33	0/612	0.79	1/825 (0.1%)
13	L	0.34	0/3198	0.97	24/4323 (0.6%)
14	O	0.40	1/347 (0.3%)	0.77	2/466 (0.4%)
14	P	0.28	0/333	0.73	0/447
All	All	0.46	5/24631 (0.0%)	0.79	66/34540 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
8	K	0	1
13	L	0	2
All	All	0	4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	K	302	LEU	C-N	-8.44	1.23	1.33
14	O	114	PRO	N-CD	6.20	1.56	1.47
9	M	754	THR	C-N	-5.35	1.26	1.33
9	M	894	GLN	C-N	5.07	1.40	1.34
9	M	994	SER	C-N	-5.03	1.27	1.33

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	L	376	LYS	N-CA-C	-17.68	80.11	109.46
8	K	114	ILE	N-CA-C	16.12	125.41	110.74
13	L	376	LYS	CB-CA-C	13.39	131.76	109.80
10	N	20	LEU	CB-CA-C	-12.89	89.50	111.41
1	E	42	ARG	N-CA-C	12.88	126.19	109.83

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	45	THR	Peptide
8	K	113	VAL	Mainchain
13	L	308	PRO	Mainchain
13	L	309	PHE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	799	0	834	8	0
1	E	793	0	829	16	0
2	B	657	0	706	7	0
2	F	694	0	742	10	0
3	C	823	0	882	10	0
3	G	823	0	882	15	0
4	D	726	0	747	10	0
5	H	725	0	745	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	I	2975	0	1639	18	0
7	J	3011	0	1639	18	0
8	K	2365	0	2376	53	0
9	M	1490	0	1513	31	0
10	N	3273	0	3136	81	0
11	R	24	0	23	6	0
12	X	604	0	605	10	0
13	L	3128	0	3011	58	0
14	O	342	0	355	6	0
14	P	329	0	341	6	0
15	M	27	0	22	2	0
16	M	1	0	0	0	0
All	All	23609	0	21027	323	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 323 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:K:256:CYS:SG	8:K:302:LEU:O	2.09	1.11
10:N:210:TYR:HB2	10:N:260:VAL:CG1	1.83	1.08
10:N:210:TYR:HE1	10:N:261:SER:HA	1.08	1.08
10:N:210:TYR:CE1	10:N:261:SER:HA	1.88	1.08
9:M:914:LEU:HB2	11:R:3:THR:HG21	1.30	1.07

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/135 (70%)	92 (97%)	3 (3%)	0	100	100
1	E	94/135 (70%)	90 (96%)	4 (4%)	0	100	100
2	B	80/102 (78%)	78 (98%)	2 (2%)	0	100	100
2	F	84/102 (82%)	80 (95%)	4 (5%)	0	100	100
3	C	105/107 (98%)	100 (95%)	5 (5%)	0	100	100
3	G	105/107 (98%)	98 (93%)	7 (7%)	0	100	100
4	D	91/125 (73%)	89 (98%)	2 (2%)	0	100	100
5	H	91/125 (73%)	87 (96%)	4 (4%)	0	100	100
8	K	301/327 (92%)	278 (92%)	23 (8%)	0	100	100
9	M	182/275 (66%)	162 (89%)	20 (11%)	0	100	100
10	N	400/439 (91%)	369 (92%)	31 (8%)	0	100	100
11	R	1/3 (33%)	1 (100%)	0	0	100	100
12	X	69/342 (20%)	64 (93%)	5 (7%)	0	100	100
13	L	376/405 (93%)	347 (92%)	28 (7%)	1 (0%)	36	65
14	O	40/134 (30%)	38 (95%)	1 (2%)	1 (2%)	4	29
14	P	39/134 (29%)	37 (95%)	2 (5%)	0	100	100
All	All	2153/2997 (72%)	2010 (93%)	141 (6%)	2 (0%)	49	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
13	L	311	PRO
14	O	114	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/110 (76%)	84 (100%)	0	100	100
1	E	84/110 (76%)	84 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	68/78 (87%)	67 (98%)	1 (2%)	57	68
2	F	71/78 (91%)	71 (100%)	0	100	100
3	C	84/84 (100%)	78 (93%)	6 (7%)	13	40
3	G	84/84 (100%)	79 (94%)	5 (6%)	17	44
4	D	79/105 (75%)	78 (99%)	1 (1%)	61	71
5	H	79/105 (75%)	79 (100%)	0	100	100
8	K	275/298 (92%)	274 (100%)	1 (0%)	84	81
9	M	161/238 (68%)	159 (99%)	2 (1%)	63	72
10	N	360/387 (93%)	358 (99%)	2 (1%)	78	79
11	R	3/3 (100%)	3 (100%)	0	100	100
12	X	70/319 (22%)	69 (99%)	1 (1%)	59	70
13	L	352/375 (94%)	351 (100%)	1 (0%)	86	83
14	O	38/120 (32%)	38 (100%)	0	100	100
14	P	36/120 (30%)	32 (89%)	4 (11%)	6	25
All	All	1928/2614 (74%)	1904 (99%)	24 (1%)	61	72

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
9	M	888	VAL
10	N	12	VAL
10	N	5	LEU
12	X	215	GLU
3	C	118	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
12	X	307	ASN
13	L	344	ASN
12	X	322	GLN
13	L	181	GLN
14	O	98	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
15	SAM	M	1101	-	27,29,29	1.09	4 (14%)	34,42,42	1.97	10 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	SAM	M	1101	-	-	7/17/33/33	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	M	1101	SAM	C2-N1	2.37	1.38	1.33
15	M	1101	SAM	OXT-C	-2.36	1.23	1.30
15	M	1101	SAM	C5-N7	-2.32	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	M	1101	SAM	C2-N3	2.12	1.37	1.33

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	M	1101	SAM	N3-C2-N1	-5.22	120.67	128.58
15	M	1101	SAM	C5-C4-N3	-4.33	120.76	126.72
15	M	1101	SAM	N9-C8-N7	-4.13	108.08	113.94
15	M	1101	SAM	C5-N7-C8	3.55	109.03	103.45
15	M	1101	SAM	C2-N3-C4	3.25	119.77	111.83

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
15	M	1101	SAM	CB-CG-SD-CE
15	M	1101	SAM	CB-CG-SD-C5'
15	M	1101	SAM	O4'-C4'-C5'-SD
15	M	1101	SAM	C3'-C4'-C5'-SD
15	M	1101	SAM	CA-CB-CG-SD

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	M	1101	SAM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20765. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 162



Y Index: 162




Z Index: 162


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 167



Y Index: 167




Z Index: 181


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.008. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 411 nm^3 ; this corresponds to an approximate mass of 371 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20765 and PDB model 6UGM. Per-residue inclusion information can be found in section [3](#) on page [8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.008 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.008).

9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.008) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9120	 0.3850
A	 0.9660	 0.5170
B	 0.9830	 0.5190
C	 0.9350	 0.4790
D	 0.9820	 0.5090
E	 0.9670	 0.5260
F	 0.9580	 0.5190
G	 0.9400	 0.4950
H	 0.9890	 0.5160
I	 0.9710	 0.3770
J	 0.9730	 0.3780
K	 0.9520	 0.3300
L	 0.7260	 0.2460
M	 0.8810	 0.3680
N	 0.9480	 0.3980
O	 0.6130	 0.2420
P	 0.4750	 0.2670
R	 0.9170	 0.3180
X	 0.8780	 0.2810

