



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:21 AM UTC

PDB ID : 5UHN / pdb_00005uhn
Title : Crystal Structure of the Tyrosine Kinase Domain of FGF Receptor 2 harboring a N549H/E565A Double Gain-of-Function Mutation
Authors : Mohammadi, M.; Chen, H.
Deposited on : 2017-01-11
Resolution : 2.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	10% 74% 13% 13%
1	B	324	12% 73% 9% 17%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	1	0
			2220	1413	377	409	21			
1	B	269	Total	C	N	O	S	0	1	0
			2109	1345	360	383	21			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	445	MET	-	initiating methionine	UNP P21802
A	446	GLY	-	expression tag	UNP P21802
A	447	SER	-	expression tag	UNP P21802
A	448	SER	-	expression tag	UNP P21802
A	449	HIS	-	expression tag	UNP P21802
A	450	HIS	-	expression tag	UNP P21802
A	451	HIS	-	expression tag	UNP P21802
A	452	HIS	-	expression tag	UNP P21802
A	453	HIS	-	expression tag	UNP P21802
A	454	HIS	-	expression tag	UNP P21802
A	455	SER	-	expression tag	UNP P21802
A	456	GLN	-	expression tag	UNP P21802
A	457	ASP	-	expression tag	UNP P21802
A	491	ALA	CYS	conflict	UNP P21802
A	549	HIS	ASN	engineered mutation	UNP P21802
A	565	ALA	GLU	engineered mutation	UNP P21802
B	445	MET	-	initiating methionine	UNP P21802
B	446	GLY	-	expression tag	UNP P21802
B	447	SER	-	expression tag	UNP P21802
B	448	SER	-	expression tag	UNP P21802
B	449	HIS	-	expression tag	UNP P21802
B	450	HIS	-	expression tag	UNP P21802
B	451	HIS	-	expression tag	UNP P21802
B	452	HIS	-	expression tag	UNP P21802
B	453	HIS	-	expression tag	UNP P21802

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	454	HIS	-	expression tag	UNP P21802
B	455	SER	-	expression tag	UNP P21802
B	456	GLN	-	expression tag	UNP P21802
B	457	ASP	-	expression tag	UNP P21802
B	491	ALA	CYS	conflict	UNP P21802
B	549	HIS	ASN	engineered mutation	UNP P21802
B	565	ALA	GLU	engineered mutation	UNP P21802

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: C₁₁H₁₈N₅O₁₂P₃).


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibroblast growth factor receptor 2



• Molecule 1: Fibroblast growth factor receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.33Å 78.56Å 116.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.28 – 2.91 39.28 – 2.91	Depositor EDS
% Data completeness (in resolution range)	98.8 (39.28-2.91) 98.8 (39.28-2.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.17 (at 2.90Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.254 , 0.311 0.254 , 0.309	Depositor DCC
R_{free} test set	1404 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.775	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4405	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4793e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MG, ACP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/2269	0.29	0/3069
1	B	0.09	0/2155	0.28	0/2912
All	All	0.09	0/4424	0.29	0/5981

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2220	0	2201	23	0
1	B	2109	0	2106	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	31	0	14	2	0
3	B	31	0	14	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
All	All	4405	0	4335	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:490:GLY:HA3	3:A:802:ACP:H3B2	1.75	0.67
1:A:610:LEU:HD13	1:A:692:LEU:HD21	1.82	0.61
1:B:610:LEU:HD13	1:B:692:LEU:HD21	1.81	0.60
1:B:625:ARG:NH1	1:B:647:LEU:O	2.35	0.59
1:A:625:ARG:NH1	1:A:647:LEU:O	2.37	0.58
1:B:644:ASP:OD2	3:B:802:ACP:O1B	2.21	0.57
1:A:485:LYS:HG2	1:A:497:MET:HE2	1.88	0.56
1:A:667:VAL:HG12	1:A:712:LEU:HD23	1.87	0.55
1:A:554:CYS:HB2	1:A:561:TYR:HB2	1.90	0.54
1:B:536:GLU:HA	1:B:539:LYS:HD2	1.89	0.54
1:B:617:LEU:HD21	1:B:645:PHE:HE1	1.74	0.53
1:B:577:ARG:NH1	1:B:700:GLY:O	2.43	0.52
1:A:710:GLU:OE1	1:A:710:GLU:N	2.43	0.50
1:A:491:ALA:HB3	1:B:491:ALA:HB3	1.94	0.49
1:A:666:PRO:O	1:A:670:MET:HG3	2.13	0.49
1:B:537:MET:HG2	1:B:648:ALA:HB3	1.95	0.48
1:A:647:LEU:HD13	1:A:665:LEU:HD23	1.96	0.48
1:B:667:VAL:HG13	1:B:675:LEU:HD11	1.95	0.48
1:B:710:GLU:OE1	1:B:710:GLU:N	2.46	0.48
1:A:665:LEU:HD12	1:A:670:MET:HE3	1.96	0.47
1:B:647:LEU:HD13	1:B:665:LEU:HD23	1.97	0.46
1:A:536:GLU:HA	1:A:539:LYS:HD2	1.97	0.46
1:B:666:PRO:O	1:B:670:MET:HG3	2.15	0.46
1:A:693:MET:HB3	1:A:736:MET:HE2	1.98	0.45
1:A:579:ARG:HG2	1:A:598:MET:HE3	1.99	0.45
1:B:625:ARG:HA	1:B:680:TYR:CZ	2.52	0.45
1:B:724:LYS:NZ	1:B:728:CYS:O	2.49	0.44
1:B:537:MET:O	1:B:541:ILE:HG12	2.18	0.44
1:A:605:SER:O	1:A:609:GLN:HG3	2.19	0.43
1:A:673:GLU:OE2	1:A:747:ARG:NH1	2.40	0.43
1:B:554:CYS:HB2	1:B:561:TYR:HB2	2.01	0.42
1:A:488:GLY:HA3	3:A:802:ACP:H4'	2.00	0.42
1:A:655:ASP:HB3	1:A:679:VAL:HG13	2.02	0.42
1:A:675:LEU:HD21	1:A:713:PHE:HD1	1.84	0.42
1:A:483:LEU:HD21	1:A:518:MET:HE1	2.02	0.41
1:A:625:ARG:HA	1:A:680:TYR:CZ	2.55	0.41
1:B:630:ARG:NH2	1:B:666:PRO:HG2	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:697:PHE:CZ	1:A:732:LEU:HD23	2.56	0.41
1:A:608:TYR:CZ	1:A:612:ARG:HD2	2.56	0.41
1:B:605:SER:O	1:B:609:GLN:HG3	2.21	0.40
1:B:680:TYR:OH	1:B:685:ASP:OD1	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/324 (86%)	265 (95%)	14 (5%)	0	100	100
1	B	264/324 (82%)	255 (97%)	8 (3%)	1 (0%)	30	59
All	All	543/648 (84%)	520 (96%)	22 (4%)	1 (0%)	43	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	727	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/283 (84%)	232 (98%)	5 (2%)	47	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	225/283 (80%)	222 (99%)	3 (1%)	61	86
All	All	462/566 (82%)	454 (98%)	8 (2%)	53	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	540	MET
1	A	597	GLN
1	A	637	ASN
1	A	678	ARG
1	A	732	LEU
1	B	617	LEU
1	B	637	ASN
1	B	675	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	556	GLN
1	A	682	HIS
1	A	683	GLN
1	B	683	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	801	-	4,4,4	0.23	0	6,6,6	0.08	0
3	ACP	B	802	4	31,33,33	1.95	10 (32%)	47,52,52	1.94	10 (21%)
2	SO4	A	801	-	4,4,4	0.24	0	6,6,6	0.09	0
3	ACP	A	802	4	31,33,33	1.96	10 (32%)	47,52,52	2.00	11 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACP	B	802	4	-	7/19/38/38	0/3/3/3
3	ACP	A	802	4	-	3/19/38/38	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	ACP	C6-N6	4.54	1.45	1.34
3	B	802	ACP	C6-N6	4.50	1.45	1.34
3	A	802	ACP	C2'-C3'	-3.98	1.42	1.53
3	B	802	ACP	C2'-C3'	-3.89	1.42	1.53
3	A	802	ACP	PG-O1G	3.35	1.57	1.50
3	B	802	ACP	PG-O1G	3.32	1.56	1.50
3	B	802	ACP	C5-N7	-3.09	1.33	1.39
3	B	802	ACP	O4'-C1'	3.07	1.49	1.42
3	A	802	ACP	C5-N7	-3.04	1.33	1.39
3	A	802	ACP	O4'-C1'	2.95	1.48	1.42
3	B	802	ACP	C3'-C4'	-2.90	1.45	1.53
3	A	802	ACP	C3'-C4'	-2.70	1.46	1.53
3	A	802	ACP	C2'-C1'	-2.68	1.45	1.53
3	B	802	ACP	C2'-C1'	-2.54	1.45	1.53
3	A	802	ACP	PB-O2B	-2.33	1.50	1.56
3	A	802	ACP	PB-O1B	2.31	1.57	1.51
3	B	802	ACP	PB-O1B	2.29	1.57	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	ACP	C1'-N9	-2.28	1.40	1.46
3	B	802	ACP	PB-O2B	-2.27	1.50	1.56
3	B	802	ACP	C1'-N9	-2.23	1.40	1.46

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	ACP	N3-C2-N1	-5.62	120.07	128.58
3	B	802	ACP	N3-C2-N1	-5.40	120.41	128.58
3	A	802	ACP	C5-C4-N3	-5.33	119.37	126.72
3	B	802	ACP	C5-C4-N3	-5.29	119.43	126.72
3	A	802	ACP	N3-C4-N9	4.32	134.52	127.17
3	B	802	ACP	N3-C4-N9	4.31	134.50	127.17
3	A	802	ACP	C5-N7-C8	3.86	109.52	103.45
3	B	802	ACP	C5-N7-C8	3.80	109.42	103.45
3	A	802	ACP	C2-N3-C4	3.70	120.87	111.83
3	B	802	ACP	C2-N3-C4	3.59	120.60	111.83
3	B	802	ACP	PB-O3A-PA	-3.48	121.02	132.37
3	A	802	ACP	PB-O3A-PA	-3.13	122.15	132.37
3	A	802	ACP	C4-C5-N7	-3.06	107.09	110.58
3	B	802	ACP	C4'-O4'-C1'	-2.93	102.99	109.47
3	B	802	ACP	C4-C5-N7	-2.93	107.23	110.58
3	A	802	ACP	N9-C8-N7	-2.93	109.78	113.94
3	B	802	ACP	N9-C8-N7	-2.92	109.80	113.94
3	A	802	ACP	C4'-O4'-C1'	-2.86	103.16	109.47
3	B	802	ACP	C3'-C2'-C1'	2.75	106.66	101.46
3	A	802	ACP	C3'-C2'-C1'	2.43	106.06	101.46
3	A	802	ACP	C2'-C3'-C4'	2.41	107.27	102.61

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	ACP	PG-C3B-PB-O1B
3	A	802	ACP	PG-C3B-PB-O2B
3	A	802	ACP	PG-C3B-PB-O3A
3	B	802	ACP	PB-C3B-PG-O1G
3	B	802	ACP	PB-C3B-PG-O2G
3	B	802	ACP	PB-C3B-PG-O3G
3	B	802	ACP	C5'-O5'-PA-O1A
3	B	802	ACP	C5'-O5'-PA-O2A
3	B	802	ACP	C5'-O5'-PA-O3A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	802	ACP	O4'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	802	ACP	1	0
3	A	802	ACP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/324 (87%)	0.95	34 (12%) 8 7	15, 32, 69, 119	1 (0%)
1	B	269/324 (83%)	0.98	39 (14%) 6 4	16, 35, 63, 109	1 (0%)
All	All	551/648 (85%)	0.97	73 (13%) 7 6	15, 33, 68, 119	2 (0%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	660	THR	6.1
1	A	662	ASN	5.0
1	A	727	ASN	4.9
1	A	661	THR	4.8
1	A	730	ASN	4.5
1	B	490	GLY	4.2
1	A	492	PHE	4.2
1	A	664	ARG	4.2
1	B	491	ALA	4.1
1	A	542	GLY	3.8
1	B	549	HIS	3.8
1	B	729	THR	3.8
1	A	490	GLY	3.7
1	B	650	ASP	3.7
1	A	491	ALA	3.6
1	A	663	GLY	3.6
1	B	469	PRO	3.6
1	A	543	LYS	3.5
1	A	728	CYS	3.5
1	B	728	CYS	3.5
1	B	636	GLU	3.4
1	B	726	ALA	3.3
1	B	637	ASN	3.2
1	B	597	GLN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	730	ASN	3.2
1	A	469	PRO	3.0
1	B	575	TYR	2.9
1	A	596	GLU	2.8
1	A	729	THR	2.8
1	B	679	VAL	2.8
1	B	488	GLY	2.8
1	B	557	ASP	2.7
1	A	541	ILE	2.7
1	A	540	MET	2.7
1	A	679	VAL	2.7
1	B	763	LEU	2.6
1	A	637	ASN	2.6
1	B	632	VAL	2.6
1	B	727	ASN	2.6
1	B	540	MET	2.6
1	A	597	GLN	2.5
1	B	701	GLY	2.5
1	B	543	LYS	2.5
1	A	631	ASN	2.5
1	A	644	ASP	2.5
1	A	488	GLY	2.5
1	A	650	ASP	2.4
1	B	576	LEU	2.4
1	B	663	GLY	2.4
1	A	713	PHE	2.4
1	B	578	ALA	2.3
1	B	715	LEU	2.3
1	A	714	LYS	2.3
1	A	763	LEU	2.2
1	B	627	LEU	2.2
1	B	700	GLY	2.2
1	A	499	GLU	2.2
1	A	632	VAL	2.2
1	B	489	GLU	2.2
1	A	595	GLU	2.2
1	B	725	PRO	2.1
1	B	521	ASP	2.1
1	B	633	LEU	2.1
1	B	577	ARG	2.1
1	B	574	GLU	2.1
1	A	715	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	762	THR	2.1
1	A	571	ASN	2.0
1	B	580	ARG	2.0
1	B	571	ASN	2.0
1	B	541	ILE	2.0
1	B	698	THR	2.0
1	A	545	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	801	5/5	0.63	0.23	79,79,80,80	0
3	ACP	B	802	31/31	0.68	0.18	49,54,68,68	0
4	MG	B	803	1/1	0.68	0.24	80,80,80,80	0
3	ACP	A	802	31/31	0.69	0.17	45,51,68,68	0
2	SO4	A	801	5/5	0.71	0.16	79,79,79,80	0
4	MG	A	803	1/1	0.80	0.13	40,40,40,40	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP B 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ACP A 802:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.