



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 03:50 AM UTC

PDB ID : 7UIB / pdb_00007uib
Title : Crystal structure of BoNT/E receptor binding domain in complex with SV2, VHH, and sialic acid
Authors : Liu, Z.; Jin, R.; Chen, P.
Deposited on : 2022-03-29
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	129	21% 85% 14% .
1	E	129	44% 71% 27% .
2	C	105	6% 73% 17% 10%
2	F	105	9% 75% 14% 10%
3	A	407	5% 80% 18% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	407	 3% 81% 17%
4	G	2	 100%
4	H	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PG4	F	604	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 10570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VHH-G6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	128	Total	C	N	O	S	0	0	0
			980	619	159	198	4			
1	E	129	Total	C	N	O	S	0	0	0
			980	621	158	197	4			

- Molecule 2 is a protein called SV2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	95	Total	C	N	O	S	0	0	0
			802	516	130	152	4			
2	F	94	Total	C	N	O	S	0	0	0
			787	508	126	149	4			

- Molecule 3 is a protein called Neurotoxin type E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	405	Total	C	N	O	S	0	0	0
			3313	2097	572	635	9			
3	A	406	Total	C	N	O	S	0	0	0
			3327	2105	574	639	9			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O_4S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	F	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			13	8	5		

- Molecule 8 is N-acetyl-beta-neuraminic acid (CCD ID: SLB) (formula: $C_{11}H_{19}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	D	1	Total	C	N	O	0	0
			21	11	1	9		
8	A	1	Total	C	N	O	0	0
			21	11	1	9		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	5	Total 5	O 5	0	0
9	C	8	Total 8	O 8	0	0
9	E	11	Total 11	O 11	0	0
9	F	9	Total 9	O 9	0	0
9	D	75	Total 75	O 75	0	0
9	A	71	Total 71	O 71	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: VHH-G6




• Molecule 1: VHH-G6





• Molecule 2: SV2





• Molecule 2: SV2



• Molecule 3: Neurotoxin type E



• Molecule 3: Neurotoxin type E



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	141.37Å 172.18Å 137.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	109.26 – 2.77 109.26 – 2.77	Depositor EDS
% Data completeness (in resolution range)	99.6 (109.26-2.77) 99.6 (109.26-2.77)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.235 , 0.261 0.235 , 0.261	Depositor DCC
R_{free} test set	2090 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	54.1	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10570	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SLB, NAG, SO4, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.43	0/1005	0.70	0/1373
1	E	0.42	0/1005	0.72	0/1376
2	C	0.39	0/822	0.71	0/1105
2	F	0.38	0/807	0.65	0/1086
3	A	0.39	0/3397	0.72	0/4606
3	D	0.40	0/3383	0.73	0/4587
All	All	0.40	0/10419	0.72	0/14133

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	980	0	900	9	0
1	E	980	0	899	20	0
2	C	802	0	747	16	0
2	F	787	0	730	12	0
3	A	3327	0	3215	45	0
3	D	3313	0	3198	60	0
4	G	28	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	28	0	25	0	0
5	C	14	0	13	0	0
5	F	42	0	39	0	0
6	A	10	0	0	0	0
6	C	10	0	0	0	0
6	D	10	0	0	0	0
6	F	5	0	0	0	0
7	F	13	0	18	7	0
8	A	21	0	18	4	0
8	D	21	0	18	8	0
9	A	71	0	0	0	0
9	B	5	0	0	5	0
9	C	8	0	0	1	0
9	D	75	0	0	3	0
9	E	11	0	0	0	0
9	F	9	0	0	0	0
All	All	10570	0	9845	150	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 150 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:F:604:PG4:H42	3:D:1100:ARG:O	1.58	1.00
2:F:578:HIS:NE2	7:F:604:PG4:H51	1.82	0.94
7:F:604:PG4:H61	3:D:1099:ASP:OD1	1.72	0.89
2:C:515:ARG:HE	3:A:1158:HIS:HD2	1.20	0.85
7:F:604:PG4:C4	3:D:1100:ARG:O	2.30	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	126/129 (98%)	122 (97%)	4 (3%)	0	100	100
1	E	127/129 (98%)	121 (95%)	6 (5%)	0	100	100
2	C	93/105 (89%)	90 (97%)	3 (3%)	0	100	100
2	F	92/105 (88%)	89 (97%)	3 (3%)	0	100	100
3	A	404/407 (99%)	380 (94%)	24 (6%)	0	100	100
3	D	403/407 (99%)	382 (95%)	21 (5%)	0	100	100
All	All	1245/1282 (97%)	1184 (95%)	61 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	104/111 (94%)	100 (96%)	4 (4%)	29	61
1	E	103/111 (93%)	95 (92%)	8 (8%)	11	32
2	C	90/97 (93%)	89 (99%)	1 (1%)	65	85
2	F	88/97 (91%)	88 (100%)	0	100	100
3	A	374/378 (99%)	360 (96%)	14 (4%)	30	62
3	D	371/378 (98%)	363 (98%)	8 (2%)	45	75
All	All	1130/1172 (96%)	1095 (97%)	35 (3%)	35	68

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	1132	VAL
3	A	1138	ASN
3	A	1192	SER
1	E	127	VAL
1	E	126	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
3	A	859	ASN
3	A	941	ASN
3	A	935	ASN
3	A	964	GLN
2	F	543	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	1	4,2	14,14,15	0.28	0	17,19,21	0.82	0
4	NAG	G	2	4	14,14,15	0.31	0	17,19,21	0.48	0
4	NAG	H	1	4,2	14,14,15	0.36	0	17,19,21	1.33	1 (5%)
4	NAG	H	2	4	14,14,15	0.36	0	17,19,21	1.47	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,2	-	4/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	NAG	H	1	4,2	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	2	NAG	C1-O5-C5	4.00	117.55	112.19
4	H	1	NAG	C1-O5-C5	3.80	117.28	112.19
4	H	2	NAG	C4-C3-C2	2.82	115.16	111.02
4	H	2	NAG	C3-C4-C5	2.64	115.02	110.23

There are no chirality outliers.


5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
4	G	2	NAG	C4-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	F	601	2	14,14,15	0.49	0	17,19,21	0.78	1 (5%)
6	SO4	D	1302	-	4,4,4	0.34	0	6,6,6	0.07	0
6	SO4	C	603	-	4,4,4	0.35	0	6,6,6	0.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	601	2	14,14,15	0.52	0	17,19,21	1.25	1 (5%)
7	PG4	F	604	-	12,12,12	0.19	0	11,11,11	0.17	0
6	SO4	F	605	-	4,4,4	0.32	0	6,6,6	0.10	0
6	SO4	A	1302	-	4,4,4	0.32	0	6,6,6	0.06	0
6	SO4	D	1303	-	4,4,4	0.34	0	6,6,6	0.07	0
6	SO4	A	1303	-	4,4,4	0.35	0	6,6,6	0.07	0
8	SLB	A	1301	-	21,21,21	2.56	5 (23%)	24,31,31	1.56	3 (12%)
5	NAG	F	603	2	14,14,15	0.49	0	17,19,21	1.12	2 (11%)
5	NAG	F	602	2	14,14,15	0.67	0	17,19,21	1.19	2 (11%)
6	SO4	C	602	-	4,4,4	0.36	0	6,6,6	0.08	0
8	SLB	D	1301	-	21,21,21	2.65	7 (33%)	24,31,31	1.24	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	F	601	2	-	4/6/23/26	0/1/1/1
5	NAG	C	601	2	-	5/6/23/26	0/1/1/1
8	SLB	A	1301	-	-	9/20/38/38	0/1/1/1
5	NAG	F	603	2	-	4/6/23/26	0/1/1/1
5	NAG	F	602	2	-	3/6/23/26	0/1/1/1
7	PG4	F	604	-	-	5/10/10/10	-
8	SLB	D	1301	-	-	6/20/38/38	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	D	1301	SLB	O6-C2	9.21	1.52	1.43
8	A	1301	SLB	O6-C2	8.99	1.52	1.43
8	D	1301	SLB	C10-N5	3.68	1.46	1.34
8	A	1301	SLB	C10-N5	3.56	1.45	1.34
8	D	1301	SLB	C3-C4	-3.21	1.48	1.53

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1301	SLB	O6-C6-C7	4.30	113.37	106.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	601	NAG	C4-C3-C2	3.59	116.28	111.02
5	F	603	NAG	C4-C3-C2	3.40	116.00	111.02
5	F	602	NAG	C1-O5-C5	3.25	116.54	112.19
8	D	1301	SLB	C11-C10-N5	3.15	121.34	116.12

There are no chirality outliers.

5 of 36 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	601	NAG	C1-C2-N2-C7
5	F	602	NAG	C3-C2-N2-C7
5	F	602	NAG	C8-C7-N2-C2
5	F	602	NAG	O7-C7-N2-C2
8	A	1301	SLB	C4-C5-N5-C10


There are no ring outliers.


3 monomers are involved in 19 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	F	604	PG4	7	0
8	A	1301	SLB	4	0
8	D	1301	SLB	8	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	128/129 (99%)	1.42	27 (21%) 2 2	46, 72, 94, 101	0
1	E	129/129 (100%)	1.81	57 (44%) 0 0	37, 79, 110, 126	0
2	C	95/105 (90%)	0.66	6 (6%) 26 21	37, 54, 76, 104	0
2	F	94/105 (89%)	0.86	9 (9%) 13 11	38, 60, 85, 101	0
3	A	406/407 (99%)	0.61	22 (5%) 31 26	34, 50, 81, 102	0
3	D	405/407 (99%)	0.38	11 (2%) 56 48	29, 42, 63, 110	0
All	All	1257/1282 (98%)	0.76	132 (10%) 11 9	29, 51, 93, 126	0

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	847	ILE	6.5
1	E	75	ALA	5.7
1	E	2	LEU	5.7
3	A	1019	ASN	4.9
3	D	1135	SER	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	G	2	14/15	0.61	0.15	92,99,103,106	0
4	NAG	H	2	14/15	0.70	0.14	92,95,99,101	0
4	NAG	G	1	14/15	0.82	0.13	77,82,86,93	0
4	NAG	H	1	14/15	0.83	0.11	68,80,87,89	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain G:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	601	14/15	0.57	0.16	88,93,96,96	0
5	NAG	F	602	14/15	0.60	0.18	76,85,88,90	0
5	NAG	F	603	14/15	0.72	0.14	86,90,92,94	0
6	SO4	A	1303	5/5	0.74	0.14	105,105,107,108	0
5	NAG	F	601	14/15	0.75	0.15	74,83,87,88	0
8	SLB	D	1301	21/21	0.75	0.23	37,39,41,43	21
7	PG4	F	604	13/13	0.78	0.16	48,53,56,57	0
8	SLB	A	1301	21/21	0.78	0.17	77,85,89,90	0
6	SO4	D	1302	5/5	0.79	0.16	90,90,92,94	0
6	SO4	C	603	5/5	0.81	0.18	79,88,88,89	0
6	SO4	D	1303	5/5	0.84	0.16	83,84,89,91	0
6	SO4	F	605	5/5	0.88	0.10	74,75,78,81	0
6	SO4	C	602	5/5	0.90	0.14	71,71,75,75	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SO4	A	1302	5/5	0.90	0.12	79,81,86,86	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around SLB D 1301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SLB A 1301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 D 1302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 C 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 D 1303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SO4 F 605:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SO4 C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.