



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:17 AM UTC

PDB ID : 5UJ2 / pdb_00005uj2
Title : Crystal structure of HCV NS5B genotype 2A JFH-1 isolate with S15G E86Q E87Q C223H V321I mutations and Delta8 neta hairpoin loop deletion in complex with GS-639476 (diphosphate version of GS-9813), Mn²⁺ and symmetrical primer template 5'-AUAAAUUU
Authors : Edwards, T.E.; Fox III, D.; Appleby, T.C.; Murakami, E.; Rey, A.; McGrath, M.E.
Deposited on : 2017-01-16
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)
RNA backbone	3983	1120 (3.10-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	8	12% 50% 25% 25%
1	T	8	25% 38% 62%

Continued on next page...

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	A	572	% 84% 9% • 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(*AP*UP*AP*AP*AP*UP*UP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	T	8	Total	C	N	O	P	0	0	1
			147	66	23	51	7			
1	P	6	Total	C	N	O	P	0	0	1
			105	47	16	37	5			

- Molecule 2 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	534	Total	C	N	O	S	0	1	0
			4061	2567	714	754	26			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP R9TEU1
A	0	SER	-	expression tag	UNP R9TEU1
A	15	GLY	SER	engineered mutation	UNP R9TEU1
A	86	GLN	GLU	engineered mutation	UNP R9TEU1
A	87	GLN	GLU	engineered mutation	UNP R9TEU1
A	223	HIS	CYS	engineered mutation	UNP R9TEU1
A	321	ILE	VAL	engineered mutation	UNP R9TEU1
A	?	-	ASN	deletion	UNP R9TEU1
A	?	-	PHE	deletion	UNP R9TEU1
A	?	-	GLU	deletion	UNP R9TEU1
A	?	-	MET	deletion	UNP R9TEU1
A	?	-	TYR	deletion	UNP R9TEU1
A	?	-	GLY	deletion	UNP R9TEU1
A	?	-	SER	deletion	UNP R9TEU1
A	?	-	VAL	deletion	UNP R9TEU1
A	445	GLY	SER	conflict	UNP R9TEU1
A	571	LEU	-	expression tag	UNP R9TEU1
A	572	GLU	-	expression tag	UNP R9TEU1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	573	HIS	-	expression tag	UNP R9TEU1
A	574	HIS	-	expression tag	UNP R9TEU1
A	575	HIS	-	expression tag	UNP R9TEU1
A	576	HIS	-	expression tag	UNP R9TEU1
A	577	HIS	-	expression tag	UNP R9TEU1
A	578	HIS	-	expression tag	UNP R9TEU1

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	P	1	Total Mn 1 1	0	0
3	A	2	Total Mn 2 2	0	0

- Molecule 4 is (1S)-1-(4-aminoimidazo[2,1-f][1,2,4]triazin-7-yl)-1,4-anhydro-2-deoxy-2-fluoro-5-O-[(S)-hydroxy(phosphonooxy)phosphoryl]-2-methyl-D-ribose (CCD ID: 8B4) (formula: C₁₁H₁₆FN₅O₉P₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Cl 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	P	2	Total 2	O 2	0	0
6	A	23	Total 23	O 23	0	0

- Molecule 1: RNA (5'-R(*AP*UP*AP*AP*UP*UP*U)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	141.76Å 141.76Å 91.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.90 50.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.90) 99.1 (50.00-2.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.90Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.227 0.191 , 0.228	Depositor DCC
R_{free} test set	1167 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	63.1	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4370	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8B4, MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	P	0.41	0/116	0.81	0/179
1	T	0.43	0/163	0.78	0/252
2	A	0.64	0/4159	0.92	1/5672 (0.0%)
All	All	0.63	0/4438	0.91	1/6103 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	495	PRO	O-C-N	5.12	123.66	121.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	105	0	52	1	0
1	T	147	0	74	2	0
2	A	4061	0	3985	23	0
3	A	2	0	0	0	0
3	P	1	0	0	0	0
4	P	28	0	0	1	0
5	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	23	0	0	0	0
6	P	2	0	0	0	0
All	All	4370	0	4111	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:85:LEU:C	2:A:85:LEU:HD23	2.25	0.61
1:P:6:U:H2'	1:P:7:U:C6	2.40	0.57
2:A:18:GLU:CB	2:A:399:THR:HG21	2.37	0.55
2:A:203:TYR:CD2	2:A:314:LEU:HD13	2.44	0.53
2:A:203:TYR:CE2	2:A:314:LEU:HD13	2.47	0.50
1:T:4:A:H2'	1:T:5:A:C8	2.48	0.49
2:A:21:LEU:HB3	2:A:22:PRO:HD2	1.95	0.49
2:A:399:THR:HG22	2:A:401:ARG:H	1.78	0.49
1:T:2:U:O2	2:A:283:GLY:HA2	2.14	0.48
2:A:86:GLN:O	2:A:90:GLN:HG2	2.14	0.47
2:A:84:THR:O	2:A:85:LEU:C	2.56	0.47
2:A:326:SER:OG	2:A:332:ASP:OD2	2.30	0.46
2:A:132:THR:O	2:A:259:ARG:NE	2.38	0.45
2:A:434:MET:HE2	2:A:507:VAL:HG13	1.97	0.45
4:P:602:8B4:O3'	4:P:602:8B4:O2B	2.35	0.44
2:A:60:LEU:HD13	2:A:64:TYR:CE1	2.53	0.44
2:A:174:ALA:HB1	2:A:253:ILE:HD13	1.99	0.44
2:A:219:TYR:HB3	2:A:320:LEU:HD21	1.99	0.44
2:A:90:GLN:HE21	2:A:90:GLN:HA	1.83	0.43
2:A:111:LEU:O	2:A:112:SER:C	2.62	0.41
2:A:48:ARG:HG2	2:A:159:LEU:HG	2.01	0.41
2:A:175:LEU:O	2:A:176:TYR:C	2.64	0.41
2:A:31:LEU:HD12	2:A:31:LEU:C	2.45	0.41
2:A:399:THR:O	2:A:400:VAL:C	2.63	0.41
2:A:131:GLN:O	2:A:132:THR:C	2.63	0.40
2:A:524:TYR:CZ	2:A:539:LEU:HD21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	533/572 (93%)	513 (96%)	19 (4%)	1 (0%)	43 72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	130	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	428/486 (88%)	409 (96%)	19 (4%)	25 59

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	5	SER
2	A	19	GLU
2	A	29	SER
2	A	83	LEU
2	A	90	GLN
2	A	112	SER
2	A	117	ASN
2	A	139[A]	MET
2	A	139[B]	MET
2	A	314	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	315	VAL
2	A	324	SER
2	A	331	GLU
2	A	336	LEU
2	A	384	LEU
2	A	473	SER
2	A	483	THR
2	A	491	LYS
2	A	513	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	90	GLN
2	A	406	ASN
2	A	411	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	P	4/8 (50%)	0	0
1	T	6/8 (75%)	2 (33%)	0
All	All	10/16 (62%)	2 (20%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	T	7	U
1	T	8	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	8B4	P	602	3	24,30,30	1.25	4 (16%)	30,48,48	2.04	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8B4	P	602	3	-	2/14/36/36	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	P	602	8B4	C5-N4	-3.36	1.33	1.38
4	P	602	8B4	C9-N4	-2.86	1.32	1.37
4	P	602	8B4	PA-O3A	2.51	1.62	1.59
4	P	602	8B4	C5-N7	2.06	1.37	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	602	8B4	C8-C9-N4	5.80	108.02	103.88
4	P	602	8B4	C9-C8-N7	-5.26	105.94	111.62
4	P	602	8B4	C1'-C9-N4	4.26	131.07	123.09
4	P	602	8B4	C5-N4-N3	-3.59	123.57	127.27
4	P	602	8B4	C8-N7-C5	2.54	108.28	105.13

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	P	602	8B4	PA-O3A-PB-O1B
4	P	602	8B4	PA-O3A-PB-O2B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	P	602	8B4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	6/8 (75%)	0.20	1 (16%) 4 3	72, 74, 98, 123	0
1	T	8/8 (100%)	0.59	2 (25%) 2 1	57, 60, 100, 134	0
2	A	534/572 (93%)	-0.16	5 (0%) 81 75	34, 57, 83, 105	1 (0%)
All	All	548/588 (93%)	-0.15	8 (1%) 72 64	34, 57, 84, 134	1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	14	CYS	3.5
1	T	1	A	3.0
2	A	541	GLU	2.9
2	A	95	HIS	2.7
1	T	8	U	2.7
2	A	149	PRO	2.5
1	P	3	A	2.2
2	A	534	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	A	602	1/1	0.94	0.09	100,100,100,100	0
5	CL	A	603	1/1	0.94	0.07	70,70,70,70	0
4	8B4	P	602	28/28	0.98	0.06	45,55,58,59	0
3	MN	A	601	1/1	1.00	0.01	53,53,53,53	0
3	MN	P	601	1/1	1.00	0.03	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.