



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:59 AM UTC

PDB ID : 6UJ7 / pdb_00006uj7
Title : Crystal structure of HLA-B*07:02 with R140Q mutant IDH2 peptide
Authors : Miller, M.S.; Thirawatananond, P.; Gabelli, S.B.
Deposited on : 2019-10-02
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	5% 84% 9% 6%
1	D	298	2% 83% 9% 7%
2	B	119	2% 74% 10% 15%
2	E	119	3% 78% 7% 15%
3	C	10	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	10	 90%10%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, B-7 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	1	0
			2293	1421	419	447	6			
1	D	278	Total	C	N	O	S	0	0	0
			2274	1410	416	442	6			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P01889
A	281	GLY	-	expression tag	UNP P01889
A	282	SER	-	expression tag	UNP P01889
A	283	LEU	-	expression tag	UNP P01889
A	284	HIS	-	expression tag	UNP P01889
A	285	HIS	-	expression tag	UNP P01889
A	286	ILE	-	expression tag	UNP P01889
A	287	LEU	-	expression tag	UNP P01889
A	288	ASP	-	expression tag	UNP P01889
A	289	ALA	-	expression tag	UNP P01889
A	290	GLN	-	expression tag	UNP P01889
A	291	LYS	-	expression tag	UNP P01889
A	292	MET	-	expression tag	UNP P01889
A	293	VAL	-	expression tag	UNP P01889
A	294	TRP	-	expression tag	UNP P01889
A	295	ASN	-	expression tag	UNP P01889
A	296	HIS	-	expression tag	UNP P01889
A	297	ARG	-	expression tag	UNP P01889
D	0	MET	-	initiating methionine	UNP P01889
D	281	GLY	-	expression tag	UNP P01889
D	282	SER	-	expression tag	UNP P01889
D	283	LEU	-	expression tag	UNP P01889
D	284	HIS	-	expression tag	UNP P01889
D	285	HIS	-	expression tag	UNP P01889
D	286	ILE	-	expression tag	UNP P01889

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	287	LEU	-	expression tag	UNP P01889
D	288	ASP	-	expression tag	UNP P01889
D	289	ALA	-	expression tag	UNP P01889
D	290	GLN	-	expression tag	UNP P01889
D	291	LYS	-	expression tag	UNP P01889
D	292	MET	-	expression tag	UNP P01889
D	293	VAL	-	expression tag	UNP P01889
D	294	TRP	-	expression tag	UNP P01889
D	295	ASN	-	expression tag	UNP P01889
D	296	HIS	-	expression tag	UNP P01889
D	297	ARG	-	expression tag	UNP P01889

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	0	0
			843	536	142	162	3			
2	E	101	Total	C	N	O	S	0	0	0
			843	536	142	162	3			

- Molecule 3 is a protein called Isocitrate dehydrogenase [NADP], mitochondrial.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	0	0	0
			74	45	13	16			
3	F	10	Total	C	N	O	0	0	0
			74	45	13	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	7	GLN	ARG	engineered mutation	UNP P48735
F	7	GLN	ARG	engineered mutation	UNP P48735


- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		
5	B	1	Total	Na	0	0
			1	1		
5	D	1	Total	Na	0	0
			1	1		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	175	Total	O	0	0
			175	175		
7	B	53	Total	O	0	0
			53	53		
7	C	9	Total	O	0	0
			9	9		
7	D	184	Total	O	0	0
			184	184		
7	E	64	Total	O	0	0
			64	64		
7	F	11	Total	O	0	0
			11	11		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, B-7 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, B-7 alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Isocitrate dehydrogenase [NADP], mitochondrial

Chain C:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: Isocitrate dehydrogenase [NADP], mitochondrial

Chain F:  90% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.04Å 70.67Å 87.48Å 90.00° 107.39° 90.00°	Depositor
Resolution (Å)	47.43 – 1.90 47.43 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.43-1.90) 99.4 (47.43-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.182 , 0.238 0.189 , 0.242	Depositor DCC
R_{free} test set	3062 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6906	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, SO4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	2/2359 (0.1%)	1.30	7/3205 (0.2%)
1	D	1.02	0/2337	1.31	4/3175 (0.1%)
2	B	0.98	0/866	1.20	0/1171
2	E	0.95	0/866	1.22	0/1171
3	C	1.11	0/74	1.36	0/99
3	F	1.02	0/74	1.35	0/99
All	All	1.01	2/6576 (0.0%)	1.28	11/8920 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	50	PRO	C-O	-5.46	1.17	1.24
1	A	93	HIS	CE1-NE2	5.36	1.38	1.32

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ASP	CA-CB-CG	5.66	118.26	112.60
1	D	129	ASP	CA-CB-CG	5.60	118.20	112.60
1	A	224	GLN	CA-C-N	5.51	129.09	120.82
1	A	224	GLN	C-N-CA	5.51	129.09	120.82
1	A	225	THR	CA-C-N	5.51	127.66	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2293	0	2135	19	0
1	D	2274	0	2115	28	0
2	B	843	0	802	7	0
2	E	843	0	802	5	0
3	C	74	0	77	0	0
3	F	74	0	77	2	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
6	D	5	0	0	0	0
7	A	175	0	0	6	0
7	B	53	0	0	1	0
7	C	9	0	0	0	0
7	D	184	0	0	5	0
7	E	64	0	0	0	0
7	F	11	0	0	0	0
All	All	6906	0	6008	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:177:ASP:O	1:D:181:ARG:HB3	1.57	1.05
1:D:72:GLN:HG3	7:D:545:HOH:O	1.84	0.78
1:A:216[B]:THR:HG22	7:A:448:HOH:O	1.86	0.75
2:B:19:GLU:O	2:B:19:GLU:HG2	1.91	0.69
1:D:192:HIS:CE1	2:E:118:ASP:HB3	2.34	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/298 (94%)	272 (98%)	7 (2%)	0	100	100
1	D	276/298 (93%)	269 (98%)	7 (2%)	0	100	100
2	B	99/119 (83%)	97 (98%)	2 (2%)	0	100	100
2	E	99/119 (83%)	98 (99%)	0	1 (1%)	12	5
3	C	8/10 (80%)	8 (100%)	0	0	100	100
3	F	8/10 (80%)	8 (100%)	0	0	100	100
All	All	769/854 (90%)	752 (98%)	16 (2%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	20	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/254 (94%)	234 (98%)	5 (2%)	47	44
1	D	236/254 (93%)	229 (97%)	7 (3%)	36	30
2	B	95/109 (87%)	89 (94%)	6 (6%)	16	8
2	E	95/109 (87%)	93 (98%)	2 (2%)	47	44
3	C	9/9 (100%)	9 (100%)	0	100	100
3	F	9/9 (100%)	9 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	683/744 (92%)	663 (97%)	20 (3%)	37	31

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	225	THR
1	D	278	SER
2	E	69	VAL
2	E	21	ILE
2	B	39	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	115	GLN
1	D	141	GLN
1	D	224	GLN
1	A	144	GLN
1	A	224	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	D	302	-	4,4,4	0.30	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/298 (93%)	0.29	14 (5%) 34 36	19, 34, 62, 102	1 (0%)
1	D	278/298 (93%)	0.19	6 (2%) 62 66	21, 34, 61, 79	0
2	B	101/119 (84%)	0.20	2 (1%) 65 69	23, 36, 58, 75	0
2	E	101/119 (84%)	0.32	3 (2%) 52 56	22, 37, 58, 84	0
3	C	10/10 (100%)	-0.25	0 100 100	23, 30, 31, 33	0
3	F	10/10 (100%)	-0.23	0 100 100	24, 29, 31, 34	0
All	All	780/854 (91%)	0.23	25 (3%) 50 54	19, 34, 61, 102	1 (0%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	19	GLU	4.2
1	A	267	PRO	3.3
2	B	19	GLU	3.2
1	A	182	ALA	3.0
2	E	20	ALA	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	D	302	5/5	0.66	0.10	67,78,88,95	0
4	K	A	301	1/1	0.82	0.15	92,92,92,92	0
5	NA	B	201	1/1	0.93	0.08	46,46,46,46	0
5	NA	D	301	1/1	0.94	0.15	38,38,38,38	0
5	NA	A	302	1/1	0.97	0.10	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.