



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:56 PM UTC

PDB ID : 3ULF / pdb\_00003ulf  
Title : The light state structure of the blue-light photoreceptor Aureochrome1 LOV  
Authors : Mitra, D.; Yang, X.; Moffat, K.  
Deposited on : 2011-11-10  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	 59% 19% 2% 20%
1	B	170	 48% 26% 2% 24%
1	C	170	 48% 21% 3% 28%
1	D	170	 49% 21% 5% 24%
1	E	170	 52% 26% 2% 21%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	170	 49% 22% 25%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aureochrome1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	135	1067	675	181	208	3	0	0	0
1	B	128	1002	634	174	191	3	0	0	0
1	C	120	946	597	164	183	2	0	0	0
1	D	129	1014	643	175	193	3	0	0	0
1	E	135	1067	675	181	208	3	0	0	0
1	F	127	1003	636	174	190	3	0	1	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	168	GLY	-	expression tag	UNP A8QW55
A	169	SER	-	expression tag	UNP A8QW55
A	170	HIS	-	expression tag	UNP A8QW55
A	171	MET	-	expression tag	UNP A8QW55
A	172	GLY	-	expression tag	UNP A8QW55
A	173	HIS	-	expression tag	UNP A8QW55
A	174	SER	-	expression tag	UNP A8QW55
A	175	MET	-	expression tag	UNP A8QW55
B	168	GLY	-	expression tag	UNP A8QW55
B	169	SER	-	expression tag	UNP A8QW55
B	170	HIS	-	expression tag	UNP A8QW55
B	171	MET	-	expression tag	UNP A8QW55
B	172	GLY	-	expression tag	UNP A8QW55
B	173	HIS	-	expression tag	UNP A8QW55
B	174	SER	-	expression tag	UNP A8QW55
B	175	MET	-	expression tag	UNP A8QW55
C	168	GLY	-	expression tag	UNP A8QW55

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	169	SER	-	expression tag	UNP A8QW55
C	170	HIS	-	expression tag	UNP A8QW55
C	171	MET	-	expression tag	UNP A8QW55
C	172	GLY	-	expression tag	UNP A8QW55
C	173	HIS	-	expression tag	UNP A8QW55
C	174	SER	-	expression tag	UNP A8QW55
C	175	MET	-	expression tag	UNP A8QW55
D	168	GLY	-	expression tag	UNP A8QW55
D	169	SER	-	expression tag	UNP A8QW55
D	170	HIS	-	expression tag	UNP A8QW55
D	171	MET	-	expression tag	UNP A8QW55
D	172	GLY	-	expression tag	UNP A8QW55
D	173	HIS	-	expression tag	UNP A8QW55
D	174	SER	-	expression tag	UNP A8QW55
D	175	MET	-	expression tag	UNP A8QW55
E	168	GLY	-	expression tag	UNP A8QW55
E	169	SER	-	expression tag	UNP A8QW55
E	170	HIS	-	expression tag	UNP A8QW55
E	171	MET	-	expression tag	UNP A8QW55
E	172	GLY	-	expression tag	UNP A8QW55
E	173	HIS	-	expression tag	UNP A8QW55
E	174	SER	-	expression tag	UNP A8QW55
E	175	MET	-	expression tag	UNP A8QW55
F	168	GLY	-	expression tag	UNP A8QW55
F	169	SER	-	expression tag	UNP A8QW55
F	170	HIS	-	expression tag	UNP A8QW55
F	171	MET	-	expression tag	UNP A8QW55
F	172	GLY	-	expression tag	UNP A8QW55
F	173	HIS	-	expression tag	UNP A8QW55
F	174	SER	-	expression tag	UNP A8QW55
F	175	MET	-	expression tag	UNP A8QW55

- Molecule 2 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula:  $C_{17}H_{21}N_4O_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	E	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	F	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		
3	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total	O	0	0
			26	26		
4	B	19	Total	O	0	0
			19	19		
4	C	1	Total	O	0	0
			1	1		
4	E	14	Total	O	0	0
			14	14		
4	F	15	Total	O	0	0
			15	15		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aureochrome1

Chain A: 




- Molecule 1: Aureochrome1

Chain B: 




- Molecule 1: Aureochrome1

Chain C: 



- Molecule 1: Aureochrome1


Chain D: 






- Molecule 1: Aureochromel

Chain E: 52% 26% 21%



- Molecule 1: Aureochromel

Chain F: 49% 22% 25%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.98Å 73.98Å 176.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.10 – 2.90 34.10 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (34.10-2.90) 99.5 (34.10-2.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.09 (at 2.90Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, $R_{free}$	0.201 , 0.264 (Not available) , 0.237	Depositor DCC
$R_{free}$ test set	1077 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.4	Xtrriage
Anisotropy	0.469	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 70.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.479 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6370	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	106.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.29	0/1086	0.72	2/1475 (0.1%)
1	B	0.29	0/1018	0.75	0/1381
1	C	0.27	0/962	0.73	0/1306
1	D	0.27	0/1031	0.72	2/1399 (0.1%)
1	E	0.29	0/1086	0.72	2/1475 (0.1%)
1	F	0.30	0/1019	0.71	0/1383
All	All	0.29	0/6202	0.72	6/8419 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	203	ASP	CA-C-N	6.49	126.40	119.85
1	A	203	ASP	C-N-CA	6.49	126.40	119.85
1	D	259	GLY	CA-C-N	5.94	125.42	119.24
1	D	259	GLY	C-N-CA	5.94	125.42	119.24
1	E	259	GLY	CA-C-N	5.45	125.54	119.32

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1067	0	1048	33	0
1	B	1002	0	1003	40	0
1	C	946	0	933	31	0
1	D	1014	0	1012	36	0
1	E	1067	0	1048	39	0
1	F	1003	0	1006	37	0
2	A	31	0	19	3	0
2	B	31	0	19	1	0
2	C	31	0	19	2	0
2	D	31	0	19	1	0
2	E	31	0	19	1	0
2	F	31	0	19	5	0
3	B	5	0	0	0	0
3	F	5	0	0	0	0
4	A	26	0	0	0	0
4	B	19	0	0	0	0
4	C	1	0	0	0	0
4	E	14	0	0	0	0
4	F	15	0	0	0	0
All	All	6370	0	6164	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:CYS:H	1:A:332:GLN:HE22	1.07	0.98
1:D:211:ALA:HA	1:D:214:MET:HE3	1.53	0.87
1:C:222:THR:HG22	1:C:231:ILE:HG22	1.58	0.83
1:E:212:LEU:HD13	1:F:314:VAL:HG21	1.61	0.82
1:E:283:CYS:H	1:E:332:GLN:HE22	1.24	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/170 (78%)	131 (98%)	2 (2%)	0	100	100
1	B	126/170 (74%)	117 (93%)	9 (7%)	0	100	100
1	C	118/170 (69%)	110 (93%)	8 (7%)	0	100	100
1	D	127/170 (75%)	116 (91%)	10 (8%)	1 (1%)	16	44
1	E	133/170 (78%)	129 (97%)	4 (3%)	0	100	100
1	F	126/170 (74%)	121 (96%)	5 (4%)	0	100	100
All	All	763/1020 (75%)	724 (95%)	38 (5%)	1 (0%)	48	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	230	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/148 (80%)	111 (94%)	7 (6%)	18	48
1	B	111/148 (75%)	105 (95%)	6 (5%)	20	51
1	C	105/148 (71%)	95 (90%)	10 (10%)	8	26
1	D	112/148 (76%)	100 (89%)	12 (11%)	6	22
1	E	118/148 (80%)	115 (98%)	3 (2%)	42	74
1	F	111/148 (75%)	102 (92%)	9 (8%)	11	33
All	All	675/888 (76%)	628 (93%)	47 (7%)	14	40

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	270	ILE
1	E	239	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	278	VAL
1	D	316	VAL
1	E	270	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	317	GLN
1	F	272	ASN
1	D	289	GLN
1	F	312	ASN
1	E	332	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMN	D	501	1	33,33,33	1.70	4 (12%)	48,50,50	2.04	9 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FMN	C	501	1	33,33,33	1.69	4 (12%)	48,50,50	2.07	8 (16%)
3	PO4	F	502	-	4,4,4	0.84	0	6,6,6	0.65	0
3	PO4	B	502	-	4,4,4	0.83	0	6,6,6	0.69	0
2	FMN	A	501	1	33,33,33	1.72	4 (12%)	48,50,50	2.02	9 (18%)
2	FMN	F	501	1	33,33,33	1.69	4 (12%)	48,50,50	1.94	8 (16%)
2	FMN	E	501	1	33,33,33	1.72	4 (12%)	48,50,50	2.07	10 (20%)
2	FMN	B	501	1	33,33,33	1.68	4 (12%)	48,50,50	2.11	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	D	501	1	-	2/18/18/18	0/3/3/3
2	FMN	C	501	1	-	7/18/18/18	0/3/3/3
2	FMN	A	501	1	-	5/18/18/18	0/3/3/3
2	FMN	F	501	1	-	4/18/18/18	0/3/3/3
2	FMN	E	501	1	-	3/18/18/18	0/3/3/3
2	FMN	B	501	1	-	2/18/18/18	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FMN	C4A-N5	7.85	1.47	1.30
2	E	501	FMN	C4A-N5	7.76	1.47	1.30
2	D	501	FMN	C4A-N5	7.60	1.47	1.30
2	F	501	FMN	C4A-N5	7.59	1.47	1.30
2	B	501	FMN	C4A-N5	7.56	1.47	1.30

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FMN	C10-C4A-N5	-9.11	106.22	124.81
2	C	501	FMN	C10-C4A-N5	-9.03	106.37	124.81
2	D	501	FMN	C10-C4A-N5	-8.94	106.55	124.81
2	A	501	FMN	C10-C4A-N5	-8.78	106.87	124.81
2	E	501	FMN	C10-C4A-N5	-8.40	107.65	124.81

There are no chirality outliers.

5 of 23 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	FMN	C3'-C4'-C5'-O5'
2	A	501	FMN	C5'-O5'-P-O1P
2	C	501	FMN	C1'-C2'-C3'-C4'
2	C	501	FMN	C5'-O5'-P-O1P
2	C	501	FMN	C5'-O5'-P-O2P


There are no ring outliers.


6 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	FMN	1	0
2	C	501	FMN	2	0
2	A	501	FMN	3	0
2	F	501	FMN	5	0
2	E	501	FMN	1	0
2	B	501	FMN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	135/170 (79%)	-1.16	0 100 100	49, 66, 103, 178	0
1	B	128/170 (75%)	-1.03	0 100 100	49, 74, 153, 173	0
1	C	120/170 (70%)	-0.44	0 100 100	117, 169, 212, 222	0
1	D	129/170 (75%)	-0.46	0 100 100	108, 169, 217, 225	0
1	E	135/170 (79%)	-1.15	0 100 100	48, 65, 106, 188	0
1	F	127/170 (74%)	-1.07	0 100 100	29, 74, 125, 177	1 (0%)
All	All	774/1020 (75%)	-0.89	0 100 100	29, 82, 200, 225	1 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FMN	D	501	31/31	0.95	0.09	150,167,181,182	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMN	C	501	31/31	0.96	0.10	148,169,189,194	0
3	PO4	F	502	5/5	0.97	0.08	100,118,125,133	0
2	FMN	B	501	31/31	0.99	0.05	48,63,69,72	0
2	FMN	E	501	31/31	0.99	0.05	33,58,65,67	0
2	FMN	F	501	31/31	0.99	0.05	50,64,72,74	0
3	PO4	B	502	5/5	0.99	0.04	117,124,131,141	0
2	FMN	A	501	31/31	0.99	0.04	33,57,64,72	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around FMN B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FMN E 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FMN F 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.