



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:15 PM UTC

PDB ID : 6ULT / pdb_00006ult
Title : BRD2-BD2 in complex with the cyclic peptide 4.2_3
Authors : Patel, K.; Walshe, J.L.; Walport, L.J.; Mackay, J.P.
Deposited on : 2019-10-08
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	109	24% 81% 14% 5%
1	B	109	31% 84% 14% ..
1	C	109	17% 81% 12% 7%
1	D	109	42% 78% 19% ..
1	E	109	9% 88% 11% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	109	14% 82% 17% ..
1	G	109	56% 75% 17% • 6%
1	H	109	10% 83% 14% ..
2	I	13	85% 15%
2	J	13	62% 23% 15%
2	K	13	77% 15% 8%
2	L	13	15% 77% 23%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bromodomain-containing protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	0	0
			862	553	152	150	7			
1	B	108	Total	C	N	O	S	0	1	0
			895	571	157	160	7			
1	C	101	Total	C	N	O	S	0	0	0
			838	537	148	147	6			
1	D	107	Total	C	N	O	S	0	0	0
			887	567	156	157	7			
1	E	109	Total	C	N	O	S	0	2	0
			906	578	158	163	7			
1	F	108	Total	C	N	O	S	0	1	0
			898	574	158	159	7			
1	G	102	Total	C	N	O	S	6	1	0
			845	542	147	149	7			
1	H	108	Total	C	N	O	S	0	1	0
			897	572	157	161	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	346	SER	-	expression tag	UNP P25440
B	346	SER	-	expression tag	UNP P25440
C	346	SER	-	expression tag	UNP P25440
D	346	SER	-	expression tag	UNP P25440
E	346	SER	-	expression tag	UNP P25440
F	346	SER	-	expression tag	UNP P25440
G	346	SER	-	expression tag	UNP P25440
H	346	SER	-	expression tag	UNP P25440

- Molecule 2 is a protein called Cyclic peptide 4.2_3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	13	Total	C	N	O	S	0	0	0
			118	78	22	17	1			
2	J	11	Total	C	N	O	S	0	0	0
			101	67	19	14	1			
2	K	12	Total	C	N	O	S	0	0	0
			109	73	20	15	1			
2	L	10	Total	C	N	O	S	0	0	0
			90	61	15	13	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total	O	0	0
			11	11		
3	B	12	Total	O	0	0
			12	12		
3	C	7	Total	O	0	0
			7	7		
3	D	3	Total	O	0	0
			3	3		
3	E	12	Total	O	0	0
			12	12		
3	F	10	Total	O	0	0
			10	10		
3	G	12	Total	O	0	0
			12	12		
3	H	8	Total	O	0	0
			8	8		
3	J	1	Total	O	0	0
			1	1		
3	K	3	Total	O	0	0
			3	3		
3	L	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bromodomain-containing protein 2




- Molecule 1: Bromodomain-containing protein 2




- Molecule 1: Bromodomain-containing protein 2





- Molecule 1: Bromodomain-containing protein 2









- Molecule 1: Bromodomain-containing protein 2




- Molecule 1: Bromodomain-containing protein 2





- Molecule 1: Bromodomain-containing protein 2



- Molecule 1: Bromodomain-containing protein 2



- Molecule 2: Cyclic peptide 4.2_3



- Molecule 2: Cyclic peptide 4.2_3




- Molecule 2: Cyclic peptide 4.2_3

Chain K:  77% 15% 8%



- Molecule 2: Cyclic peptide 4.2_3

Chain L:  15% 77% 23%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.08Å 63.63Å 120.34Å 90.00° 108.39° 90.00°	Depositor
Resolution (Å)	42.50 – 2.80 42.50 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.2 (42.50-2.80) 98.5 (42.50-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.14_3260, PHENIX 1.14_3260	Depositor
R, R_{free}	0.284 , 0.335 0.283 , 0.335	Depositor DCC
R_{free} test set	1390 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.503	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 72.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	7526	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2657e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALY, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/887	0.32	0/1195
1	B	0.12	0/923	0.33	0/1242
1	C	0.11	0/862	0.32	0/1162
1	D	0.13	0/911	0.34	0/1226
1	E	0.13	0/939	0.32	0/1265
1	F	0.15	0/926	0.36	0/1246
1	G	0.14	0/873	0.36	0/1177
1	H	0.13	0/925	0.31	0/1245
2	I	0.16	0/92	0.39	0/120
2	J	0.16	0/75	0.42	0/97
2	K	0.14	0/83	0.40	0/108
2	L	0.14	0/65	0.40	0/86
All	All	0.13	0/7561	0.34	0/10169

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	862	0	844	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	895	0	873	11	0
1	C	838	0	815	8	0
1	D	887	0	867	15	0
1	E	906	0	880	8	0
1	F	898	0	881	10	0
1	G	845	0	825	13	0
1	H	897	0	872	10	0
2	I	118	0	123	2	0
2	J	101	0	104	1	0
2	K	109	0	115	2	0
2	L	90	0	91	0	0
3	A	11	0	0	0	0
3	B	12	0	0	0	0
3	C	7	0	0	0	0
3	D	3	0	0	0	0
3	E	12	0	0	0	0
3	F	10	0	0	0	0
3	G	12	0	0	2	0
3	H	8	0	0	0	0
3	J	1	0	0	0	0
3	K	3	0	0	0	0
3	L	1	0	0	0	0
All	All	7526	0	7290	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:396:LEU:HG	1:D:421:MET:HE1	1.57	0.85
1:B:433:HIS:HB3	1:B:436:VAL:HG23	1.73	0.70
1:H:380:ALA:HB3	1:H:383:LEU:HD12	1.76	0.68
1:G:356:ILE:HG12	1:G:449:ARG:HG3	1.79	0.64
1:A:356:ILE:HG12	1:A:449:ARG:HD2	1.82	0.62

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/109 (94%)	97 (95%)	5 (5%)	0	100	100
1	B	107/109 (98%)	103 (96%)	4 (4%)	0	100	100
1	C	99/109 (91%)	97 (98%)	2 (2%)	0	100	100
1	D	105/109 (96%)	100 (95%)	4 (4%)	1 (1%)	12	38
1	E	109/109 (100%)	103 (94%)	6 (6%)	0	100	100
1	F	107/109 (98%)	104 (97%)	3 (3%)	0	100	100
1	G	101/109 (93%)	97 (96%)	4 (4%)	0	100	100
1	H	107/109 (98%)	102 (95%)	5 (5%)	0	100	100
2	I	9/13 (69%)	9 (100%)	0	0	100	100
2	J	7/13 (54%)	7 (100%)	0	0	100	100
2	K	8/13 (62%)	8 (100%)	0	0	100	100
2	L	7/13 (54%)	7 (100%)	0	0	100	100
All	All	868/924 (94%)	834 (96%)	33 (4%)	1 (0%)	48	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	381	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/96 (95%)	90 (99%)	1 (1%)	65	87
1	B	96/96 (100%)	95 (99%)	1 (1%)	68	88
1	C	88/96 (92%)	88 (100%)	0	100	100
1	D	94/96 (98%)	93 (99%)	1 (1%)	65	87
1	E	98/96 (102%)	96 (98%)	2 (2%)	48	80
1	F	96/96 (100%)	95 (99%)	1 (1%)	68	88
1	G	90/96 (94%)	86 (96%)	4 (4%)	25	59
1	H	96/96 (100%)	93 (97%)	3 (3%)	35	70
2	I	9/9 (100%)	9 (100%)	0	100	100
2	J	7/9 (78%)	6 (86%)	1 (14%)	3	11
2	K	8/9 (89%)	8 (100%)	0	100	100
2	L	6/9 (67%)	6 (100%)	0	100	100
All	All	779/804 (97%)	765 (98%)	14 (2%)	51	82

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	376	VAL
1	G	402	LYS
2	J	8	ARG
1	H	409	ARG
1	H	436	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	405	ASN
1	C	429	ASN
1	G	429	ASN
1	F	387	HIS
1	F	429	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALY	L	2	2	10,11,12	0.90	0	7,12,14	0.97	0
2	ALY	L	9	2	10,11,12	0.90	0	7,12,14	0.84	0
2	ALY	I	9	2	10,11,12	0.90	0	7,12,14	0.71	0
2	ALY	K	2	2	10,11,12	0.86	0	7,12,14	0.53	0
2	ALY	J	9	2	10,11,12	0.87	0	7,12,14	0.95	0
2	ALY	I	2	2	10,11,12	0.90	0	7,12,14	0.74	0
2	ALY	K	9	2	10,11,12	0.89	0	7,12,14	0.64	0
2	ALY	J	2	2,1	10,11,12	0.90	0	7,12,14	1.58	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALY	L	2	2	-	5/9/10/12	-
2	ALY	L	9	2	-	2/9/10/12	-
2	ALY	I	9	2	-	1/9/10/12	-
2	ALY	K	2	2	-	1/9/10/12	-
2	ALY	J	9	2	-	6/9/10/12	-
2	ALY	I	2	2	-	5/9/10/12	-
2	ALY	K	9	2	-	1/9/10/12	-
2	ALY	J	2	2,1	-	4/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	2	ALY	CD-CG-CB	-3.66	99.83	113.62

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	2	ALY	N-CA-CB-CG
2	J	2	ALY	C-CA-CB-CG
2	L	2	ALY	N-CA-CB-CG
2	L	2	ALY	C-CA-CB-CG
2	L	2	ALY	O-C-CA-CB

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	9	ALY	1	0
2	J	9	ALY	1	0
2	I	2	ALY	1	0
2	K	9	ALY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/109 (95%)	1.46	26 (25%) 2 1	32, 57, 80, 99	0
1	B	108/109 (99%)	1.70	34 (31%) 1 1	52, 65, 75, 81	1 (0%)
1	C	101/109 (92%)	1.29	18 (17%) 4 3	38, 56, 86, 95	0
1	D	107/109 (98%)	2.01	46 (42%) 0 0	55, 74, 90, 108	0
1	E	109/109 (100%)	0.72	10 (9%) 14 10	16, 38, 63, 82	2 (1%)
1	F	108/109 (99%)	1.01	15 (13%) 6 5	19, 38, 67, 82	1 (0%)
1	G	102/109 (93%)	2.31	61 (59%) 0 0	39, 72, 94, 103	3 (2%)
1	H	108/109 (99%)	0.90	11 (10%) 12 9	25, 47, 79, 90	1 (0%)
2	I	10/13 (76%)	0.75	0 100 100	30, 35, 50, 62	0
2	J	8/13 (61%)	1.15	0 100 100	54, 56, 62, 63	0
2	K	9/13 (69%)	1.15	0 100 100	43, 49, 64, 65	0
2	L	7/13 (53%)	1.76	2 (28%) 1 1	54, 62, 70, 73	0
All	All	881/924 (95%)	1.41	223 (25%) 1 1	16, 58, 87, 108	8 (0%)

The worst 5 of 223 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	381	LEU	5.6
1	B	379[A]	SER	4.9
1	G	399	VAL	4.9
1	D	394	MET	4.7
1	B	428	TYR	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ALY	J	2	12/13	0.72	0.24	50,56,67,80	0
2	ALY	L	9	12/13	0.76	0.17	15,36,59,63	0
2	ALY	L	2	12/13	0.81	0.20	54,57,64,75	0
2	ALY	J	9	12/13	0.82	0.20	51,53,55,66	0
2	ALY	I	2	12/13	0.83	0.17	23,37,54,62	0
2	ALY	K	9	12/13	0.84	0.13	39,47,54,58	0
2	ALY	K	2	12/13	0.88	0.16	37,39,63,79	0
2	ALY	I	9	12/13	0.90	0.10	23,24,44,58	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.