



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 05:50 PM UTC

PDB ID : 6ULX / pdb_00006ulx
Title : Adenylation domain of a keto acid-selecting NRPS module bound to keto acyl
adenylate space group P43212
Authors : Alonzo, D.A.; Chiche-Lapierre, C.; Schmeing, T.M.
Deposited on : 2019-10-08
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	24% 88% 7% ••

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8709 atoms, of which 4239 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amino acid adenylation domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	551	Total	C	H	N	O	S	0	0	0
			8513	2733	4218	737	805	20			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	650	ALA	-	expression tag	UNP M5R382
A	651	ALA	-	expression tag	UNP M5R382
A	652	ALA	-	expression tag	UNP M5R382
A	653	GLU	-	expression tag	UNP M5R382
A	654	ASN	-	expression tag	UNP M5R382
A	655	LEU	-	expression tag	UNP M5R382
A	656	TYR	-	expression tag	UNP M5R382
A	657	PHE	-	expression tag	UNP M5R382
A	658	GLN	-	expression tag	UNP M5R382

- Molecule 2 is 5'-O-[(S)-hydroxy[(4-methyl-2-oxopentanoyl)oxy]phosphoryl]adenosine (CCD ID: QA7) (formula: C₁₆H₂₂N₅O₉P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	P	0	0
			52	16	21	5	9	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		
3	A	1	Total	O S	0	0
			5	4 1		
3	A	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	59	Total	O	0	0
			59	59		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amino acid adenylation domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.27Å 103.27Å 172.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.03 – 2.31 73.03 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.5 (73.03-2.31) 93.3 (73.03-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.32Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.222 , 0.249 0.227 , 0.254	Depositor DCC
R_{free} test set	1905 reflections (4.51%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.697	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 70.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8709	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, QA7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/4394	0.37	0/5967

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4295	4218	4217	32	2
2	A	31	21	0	0	0
3	A	85	0	0	1	0
4	A	59	0	0	0	0
All	All	4470	4239	4217	32	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:636:LYS:C	1:A:637:ILE:HD13	1.76	1.10
1:A:545:ILE:HG12	1:A:616:LEU:HD11	1.66	0.77
1:A:565:VAL:HG21	1:A:606:ILE:HD11	1.69	0.74
1:A:587:LEU:HD23	1:A:606:ILE:CG2	2.19	0.73
1:A:107:ALA:O	1:A:110:THR:HG22	1.89	0.72
1:A:342:MET:HE3	1:A:377:MET:HE3	1.72	0.72
1:A:587:LEU:HD23	1:A:606:ILE:HG21	1.73	0.71
1:A:637:ILE:HD13	1:A:637:ILE:N	2.07	0.68
1:A:636:LYS:O	1:A:637:ILE:HD13	1.95	0.66
1:A:636:LYS:C	1:A:637:ILE:CD1	2.62	0.65
1:A:309:ILE:HG23	1:A:310:VAL:HG23	1.79	0.63
1:A:559:ALA:HB1	1:A:587:LEU:HD21	1.83	0.61
1:A:100:LEU:HD22	1:A:256:VAL:HG21	1.83	0.59
1:A:210:THR:HG22	1:A:211:ASP:H	1.68	0.58
1:A:636:LYS:O	1:A:637:ILE:CD1	2.50	0.58
1:A:335:PRO:HG3	1:A:359:MET:HE2	1.87	0.57
1:A:190:ASN:OD1	1:A:193:VAL:HG23	2.05	0.56
1:A:587:LEU:HD23	1:A:606:ILE:HG23	1.89	0.55
1:A:545:ILE:CG1	1:A:616:LEU:HD11	2.38	0.54
1:A:210:THR:HG22	1:A:211:ASP:N	2.22	0.54
1:A:208:ILE:HD12	1:A:230:LEU:HD11	1.93	0.50
1:A:565:VAL:CG2	1:A:606:ILE:HD11	2.39	0.50
1:A:383:GLY:O	1:A:384:ALA:CB	2.60	0.50
1:A:559:ALA:CB	1:A:587:LEU:HD21	2.42	0.48
1:A:383:GLY:O	1:A:384:ALA:HB3	2.13	0.48
1:A:568:SER:HB2	1:A:587:LEU:HD11	1.95	0.48
1:A:491:ARG:NH1	3:A:709:SO4:O4	2.47	0.47
1:A:636:LYS:HE2	1:A:636:LYS:HB2	1.57	0.47
1:A:570:VAL:HG23	1:A:586:ILE:O	2.18	0.43
1:A:400:ALA:N	1:A:401:PRO:HD2	2.34	0.42
1:A:565:VAL:HG13	1:A:589:PHE:HB2	2.02	0.42
1:A:544:ILE:HG12	1:A:549:ASN:OD1	2.21	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:573:CYS:SG	1:A:573:CYS:HG[8_554]	1.20	0.40
1:A:573:CYS:CB	1:A:573:CYS:SG[8_554]	1.89	0.31

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	543/576 (94%)	533 (98%)	8 (2%)	2 (0%)	30	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	384	ALA
1	A	420	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/478 (95%)	445 (98%)	10 (2%)	45	63

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	VAL
1	A	262	SER
1	A	366	GLU
1	A	479	HIS
1	A	624	VAL
1	A	625	GLN
1	A	636	LYS
1	A	637	ILE
1	A	644	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	653	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	281	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	702	-	4,4,4	0.20	0	6,6,6	0.09	0
3	SO4	A	710	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	A	712	-	4,4,4	0.24	0	6,6,6	0.06	0
3	SO4	A	714	-	4,4,4	0.25	0	6,6,6	0.09	0
3	SO4	A	715	-	4,4,4	0.24	0	6,6,6	0.06	0
3	SO4	A	704	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	A	713	-	4,4,4	0.24	0	6,6,6	0.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	706	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	A	717	-	4,4,4	0.25	0	6,6,6	0.06	0
3	SO4	A	716	-	4,4,4	0.23	0	6,6,6	0.09	0
3	SO4	A	711	-	4,4,4	0.24	0	6,6,6	0.08	0
2	QA7	A	701	-	33,33,33	3.06	12 (36%)	48,49,49	1.92	12 (25%)
3	SO4	A	709	-	4,4,4	0.27	0	6,6,6	0.07	0
3	SO4	A	707	-	4,4,4	0.25	0	6,6,6	0.06	0
3	SO4	A	708	-	4,4,4	0.25	0	6,6,6	0.07	0
3	SO4	A	718	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	A	705	-	4,4,4	0.23	0	6,6,6	0.09	0
3	SO4	A	703	-	4,4,4	0.21	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QA7	A	701	-	-	2/23/39/39	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	QA7	C13-C12	-11.02	1.23	1.53
2	A	701	QA7	O25-C14	-5.42	1.29	1.42
2	A	701	QA7	C13-C14	4.76	1.68	1.53
2	A	701	QA7	O07-C06	4.47	1.42	1.37
2	A	701	QA7	P08-O07	4.28	1.68	1.60
2	A	701	QA7	C19-N24	4.20	1.44	1.34
2	A	701	QA7	C10-C11	-4.05	1.39	1.51
2	A	701	QA7	O25-C11	3.43	1.52	1.45
2	A	701	QA7	O27-C12	3.24	1.51	1.43
2	A	701	QA7	C12-C11	3.04	1.60	1.53
2	A	701	QA7	O26-C13	2.21	1.48	1.43
2	A	701	QA7	O31-C05	-2.18	1.18	1.23

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	QA7	O30-C06-C05	-5.32	115.94	122.05
2	A	701	QA7	C18-C23-N22	-5.18	119.58	126.72
2	A	701	QA7	N22-C21-N20	-4.30	122.08	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	QA7	N15-C16-N17	-3.71	108.67	113.94
2	A	701	QA7	N22-C23-N15	3.66	133.39	127.17
2	A	701	QA7	C21-N22-C23	3.55	120.50	111.83
2	A	701	QA7	O07-P08-O09	3.44	113.35	103.00
2	A	701	QA7	C18-N17-C16	2.81	107.87	103.45
2	A	701	QA7	C23-N15-C16	2.52	108.38	105.74
2	A	701	QA7	C04-C05-C06	2.25	119.91	115.89
2	A	701	QA7	C23-C18-N17	-2.22	108.05	110.58
2	A	701	QA7	C12-C13-C14	2.04	105.31	101.46

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	QA7	C06-O07-P08-O09
2	A	701	QA7	C06-O07-P08-O28

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	709	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/576 (95%)	1.40	136 (24%) 2 2	36, 59, 136, 160	0

All (136) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	575	VAL	11.8
1	A	569	PHE	10.5
1	A	622	ILE	10.4
1	A	635	GLY	8.8
1	A	585	LEU	8.7
1	A	616	LEU	8.5
1	A	624	VAL	8.1
1	A	602	ALA	7.9
1	A	587	LEU	7.7
1	A	574	SER	7.4
1	A	588	PHE	7.4
1	A	572	ALA	7.3
1	A	606	ILE	7.2
1	A	603	SER	7.1
1	A	593	LEU	7.1
1	A	646	TRP	7.1
1	A	589	PHE	6.9
1	A	620	ARG	6.9
1	A	263	GLY	6.8
1	A	571	ALA	6.8
1	A	600	MET	6.8
1	A	623	PRO	6.5
1	A	625	GLN	6.3
1	A	558	ILE	6.3
1	A	570	VAL	6.0
1	A	555	ILE	5.8
1	A	563	PRO	5.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	586	ILE	5.8
1	A	559	ALA	5.7
1	A	599	ILE	5.6
1	A	565	VAL	5.6
1	A	605	HIS	5.6
1	A	590	THR	5.5
1	A	594	TYR	5.4
1	A	618	ALA	5.2
1	A	596	PRO	5.2
1	A	621	ILE	5.1
1	A	644	THR	5.0
1	A	597	ALA	4.9
1	A	601	ARG	4.9
1	A	652	ALA	4.9
1	A	617	SER	4.9
1	A	610	ILE	4.8
1	A	637	ILE	4.7
1	A	573	CYS	4.7
1	A	562	VAL	4.7
1	A	568	SER	4.5
1	A	598	TYR	4.4
1	A	223	PHE	4.4
1	A	612	THR	4.3
1	A	564	GLY	4.3
1	A	619	SER	4.3
1	A	591	PRO	4.1
1	A	636	LYS	4.0
1	A	595	GLU	4.0
1	A	608	SER	4.0
1	A	266	GLY	4.0
1	A	642	LEU	3.9
1	A	567	THR	3.9
1	A	592	LYS	3.9
1	A	210	THR	3.8
1	A	638	GLU	3.8
1	A	609	HIS	3.8
1	A	604	GLN	3.8
1	A	611	ALA	3.7
1	A	92	ASP	3.6
1	A	262	SER	3.6
1	A	653	GLU	3.6
1	A	544	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	641	GLN	3.5
1	A	640	ALA	3.5
1	A	614	MET	3.3
1	A	615	GLY	3.2
1	A	566	GLU	3.2
1	A	450	ARG	3.2
1	A	211	ASP	3.2
1	A	557	ALA	3.1
1	A	643	LYS	3.1
1	A	540	LYS	3.1
1	A	607	LYS	3.1
1	A	649	GLY	3.0
1	A	517	ASP	2.9
1	A	552	ASN	2.9
1	A	613	LYS	2.9
1	A	187	ARG	2.9
1	A	547	GLY	2.8
1	A	560	GLU	2.8
1	A	546	ASN	2.7
1	A	561	GLU	2.7
1	A	543	ILE	2.7
1	A	121	LYS	2.7
1	A	645	ARG	2.7
1	A	650	ALA	2.6
1	A	209	LEU	2.6
1	A	230	LEU	2.6
1	A	364	GLU	2.6
1	A	215	ILE	2.6
1	A	550	TYR	2.6
1	A	231	ARG	2.5
1	A	212	ASP	2.5
1	A	95	GLN	2.5
1	A	457	THR	2.5
1	A	232	VAL	2.5
1	A	217	GLU	2.4
1	A	294	GLU	2.4
1	A	456	GLU	2.4
1	A	397	HIS	2.4
1	A	146	GLY	2.4
1	A	454	ALA	2.3
1	A	229	GLN	2.3
1	A	556	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	224	LEU	2.3
1	A	545	ILE	2.3
1	A	239	LEU	2.2
1	A	222	ALA	2.2
1	A	554	GLU	2.2
1	A	639	ARG	2.2
1	A	365	LYS	2.2
1	A	124	LEU	2.2
1	A	228	ASP	2.1
1	A	242	GLU	2.1
1	A	233	ALA	2.1
1	A	188	GLU	2.1
1	A	549	ASN	2.1
1	A	446	THR	2.1
1	A	216	GLU	2.1
1	A	147	LEU	2.1
1	A	241	LEU	2.1
1	A	511	GLN	2.1
1	A	647	GLN	2.1
1	A	227	THR	2.0
1	A	130	GLN	2.0
1	A	479	HIS	2.0
1	A	149	ALA	2.0
1	A	480	GLN	2.0
1	A	455	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	709	5/5	0.49	0.17	75,97,103,114	0
3	SO4	A	713	5/5	0.63	0.20	66,75,94,107	0
3	SO4	A	718	5/5	0.64	0.14	94,99,103,110	0
3	SO4	A	714	5/5	0.67	0.14	84,93,111,111	0
3	SO4	A	717	5/5	0.68	0.19	82,86,107,115	0
3	SO4	A	711	5/5	0.76	0.16	55,74,94,104	0
3	SO4	A	712	5/5	0.77	0.15	78,82,97,118	0
3	SO4	A	706	5/5	0.80	0.12	80,80,93,95	0
3	SO4	A	716	5/5	0.80	0.17	67,81,88,105	0
3	SO4	A	708	5/5	0.81	0.12	93,102,113,123	0
3	SO4	A	703	5/5	0.81	0.15	77,85,95,96	0
3	SO4	A	715	5/5	0.81	0.18	75,79,111,111	0
3	SO4	A	707	5/5	0.82	0.17	63,64,86,95	0
3	SO4	A	710	5/5	0.89	0.10	66,69,86,91	0
2	QA7	A	701	31/31	0.92	0.10	33,45,57,62	0
3	SO4	A	704	5/5	0.93	0.16	64,64,67,74	0
3	SO4	A	705	5/5	0.94	0.08	64,65,72,75	0
3	SO4	A	702	5/5	0.97	0.13	50,53,59,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around QA7 A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.