



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 08:40 AM EDT

PDB ID : 7ULL / pdb_00007ull
Title : Human Grp94 N-terminal domain in complex with 42C
Authors : Stachowski, T.R.; Nithianantham, S.; Vanarotti, M.; Fischer, M.
Deposited on : 2022-04-05
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	18% 70% 10% 20%
1	B	272	15% 62% 16% 22%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endoplasmin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	1	0
			1676	1061	283	326	6			
1	B	211	Total	C	N	O	S	0	3	0
			1627	1036	270	317	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	66	GLY	-	expression tag	UNP P14625
A	67	SER	-	expression tag	UNP P14625
A	68	HIS	-	expression tag	UNP P14625
B	66	GLY	-	expression tag	UNP P14625
B	67	SER	-	expression tag	UNP P14625
B	68	HIS	-	expression tag	UNP P14625

- Molecule 2 is N,N-dimethyl-7H-purin-6-amine (CCD ID: 42C) (formula: C₇H₉N₅) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			12	7	5		
2	B	1	Total	C	N	0	0
			12	7	5		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	17	Total	O	0	0
			17	17		
6	B	19	Total	O	0	0
			19	19		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endoplasmin



• Molecule 1: Endoplasmin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	63.92Å 89.46Å 98.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.73 – 2.31 44.73 – 2.31	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.73-2.31) 98.8 (44.73-2.31)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 2.32Å)	Xtriage
Refinement program	PHENIX 1.18	Depositor
R, R_{free}	0.258 , 0.296 0.261 , 0.301	Depositor DCC
R_{free} test set	1988 reflections (7.79%)	wwPDB-VP
Wilson B-factor (Å ²)	59.7	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 72.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3392	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.20 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4490e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DMS, PG4, 42C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/1701	0.34	0/2299
1	B	0.16	0/1658	0.32	0/2243
All	All	0.16	0/3359	0.33	0/4542

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1676	0	1640	20	0
1	B	1627	0	1569	27	0
2	A	12	0	9	0	0
2	B	12	0	9	0	0
3	A	13	0	18	1	0
4	A	4	0	6	1	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	17	0	0	0	0
6	B	19	0	0	0	0
All	All	3392	0	3267	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:402:PG4:H31	1:B:87:LYS:HE3	1.72	0.70
1:B:220:GLN:HE22	1:B:240:THR:HB	1.58	0.68
1:A:134:VAL:HB	1:A:279:ILE:CD1	2.26	0.65
1:A:192:ILE:HG23	1:A:197:VAL:HG23	1.88	0.56
1:B:270:LYS:HE3	1:B:271:TYR:CZ	2.40	0.56
1:A:210:ILE:HB	1:A:248:THR:HB	1.88	0.56
1:B:211:VAL:HB	1:B:223:TRP:HB3	1.87	0.56
1:B:272:SER:HB2	1:B:279:ILE:HG13	1.88	0.55
1:A:134:VAL:HB	1:A:279:ILE:HD13	1.88	0.55
1:A:166:ILE:HD12	1:A:166:ILE:O	2.07	0.55
1:A:211:VAL:HB	1:A:223:TRP:HB3	1.90	0.54
1:B:99:ILE:HD12	1:B:102:ARG:CZ	2.38	0.53
1:A:181:ALA:O	1:A:186:GLN:N	2.42	0.51
1:B:133:THR:HG22	1:B:278:PRO:HG2	1.91	0.51
1:B:275:ILE:HG21	1:B:279:ILE:HD11	1.92	0.51
1:A:237:ARG:HH12	4:A:403:DMS:H12	1.75	0.50
1:B:109:SER:HB2	1:B:275:ILE:HD12	1.94	0.50
1:B:192:ILE:HG23	1:B:197:VAL:HG23	1.95	0.49
1:A:89:ILE:HG23	1:A:192:ILE:HD13	1.94	0.49
1:A:334:GLU:HB3	1:A:336:MET:HE2	1.94	0.49
1:B:138:CYS:SG	1:B:264:ILE:HG13	2.53	0.48
1:A:190:GLU:O	1:A:194:GLN:HB2	2.14	0.47
1:B:260:GLU:HB2	1:B:263:THR:HG22	1.96	0.47
1:B:210:ILE:HB	1:B:248:THR:HB	1.95	0.47
1:B:275:ILE:HG22	1:B:277:PHE:H	1.79	0.47
1:A:135:LYS:HB3	1:A:333:TRP:CH2	2.50	0.46
1:B:98:GLU:H	1:B:98:GLU:CD	2.23	0.46
1:B:113:ASP:O	1:B:117:LEU:HG	2.16	0.46
1:A:254:GLU:H	1:A:254:GLU:CD	2.25	0.45
1:B:135:LYS:HB3	1:B:333:TRP:CZ3	2.52	0.45
1:B:131:GLU:HG3	1:B:150:THR:HG21	1.99	0.44
1:B:135:LYS:HB3	1:B:333:TRP:CH2	2.53	0.44
1:A:134:VAL:HB	1:A:279:ILE:HD12	1.99	0.44
1:A:284:SER:HB3	1:A:331:TRP:CE3	2.52	0.44
1:B:257:ASP:O	1:B:263:THR:HG21	2.17	0.44
1:A:254:GLU:OE1	1:A:254:GLU:N	2.51	0.43
1:B:219:THR:O	1:B:221:HIS:ND1	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ILE:HA	1:B:204:LEU:HD21	2.00	0.43
1:B:140:LYS:HA	1:B:259:LEU:HD13	2.00	0.43
1:A:272:SER:HB2	1:A:279:ILE:HG13	2.00	0.43
1:A:337:ASN:OD1	1:A:337:ASN:N	2.51	0.43
1:B:254:GLU:H	1:B:254:GLU:CD	2.27	0.42
1:B:254:GLU:N	1:B:254:GLU:OE1	2.51	0.42
1:A:277:PHE:O	1:A:337:ASN:ND2	2.53	0.42
1:B:251:LEU:HD12	1:B:259:LEU:HG	2.02	0.41
1:B:225:SER:HB2	1:B:230:PHE:HB3	2.02	0.41
1:A:129:ASN:OD1	1:A:131:GLU:HG2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/272 (78%)	204 (96%)	8 (4%)	0	100	100
1	B	206/272 (76%)	198 (96%)	8 (4%)	0	100	100
All	All	418/544 (77%)	402 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/244 (74%)	180 (100%)	0	100	100
1	B	173/244 (71%)	172 (99%)	1 (1%)	78	88
All	All	353/488 (72%)	352 (100%)	1 (0%)	86	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	134	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	42C	B	501	-	13,13,13	1.93	4 (30%)	17,18,18	1.16	1 (5%)
5	GOL	B	502	-	5,5,5	0.97	0	5,5,5	1.04	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	42C	A	401	-	13,13,13	1.98	4 (30%)	17,18,18	1.24	1 (5%)
4	DMS	A	403	-	3,3,3	0.67	0	3,3,3	0.52	0
3	PG4	A	402	-	12,12,12	0.54	0	11,11,11	0.27	0
5	GOL	A	404	-	5,5,5	1.00	0	5,5,5	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	502	-	-	0/4/4/4	-
2	42C	B	501	-	-	1/4/4/4	0/2/2/2
2	42C	A	401	-	-	3/4/4/4	0/2/2/2
3	PG4	A	402	-	-	4/10/10/10	-
5	GOL	A	404	-	-	0/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	42C	C3-N9	4.06	1.41	1.38
2	B	501	42C	C3-N9	3.17	1.40	1.38
2	B	501	42C	C2-C3	3.14	1.45	1.40
2	A	401	42C	C2-C3	3.12	1.45	1.40
2	B	501	42C	C1-N10	2.78	1.44	1.36
2	A	401	42C	C1-N10	2.55	1.43	1.36
2	B	501	42C	C5-N6	2.18	1.37	1.33
2	A	401	42C	C8-N7	2.14	1.40	1.35

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	42C	C12-N10-C11	2.65	124.70	116.18
2	B	501	42C	C12-N10-C11	2.38	123.82	116.18

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	42C	C2-C1-N10-C11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401	42C	N6-C1-N10-C11
3	A	402	PG4	O3-C5-C6-O4
3	A	402	PG4	O1-C1-C2-O2
2	A	401	42C	C2-C1-N10-C12
3	A	402	PG4	C4-C3-O2-C2
2	B	501	42C	C2-C1-N10-C11
3	A	402	PG4	C3-C4-O3-C5


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	DMS	1	0
3	A	402	PG4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 42C B 501




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/272 (79%)	1.40	48 (22%) 2 2	50, 78, 114, 137	1 (0%)
1	B	211/272 (77%)	1.23	42 (19%) 3 3	38, 65, 147, 199	3 (1%)
All	All	428/544 (78%)	1.31	90 (21%) 2 3	38, 73, 121, 199	4 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	174	PHE	7.8
1	A	330	VAL	5.1
1	A	174	PHE	5.0
1	A	175	LEU	5.0
1	B	178	MET	4.9
1	B	191	LEU	4.9
1	B	192	ILE	4.9
1	B	187	SER	4.7
1	A	176[A]	ASN	4.5
1	B	172	SER	4.4
1	A	285	LYS	4.3
1	B	175	LEU	4.3
1	A	88	LEU	4.2
1	B	188	THR	4.1
1	B	329	THR	4.0
1	A	167	ALA	3.9
1	B	195	PHE	3.8
1	B	285	LYS	3.8
1	A	132	LEU	3.8
1	B	330	VAL	3.7
1	A	195	PHE	3.7
1	B	179	THR	3.7
1	B	186	GLN	3.6
1	A	329	THR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	94	TYR	3.4
1	B	166	ILE	3.3
1	B	176	ASN	3.2
1	A	169	SER	3.2
1	A	261	LEU	3.2
1	A	179	THR	3.2
1	B	90	ILE	3.1
1	B	262	ASP	3.1
1	A	184	ASP	3.1
1	B	91	ASN	3.0
1	B	180	GLU	3.0
1	A	272	SER	3.0
1	B	240	THR	3.0
1	B	274	PHE	2.9
1	B	92	SER	2.8
1	B	177	LYS	2.8
1	B	261	LEU	2.8
1	A	194	GLN	2.8
1	A	274	PHE	2.7
1	A	177	LYS	2.7
1	A	238	GLY	2.6
1	A	108	ALA	2.6
1	A	120	LEU	2.6
1	B	226	ASP	2.6
1	B	173	GLU	2.5
1	B	189	SER	2.5
1	A	259	LEU	2.4
1	A	186	GLN	2.4
1	A	277	PHE	2.4
1	B	331	TRP	2.4
1	B	190	GLU	2.4
1	A	235	ASP	2.3
1	A	266	ASN	2.3
1	A	181	ALA	2.3
1	A	240	THR	2.3
1	A	106	SER	2.3
1	B	141	GLU	2.3
1	A	239	ASN	2.2
1	A	178	MET	2.2
1	A	182	GLN	2.2
1	A	126	LEU	2.2
1	B	219	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	189	SER	2.2
1	A	89	ILE	2.2
1	B	335	LEU	2.2
1	B	336	MET	2.2
1	A	267	LEU	2.2
1	B	259	LEU	2.2
1	A	262	ASP	2.2
1	A	271	TYR	2.1
1	B	271	TYR	2.1
1	A	134	VAL	2.1
1	B	198	GLY	2.1
1	A	141	GLU	2.1
1	A	185	GLY	2.1
1	A	331	TRP	2.1
1	A	92	SER	2.0
1	A	188	THR	2.0
1	B	138	CYS	2.0
1	A	124	ASN	2.0
1	B	88	LEU	2.0
1	A	192	ILE	2.0
1	B	99	ILE	2.0
1	A	250	VAL	2.0
1	B	142	LYS	2.0
1	A	145	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DMS	A	403	4/4	0.81	0.26	77,82,98,101	0
5	GOL	B	502	6/6	0.89	0.16	64,72,82,84	0
5	GOL	A	404	6/6	0.90	0.13	60,70,75,91	0
3	PG4	A	402	13/13	0.94	0.09	45,60,72,76	0
2	42C	A	401	12/12	0.94	0.12	53,64,71,72	0
2	42C	B	501	12/12	0.97	0.06	24,36,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 42C A 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 42C B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.