



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:14 AM UTC

PDB ID : 4UMX / pdb_00004umx
Title : IDH1 R132H in complex with cpd 1
Authors : Mathieu, M.; Marquette, J.P.
Deposited on : 2014-05-22
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	18% 90% 6% •
1	B	425	8% 87% 8% • 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ISOCITRATE DEHYDROGENASE [NADP] CYTOPLASMIC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	412	Total	C	N	O	S	0	0	0
			3265	2077	553	617	18			
1	B	405	Total	C	N	O	S	0	0	0
			3214	2046	545	605	18			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	415	SER	-	expression tag	UNP O75874
A	416	LEU	-	expression tag	UNP O75874
A	417	GLU	-	expression tag	UNP O75874
A	418	HIS	-	expression tag	UNP O75874
A	419	HIS	-	expression tag	UNP O75874
A	420	HIS	-	expression tag	UNP O75874
A	421	HIS	-	expression tag	UNP O75874
A	422	HIS	-	expression tag	UNP O75874
A	423	HIS	-	expression tag	UNP O75874
A	424	HIS	-	expression tag	UNP O75874
A	425	HIS	-	expression tag	UNP O75874
A	132	HIS	ARG	engineered mutation	UNP O75874
B	415	SER	-	expression tag	UNP O75874
B	416	LEU	-	expression tag	UNP O75874
B	417	GLU	-	expression tag	UNP O75874
B	418	HIS	-	expression tag	UNP O75874
B	419	HIS	-	expression tag	UNP O75874
B	420	HIS	-	expression tag	UNP O75874
B	421	HIS	-	expression tag	UNP O75874
B	422	HIS	-	expression tag	UNP O75874
B	423	HIS	-	expression tag	UNP O75874
B	424	HIS	-	expression tag	UNP O75874
B	425	HIS	-	expression tag	UNP O75874
B	132	HIS	ARG	engineered mutation	UNP O75874

- Molecule 2 is 2,6-bis(1H-imidazol-1-ylmethyl)-4-(2,4,4-trimethylpentan-2-yl)phenol (CCD ID: VVS) (formula: C₂₂H₃₀N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	27	22	4	1	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	17	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	377	Total	O	0	0
			377	377		
5	B	389	Total	O	0	0
			389	389		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ISOCITRATE DEHYDROGENASE [NADP] CYTOPLASMIC



• Molecule 1: ISOCITRATE DEHYDROGENASE [NADP] CYTOPLASMIC



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.41Å 81.41Å 305.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.67 – 1.88 78.67 – 1.88	Depositor EDS
% Data completeness (in resolution range)	100.0 (78.67-1.88) 100.0 (78.67-1.88)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.76 (at 1.87Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.182 , 0.211 0.184 , 0.216	Depositor DCC
R_{free} test set	4278 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7374	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VVS, GOL, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	3/3334 (0.1%)	1.15	5/4496 (0.1%)
1	B	0.87	4/3281 (0.1%)	1.11	5/4422 (0.1%)
All	All	0.86	7/6615 (0.1%)	1.13	10/8918 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	180	MET	SD-CE	-8.33	1.58	1.79
1	A	318	MET	SD-CE	-7.83	1.59	1.79
1	B	199	MET	SD-CE	-6.79	1.62	1.79
1	B	318	MET	SD-CE	-5.94	1.64	1.79
1	B	96	ASN	CG-ND2	-5.23	1.22	1.33
1	A	385	ASN	CG-ND2	-5.10	1.22	1.33
1	A	101	ASN	CG-OD1	5.04	1.33	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	271	ASN	N-CA-C	6.17	119.18	110.23
1	B	125	VAL	N-CA-C	-6.09	106.57	111.81
1	B	169	VAL	N-CA-C	-5.73	106.12	111.45
1	B	38	ASP	N-CA-C	-5.65	98.73	108.56
1	A	192	PHE	CA-CB-CG	5.40	119.20	113.80
1	A	38	ASP	N-CA-C	-5.36	99.23	108.56
1	A	79	ASP	CA-CB-CG	5.21	117.81	112.60
1	A	385	ASN	CA-CB-CG	5.16	117.76	112.60
1	B	299	ASP	CA-CB-CG	5.07	117.67	112.60
1	B	140	ARG	N-CA-C	5.00	116.41	108.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3265	0	3233	10	0
1	B	3214	0	3189	25	0
2	A	27	0	30	0	0
3	A	48	0	25	1	0
3	B	48	0	25	1	0
4	B	6	0	8	2	0
5	A	377	0	0	0	0
5	B	389	0	0	4	0
All	All	7374	0	6510	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:TYR:HB3	1:A:254:MET:HE1	1.44	0.99
1:A:315:HIS:HA	1:A:318:MET:HE2	1.54	0.89
1:B:315:HIS:HA	1:B:318:MET:HE2	1.55	0.86
1:B:135:TYR:CB	1:B:139:TYR:HD2	2.04	0.71
1:A:208:TYR:HB3	1:A:254:MET:CE	2.24	0.65
1:B:135:TYR:HB2	1:B:139:TYR:HD2	1.62	0.64
1:B:94:SER:OG	4:B:1416:GOL:H31	1.99	0.62
1:B:135:TYR:HB3	1:B:139:TYR:CD2	2.41	0.56
1:A:121:VAL:HB	1:A:124:TRP:CE2	2.41	0.56
1:B:135:TYR:HB3	1:B:139:TYR:HD2	1.70	0.55
1:B:182:MET:SD	1:B:273:ASP:HB3	2.47	0.55
1:B:286:GLY:HA2	1:B:287:SER:C	2.33	0.54
1:A:75:THR:O	3:A:1416:NAP:H2N	2.09	0.53
1:B:135:TYR:CB	1:B:139:TYR:CD2	2.89	0.53
1:B:255:VAL:O	1:B:259:MET:HG2	2.08	0.52
1:B:96:ASN:HB2	4:B:1416:GOL:H11	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:SER:HA	1:B:249:ARG:O	2.11	0.51
1:B:330:ILE:HD12	1:B:363:SER:HB3	1.91	0.51
1:B:75:THR:O	3:B:1415:NAP:H2N	2.11	0.50
1:B:141:ALA:HB1	1:B:180:MET:CE	2.42	0.50
1:B:292:THR:HB	5:B:2278:HOH:O	2.14	0.48
1:B:318:MET:HE3	1:B:325:THR:HG22	1.96	0.47
1:A:330:ILE:HD12	1:A:363:SER:HB3	1.95	0.47
1:B:22:ILE:HD11	1:B:327:THR:HB	1.97	0.47
1:A:156:TYR:CE2	1:B:147:PRO:HD2	2.51	0.46
1:B:141:ALA:HB1	1:B:180:MET:HE2	1.98	0.45
1:A:318:MET:HE3	1:A:325:THR:HG22	1.97	0.45
1:B:338:ARG:NH1	5:B:2270:HOH:O	2.50	0.44
1:B:338:ARG:HG2	5:B:2270:HOH:O	2.18	0.43
1:A:216:LEU:HD22	1:B:180:MET:HE1	2.01	0.42
1:B:133:HIS:HB2	1:B:192:PHE:CE2	2.55	0.41
1:A:255:VAL:O	1:A:259:MET:HG2	2.21	0.40
1:B:294:VAL:HG23	5:B:2270:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/425 (96%)	401 (98%)	8 (2%)	1 (0%)	43	34
1	B	401/425 (94%)	393 (98%)	8 (2%)	0	100	100
All	All	811/850 (95%)	794 (98%)	16 (2%)	1 (0%)	48	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/361 (96%)	340 (98%)	8 (2%)	44	29
1	B	343/361 (95%)	341 (99%)	2 (1%)	78	72
All	All	691/722 (96%)	681 (99%)	10 (1%)	59	48

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	ASN
1	A	187	LYS
1	A	236	LYS
1	A	251	ILE
1	A	260	LYS
1	A	270	LYS
1	A	312	VAL
1	A	385	ASN
1	B	171	ASN
1	B	312	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	53	ASN
1	A	96	ASN
1	A	163	GLN
1	A	348	ASN
1	B	53	ASN
1	B	132	HIS
1	B	184	ASN
1	B	198	GLN
1	B	271	ASN
1	B	277	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GOL	B	1416	-	5,5,5	0.06	0	5,5,5	0.38	0
2	VVS	A	1415	-	29,29,29	0.34	0	43,43,43	0.55	1 (2%)
3	NAP	A	1416	-	50,52,52	0.90	2 (4%)	71,80,80	0.78	2 (2%)
3	NAP	B	1415	-	50,52,52	1.07	2 (4%)	71,80,80	0.74	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	1416	-	-	2/4/4/4	-
2	VVS	A	1415	-	-	0/20/20/20	0/3/3/3
3	NAP	A	1416	-	-	8/35/67/67	0/5/5/5
3	NAP	B	1415	-	-	8/35/67/67	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1415	NAP	C2N-N1N	5.97	1.41	1.35
3	A	1416	NAP	C2N-N1N	4.14	1.39	1.35
3	A	1416	NAP	O7N-C7N	2.89	1.29	1.24
3	B	1415	NAP	O7N-C7N	2.88	1.29	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1416	NAP	C6N-N1N-C1D	2.91	125.43	119.73
3	A	1416	NAP	C2N-N1N-C1D	-2.64	113.31	119.13
3	B	1415	NAP	C6N-N1N-C1D	2.49	124.61	119.73
3	B	1415	NAP	C6N-N1N-C2N	-2.19	120.01	121.88
2	A	1415	VVS	C20-C14-C10	2.03	131.64	123.78

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1416	NAP	C5D-O5D-PN-O3
3	A	1416	NAP	C5D-O5D-PN-O2N
3	A	1416	NAP	O4D-C1D-N1N-C2N
3	A	1416	NAP	O4D-C1D-N1N-C6N
3	A	1416	NAP	C2D-C1D-N1N-C2N
3	B	1415	NAP	C5B-O5B-PA-O1A
3	B	1415	NAP	C5D-O5D-PN-O3
3	B	1415	NAP	C5D-O5D-PN-O2N
3	B	1415	NAP	O4D-C1D-N1N-C2N
3	B	1415	NAP	O4D-C1D-N1N-C6N
3	B	1415	NAP	C2D-C1D-N1N-C2N
4	B	1416	GOL	C1-C2-C3-O3
4	B	1416	GOL	O2-C2-C3-O3
3	A	1416	NAP	C5B-O5B-PA-O1A
3	A	1416	NAP	C5D-O5D-PN-O1N
3	B	1415	NAP	C5D-O5D-PN-O1N
3	A	1416	NAP	C2D-C1D-N1N-C6N
3	B	1415	NAP	C2D-C1D-N1N-C6N


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1416	GOL	2	0
3	A	1416	NAP	1	0
3	B	1415	NAP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/425 (96%)	0.62	77 (18%) 3 3	16, 27, 60, 85	0
1	B	405/425 (95%)	0.23	32 (7%) 18 20	16, 25, 55, 99	0
All	All	817/850 (96%)	0.43	109 (13%) 7 7	16, 26, 59, 99	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	414	LEU	8.6
1	A	285	TYR	6.8
1	A	284	GLY	6.5
1	B	135	TYR	6.1
1	B	286	GLY	6.0
1	A	414	LEU	5.9
1	B	139	TYR	5.7
1	B	276	VAL	5.3
1	A	277	GLN	4.9
1	A	272	TYR	4.9
1	A	239	PHE	4.9
1	A	241	ALA	4.8
1	A	245	TRP	4.8
1	A	251	ILE	4.7
1	B	278	SER	4.7
1	A	276	VAL	4.5
1	A	244	ILE	4.4
1	A	280	SER	4.3
1	A	120	LEU	4.2
1	A	286	GLY	4.2
1	A	223	PHE	4.1
1	A	243	LYS	4.1
1	A	413	LYS	4.0
1	A	134	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	259	MET	4.0
1	A	267	TRP	3.9
1	A	125	VAL	3.9
1	B	277	GLN	3.9
1	A	204	GLY	3.9
1	B	136	GLY	3.8
1	A	246	TYR	3.8
1	B	272	TYR	3.8
1	A	270	LYS	3.7
1	B	133	HIS	3.6
1	B	140	ARG	3.6
1	B	134	ALA	3.5
1	A	135	TYR	3.5
1	A	275	ASP	3.5
1	A	230	ILE	3.4
1	A	252	ASP	3.4
1	A	236	LYS	3.3
1	A	274	GLY	3.2
1	A	233	LYS	3.2
1	A	237	SER	3.2
1	B	123	GLY	3.2
1	A	231	TYR	3.2
1	A	258	ALA	3.1
1	A	201	LEU	3.1
1	B	212	LYS	3.1
1	A	250	LEU	3.0
1	A	235	TYR	3.0
1	A	278	SER	3.0
1	A	137	ASP	3.0
1	A	124	TRP	2.9
1	A	247	GLU	2.9
1	A	202	SER	2.9
1	A	205	TRP	2.8
1	A	283	GLN	2.8
1	B	274	GLY	2.8
1	A	139	TYR	2.8
1	A	136	GLY	2.7
1	A	127	PRO	2.7
1	A	3	LYS	2.7
1	B	124	TRP	2.6
1	A	249	ARG	2.6
1	B	252	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	298	PRO	2.6
1	A	215	ILE	2.5
1	B	256	ALA	2.5
1	B	273	ASP	2.5
1	A	281	VAL	2.5
1	B	137	ASP	2.5
1	B	321	LYS	2.5
1	A	269	CYS	2.4
1	B	258	ALA	2.4
1	A	140	ARG	2.4
1	A	208	TYR	2.4
1	A	273	ASP	2.4
1	A	254	MET	2.4
1	A	259	MET	2.4
1	A	299	ASP	2.4
1	B	125	VAL	2.4
1	A	232	ASP	2.3
1	B	257	GLN	2.3
1	A	256	ALA	2.3
1	A	194	HIS	2.3
1	B	269	CYS	2.3
1	A	234	GLN	2.2
1	B	3	LYS	2.2
1	A	197	PHE	2.2
1	A	174	GLU	2.2
1	A	212	LYS	2.2
1	A	160	ASP	2.1
1	A	248	HIS	2.1
1	B	171	ASN	2.1
1	A	226	ILE	2.1
1	A	203	LYS	2.1
1	A	282	ALA	2.1
1	B	122	SER	2.1
1	A	161	GLY	2.1
1	A	255	VAL	2.1
1	A	407	ILE	2.0
1	B	241	ALA	2.0
1	B	260	LYS	2.0
1	A	214	THR	2.0
1	A	238	GLN	2.0
1	A	242	GLN	2.0
1	A	121	VAL	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	255	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	VVS	A	1415	27/27	0.75	0.19	46,56,68,69	0
4	GOL	B	1416	6/6	0.80	0.18	57,58,59,59	0
3	NAP	B	1415	48/48	0.95	0.07	18,28,35,40	0
3	NAP	A	1416	48/48	0.96	0.07	16,23,32,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around VVS A 1415:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP B 1415:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.