



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:55 AM UTC

PDB ID : 5UMQ / pdb_00005umq
Title : Crystal structure of TnmS1, an antibiotic binding protein from Streptomyces sp. CB03234
Authors : Chang, C.Y.; Chang, C.; Nocek, B.; Rudolf, J.D.; Joachimiak, A.; Phillips Jr., G.N.; Shen, B.; Enzyme Discovery for Natural Product Biosynthesis (NatPro); Midwest Center for Structural Genomics (MCSG)
Deposited on : 2017-01-29
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	7% 84% 12%
1	B	142	11% 73% 11% 11%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyoxalase/bleomycin resistance protein/dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	125	Total	C	N	O	S	0	0	0
			997	634	171	188	4			
1	B	126	Total	C	N	O	S	0	0	0
			1003	637	172	190	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	HIS	-	expression tag	UNP A0A125SA24
A	-14	HIS	-	expression tag	UNP A0A125SA24
A	-13	HIS	-	expression tag	UNP A0A125SA24
A	-12	HIS	-	expression tag	UNP A0A125SA24
A	-11	HIS	-	expression tag	UNP A0A125SA24
A	-10	HIS	-	expression tag	UNP A0A125SA24
A	-9	SER	-	expression tag	UNP A0A125SA24
A	-8	SER	-	expression tag	UNP A0A125SA24
A	-7	GLY	-	expression tag	UNP A0A125SA24
A	-6	LEU	-	expression tag	UNP A0A125SA24
A	-5	VAL	-	expression tag	UNP A0A125SA24
A	-4	PRO	-	expression tag	UNP A0A125SA24
A	-3	ARG	-	expression tag	UNP A0A125SA24
A	-2	GLY	-	expression tag	UNP A0A125SA24
A	-1	SER	-	expression tag	UNP A0A125SA24
A	0	HIS	-	expression tag	UNP A0A125SA24
B	-15	HIS	-	expression tag	UNP A0A125SA24
B	-14	HIS	-	expression tag	UNP A0A125SA24
B	-13	HIS	-	expression tag	UNP A0A125SA24
B	-12	HIS	-	expression tag	UNP A0A125SA24
B	-11	HIS	-	expression tag	UNP A0A125SA24
B	-10	HIS	-	expression tag	UNP A0A125SA24
B	-9	SER	-	expression tag	UNP A0A125SA24
B	-8	SER	-	expression tag	UNP A0A125SA24
B	-7	GLY	-	expression tag	UNP A0A125SA24

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	LEU	-	expression tag	UNP A0A125SA24
B	-5	VAL	-	expression tag	UNP A0A125SA24
B	-4	PRO	-	expression tag	UNP A0A125SA24
B	-3	ARG	-	expression tag	UNP A0A125SA24
B	-2	GLY	-	expression tag	UNP A0A125SA24
B	-1	SER	-	expression tag	UNP A0A125SA24
B	0	HIS	-	expression tag	UNP A0A125SA24

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	83	Total O 84 84	0	1
2	B	48	Total O 48 48	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.57Å 66.12Å 103.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 50.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	91.9 (50.00-1.95) 91.9 (50.00-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.61 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.195 , 0.256 0.204 , 0.260	Depositor DCC
R_{free} test set	898 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2132	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.40 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.0759e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1021	0.81	0/1384
1	B	0.72	1/1027 (0.1%)	0.88	2/1392 (0.1%)
All	All	0.71	1/2048 (0.0%)	0.85	2/2776 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	103	PRO	N-CD	5.32	1.55	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	MET	CA-C-N	-5.05	113.52	119.84
1	B	102	MET	C-N-CA	-5.05	113.52	119.84

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	PRO	Peptide
1	B	57	THR	Peptide
1	B	58	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	997	0	961	4	0
1	B	1003	0	966	18	0
2	A	84	0	0	0	0
2	B	48	0	0	0	0
All	All	2132	0	1927	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:PRO:O	1:B:105:GLY:N	2.00	0.93
1:B:58:LYS:HG2	1:B:63:TYR:CD2	2.06	0.91
1:A:84:THR:HG23	1:B:1:MET:HG2	1.76	0.67
1:B:43:GLU:OE1	1:B:52:ARG:HD2	1.99	0.62
1:B:102:MET:HG3	1:B:105:GLY:O	2.06	0.56
1:B:58:LYS:HG2	1:B:63:TYR:CG	2.40	0.55
1:A:43:GLU:OE1	1:A:52:ARG:HD2	2.05	0.55
1:B:81:ILE:H	1:B:123:GLN:HE21	1.55	0.53
1:B:21:PHE:HA	1:B:25:THR:HG23	1.93	0.51
1:B:103:PRO:C	1:B:105:GLY:H	2.08	0.51
1:B:104:PHE:HD1	1:B:104:PHE:O	1.94	0.50
1:B:102:MET:HG3	1:B:106:GLY:O	2.12	0.49
1:B:7:THR:HB	1:B:54:VAL:HG23	1.98	0.46
1:B:9:ASP:OD2	1:B:62:VAL:HG11	2.16	0.46
1:B:103:PRO:C	1:B:105:GLY:N	2.71	0.44
1:B:104:PHE:O	1:B:104:PHE:CD1	2.70	0.44
1:A:108:ASN:HD22	1:A:108:ASN:C	2.26	0.44
1:B:102:MET:HE2	1:B:102:MET:HB3	1.83	0.44
1:B:58:LYS:CG	1:B:63:TYR:CD2	2.91	0.43
1:A:112:GLN:HA	1:A:117:ASN:O	2.21	0.41
1:B:103:PRO:HB2	1:B:104:PHE:H	1.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/142 (87%)	119 (97%)	4 (3%)	0	100	100
1	B	124/142 (87%)	114 (92%)	6 (5%)	4 (3%)	3	0
All	All	247/284 (87%)	233 (94%)	10 (4%)	4 (2%)	7	2

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	104	PHE
1	B	105	GLY
1	B	39	MET
1	B	103	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/122 (88%)	106 (99%)	1 (1%)	70	70
1	B	108/122 (88%)	101 (94%)	7 (6%)	15	6
All	All	215/244 (88%)	207 (96%)	8 (4%)	30	20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	B	25	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	39	MET
1	B	59	HIS
1	B	62	VAL
1	B	65	GLU
1	B	101	GLN
1	B	102	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	59	HIS
1	B	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	125/142 (88%)	0.24	10 (8%) 18 21	7, 16, 39, 50	0
1	B	126/142 (88%)	0.78	15 (11%) 9 10	12, 23, 59, 71	0
All	All	251/284 (88%)	0.51	25 (9%) 12 14	7, 20, 47, 71	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	PHE	4.9
1	B	36	PHE	4.5
1	B	59	HIS	4.4
1	B	38	PRO	3.6
1	B	39	MET	3.6
1	A	36	PHE	3.5
1	B	62	VAL	3.3
1	A	41	TRP	3.1
1	B	63	TYR	3.0
1	B	101	GLN	2.7
1	B	58	LYS	2.6
1	A	0	HIS	2.6
1	B	103	PRO	2.6
1	A	65	GLU	2.6
1	B	105	GLY	2.4
1	A	104	PHE	2.4
1	B	33	ASP	2.4
1	A	102	MET	2.3
1	A	105	GLY	2.2
1	B	69	GLY	2.2
1	B	125	SER	2.1
1	B	79	GLU	2.1
1	A	37	GLY	2.1
1	A	106	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	38	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.