



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 29, 2026 – 02:19 PM UTC

PDB ID : 1UN6 / pdb_00001un6
Title : THE CRYSTAL STRUCTURE OF A ZINC FINGER - RNA COMPLEX
REVEALS TWO MODES OF MOLECULAR RECOGNITION
Authors : Lu, D.; Searles, M.A.; Klug, A.
Deposited on : 2003-09-04
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)
RNA backbone	3983	1022 (3.32-2.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	87	53% 41% 5% .
1	C	87	47% 44% 9%
1	D	87	33% 24% 9% 33%
2	E	61	26% 70%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	61	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a green segment on the left labeled '59%', a yellow segment in the middle labeled '38%', and a small orange segment on the right labeled with a dot '•'.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4554 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTION FACTOR IIIA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	87	Total	C	N	O	S	0	0	1
			713	451	133	121	8			
1	C	87	Total	C	N	O	S	0	0	0
			721	455	133	125	8			
1	D	58	Total	C	N	O	S	0	0	0
			475	299	86	85	5			

- Molecule 2 is a RNA chain called 5S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	61	Total	C	N	O	P	0	0	0
			1304	582	240	422	60			
2	F	61	Total	C	N	O	P	0	0	0
			1304	582	240	422	60			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Zn	0	0
			3	3		
3	C	3	Total	Zn	0	0
			3	3		
3	D	2	Total	Zn	0	0
			2	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Mg	0	0
			3	3		
4	C	2	Total	Mg	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	2	Total	Mg	0	0
			2	2		
4	F	6	Total	Mg	0	0
			6	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	O	0	0
			1	1		
5	C	1	Total	O	0	0
			1	1		
5	D	2	Total	O	0	0
			2	2		
5	E	3	Total	O	0	0
			3	3		
5	F	9	Total	O	0	0
			9	9		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TRANSCRIPTION FACTOR IIIA

Chain B: 




• Molecule 1: TRANSCRIPTION FACTOR IIIA

Chain C: 





• Molecule 1: TRANSCRIPTION FACTOR IIIA

Chain D: 



• Molecule 2: 5S RIBOSOMAL RNA

Chain E: 



• Molecule 2: 5S RIBOSOMAL RNA

Chain F: 

G4	C5	U16	A17	C18	G19	G64	C68	U69	G70	G71	G75	U80	G81	A85	A86	C94	G99	A100	A101	U102	A103	C104	C105	G110	G111	C112	G113	G114	G115
----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	58.60Å 191.59Å 79.77Å 90.00° 101.51° 90.00°	Depositor
Resolution (Å)	35.19 – 3.10 35.19 – 3.10	Depositor EDS
% Data completeness (in resolution range)	97.7 (35.19-3.10) 97.7 (35.19-3.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 3.12Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.216 , 0.259 0.209 , 0.250	Depositor DCC
R_{free} test set	754 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	78.9	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 65.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4554	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.58	1/736 (0.1%)	1.09	3/988 (0.3%)
1	C	0.40	0/744	0.94	4/997 (0.4%)
1	D	0.43	0/490	0.95	0/658
2	E	0.40	0/1458	0.58	0/2273
2	F	0.44	0/1458	0.65	0/2273
All	All	0.45	1/4886 (0.0%)	0.78	7/7189 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	189	GLN	C-N	-5.45	1.25	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	133	LEU	CA-C-N	7.69	129.45	119.84
1	B	133	LEU	C-N-CA	7.69	129.45	119.84
1	B	169	SER	N-CA-C	-7.00	104.89	113.50
1	C	160	ALA	N-CA-C	6.52	118.18	111.14
1	C	130	THR	CB-CA-C	-6.43	109.14	116.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	713	0	679	45	0
1	C	721	0	683	36	0
1	D	475	0	452	33	0
2	E	1304	0	664	35	0
2	F	1304	0	664	22	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	2	0	0	0	0
4	B	3	0	0	0	0
4	C	2	0	0	0	0
4	E	2	0	0	0	0
4	F	6	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	2	0	0	0	0
5	E	3	0	0	0	0
5	F	9	0	0	0	0
All	All	4554	0	3142	163	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 163 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:THR:CG2	1:B:179:LEU:HD12	1.83	1.08
1:B:118:LYS:HB2	1:B:121:GLN:HE21	1.17	1.03
1:B:176:THR:HG23	1:B:179:LEU:HD12	1.03	1.00
1:D:158:VAL:HG12	1:D:159:HIS:H	1.27	0.98
1:C:118:LYS:H	1:C:121:GLN:HE21	1.09	0.93

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	85/87 (98%)	71 (84%)	14 (16%)	0	100	100
1	C	85/87 (98%)	70 (82%)	10 (12%)	5 (6%)	1	8
1	D	56/87 (64%)	40 (71%)	7 (12%)	9 (16%)	0	0
All	All	226/261 (87%)	181 (80%)	31 (14%)	14 (6%)	1	7

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	162	TYR
1	D	134	PRO
1	D	159	HIS
1	D	161	GLY
1	D	169	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	79/80 (99%)	73 (92%)	6 (8%)	12	39
1	C	80/80 (100%)	74 (92%)	6 (8%)	12	39
1	D	53/80 (66%)	50 (94%)	3 (6%)	18	49
All	All	212/240 (88%)	197 (93%)	15 (7%)	13	41

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	131	GLN
1	D	145	ARG
1	C	148	LEU
1	D	148	LEU
1	C	171	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	120	ASN
1	C	121	GLN
1	D	183	HIS
1	D	139	HIS
1	C	108	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	60/61 (98%)	6 (10%)	0
2	F	60/61 (98%)	4 (6%)	0
All	All	120/122 (98%)	10 (8%)	0

5 of 10 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	17	A
2	E	18	C
2	E	19	G
2	E	100	A
2	E	107	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 21 ligands modelled in this entry, 21 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	87/87 (100%)	-0.35	0	100 100	31, 53, 81, 88	0
1	C	87/87 (100%)	0.07	0	100 100	51, 94, 120, 126	0
1	D	58/87 (66%)	0.24	1 (1%)	69 48	62, 95, 136, 139	0
2	E	61/61 (100%)	0.44	0	100 100	66, 135, 159, 165	0
2	F	61/61 (100%)	-0.26	0	100 100	41, 63, 104, 123	0
All	All	354/383 (92%)	0.00	1 (0%)	90 80	31, 82, 144, 165	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	171	SER	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	E	343	1/1	0.57	0.25	83,83,83,83	0
4	MG	E	341	1/1	0.77	0.39	93,93,93,93	0
4	MG	C	344	1/1	0.78	0.15	69,69,69,69	0
4	MG	B	306	1/1	0.85	0.19	83,83,83,83	0
4	MG	F	309	1/1	0.89	0.19	44,44,44,44	0
4	MG	B	308	1/1	0.90	0.17	62,62,62,62	0
4	MG	F	303	1/1	0.92	0.29	61,61,61,61	0
4	MG	F	305	1/1	0.93	0.29	50,50,50,50	0
4	MG	F	304	1/1	0.93	0.16	60,60,60,60	0
4	MG	F	301	1/1	0.94	0.42	54,54,54,54	0
4	MG	F	302	1/1	0.94	0.49	59,59,59,59	0
4	MG	C	342	1/1	0.94	0.32	53,53,53,53	0
3	ZN	C	204	1/1	0.95	0.16	101,101,101,101	0
4	MG	B	307	1/1	0.96	0.07	54,54,54,54	0
3	ZN	B	204	1/1	0.99	0.14	51,51,51,51	0
3	ZN	C	205	1/1	0.99	0.14	75,75,75,75	0
3	ZN	C	206	1/1	0.99	0.15	68,68,68,68	0
3	ZN	D	205	1/1	0.99	0.12	70,70,70,70	0
3	ZN	D	206	1/1	0.99	0.07	103,103,103,103	0
3	ZN	B	206	1/1	0.99	0.14	61,61,61,61	0
3	ZN	B	205	1/1	1.00	0.12	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.