



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 06:17 AM UTC

PDB ID : 3UNW / pdb_00003unw
Title : Crystal Structure of Human GAC in Complex with Glutamate
Authors : DeLaBarre, B.; Gross, S.; Cheng, F.; Gao, Y.; Jha, A.; Jiang, F.; Song, J.J.;
Wie, W.; Hurov, J.
Deposited on : 2011-11-16
Resolution : 2.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1853 (2.58-2.54)
Clashscore	190562	1897 (2.58-2.54)
Ramachandran outliers	187476	1875 (2.58-2.54)
Sidechain outliers	187428	1875 (2.58-2.54)
RSRZ outliers	180081	1853 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	6% 62% 13% • 24%
1	B	534	6% 62% 10% •• 25%
1	C	534	5% 63% 11% • 25%
1	D	534	6% 60% 13% • 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12769 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaminase kidney isoform, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	9	0	0
			3162	2014	532	588	28			
1	B	400	Total	C	N	O	S	6	0	0
			3123	1993	525	577	28			
1	C	400	Total	C	N	O	S	25	0	0
			3124	1995	525	576	28			
1	D	403	Total	C	N	O	S	13	0	0
			3141	2001	528	584	28			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	599	HIS	-	expression tag	UNP O94925
A	600	HIS	-	expression tag	UNP O94925
A	601	HIS	-	expression tag	UNP O94925
A	602	HIS	-	expression tag	UNP O94925
A	603	HIS	-	expression tag	UNP O94925
A	604	HIS	-	expression tag	UNP O94925
B	599	HIS	-	expression tag	UNP O94925
B	600	HIS	-	expression tag	UNP O94925
B	601	HIS	-	expression tag	UNP O94925
B	602	HIS	-	expression tag	UNP O94925
B	603	HIS	-	expression tag	UNP O94925
B	604	HIS	-	expression tag	UNP O94925
C	599	HIS	-	expression tag	UNP O94925
C	600	HIS	-	expression tag	UNP O94925
C	601	HIS	-	expression tag	UNP O94925
C	602	HIS	-	expression tag	UNP O94925
C	603	HIS	-	expression tag	UNP O94925
C	604	HIS	-	expression tag	UNP O94925
D	599	HIS	-	expression tag	UNP O94925
D	600	HIS	-	expression tag	UNP O94925
D	601	HIS	-	expression tag	UNP O94925

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	602	HIS	-	expression tag	UNP O94925
D	603	HIS	-	expression tag	UNP O94925
D	604	HIS	-	expression tag	UNP O94925

- Molecule 2 is GLUTAMIC ACID (CCD ID: GLU) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	5	1	4		
2	B	1	Total	C	N	O	0	0
			10	5	1	4		
2	C	1	Total	C	N	O	0	0
			10	5	1	4		
2	D	1	Total	C	N	O	0	0
			10	5	1	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total	O	0	0
			46	46		
3	B	43	Total	O	0	0
			43	43		
3	C	46	Total	O	0	0
			46	46		
3	D	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutaminase kidney isoform, mitochondrial



- Molecule 1: Glutaminase kidney isoform, mitochondrial





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.66Å 139.61Å 178.38Å 90.00° 94.14° 90.00°	Depositor
Resolution (Å)	29.04 – 2.56 29.04 – 2.56	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.04-2.56) 99.7 (29.04-2.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.226 , 0.261 0.232 , 0.269	Depositor DCC
R_{free} test set	4013 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 28.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12769	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.45	5/3232 (0.2%)	1.45	27/4364 (0.6%)
1	B	1.46	6/3192 (0.2%)	1.45	32/4307 (0.7%)
1	C	1.44	5/3193 (0.2%)	1.40	21/4309 (0.5%)
1	D	1.45	7/3210 (0.2%)	1.44	30/4333 (0.7%)
All	All	1.45	23/12827 (0.2%)	1.44	110/17313 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	10
1	C	0	5
1	D	0	12
All	All	0	34

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	372	GLY	CA-C	-7.10	1.46	1.52
1	B	483	GLY	N-CA	-6.77	1.38	1.45
1	B	501	TRP	NE1-CE2	-6.10	1.30	1.37
1	C	230	HIS	CB-CG	-5.85	1.42	1.50
1	A	541	ASP	N-CA	-5.71	1.41	1.46

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	526	ASN	CA-CB-CG	12.43	125.03	112.60
1	B	306	HIS	CA-CB-CG	-8.30	105.50	113.80
1	B	259	ASP	CA-CB-CG	-7.90	104.70	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	496	MET	CA-C-N	7.76	127.53	122.18
1	D	496	MET	C-N-CA	7.76	127.53	122.18

There are no chirality outliers.

5 of 34 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	169	ARG	Sidechain
1	A	235	TYR	Sidechain
1	A	272	ARG	Sidechain
1	A	382	ARG	Sidechain
1	A	466	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3162	0	3126	18	6
1	B	3123	0	3100	20	1
1	C	3124	0	3103	17	8
1	D	3141	0	3112	19	3
2	A	10	0	5	0	0
2	B	10	0	5	0	0
2	C	10	0	5	0	0
2	D	10	0	5	1	0
3	A	46	0	0	0	0
3	B	43	0	0	0	0
3	C	46	0	0	0	0
3	D	44	0	0	0	1
All	All	12769	0	12461	69	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:532:ASN:HD22	1:A:535:HIS:H	1.18	0.91
1:C:532:ASN:HD22	1:C:535:HIS:H	1.24	0.85
1:A:267:THR:HG22	1:A:269:ASP:H	1.43	0.81
1:D:267:THR:HG22	1:D:269:ASP:H	1.46	0.80
1:B:267:THR:HG22	1:B:269:ASP:H	1.50	0.77

The worst 5 of 12 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:GLU:CD	1:C:538:LYS:NZ[1_455]	0.60	1.60
1:A:236:GLU:OE1	1:C:538:LYS:NZ[1_455]	0.69	1.51
1:C:259:ASP:OD1	1:C:307:ARG:NH2[1_655]	1.32	0.88
1:D:259:ASP:OD1	1:D:307:ARG:NH2[1_455]	1.44	0.76
1:A:236:GLU:OE1	1:C:538:LYS:CE[1_455]	1.54	0.66

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/534 (76%)	384 (95%)	17 (4%)	2 (0%)	24	33
1	B	394/534 (74%)	373 (95%)	16 (4%)	5 (1%)	9	12
1	C	394/534 (74%)	374 (95%)	15 (4%)	5 (1%)	9	12
1	D	399/534 (75%)	383 (96%)	14 (4%)	2 (0%)	24	33
All	All	1590/2136 (74%)	1514 (95%)	62 (4%)	14 (1%)	14	20

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	VAL
1	B	217	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	322	PHE
1	B	527	PHE
1	D	193	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/458 (76%)	328 (94%)	22 (6%)	16	23
1	B	347/458 (76%)	330 (95%)	17 (5%)	22	33
1	C	347/458 (76%)	332 (96%)	15 (4%)	26	38
1	D	349/458 (76%)	327 (94%)	22 (6%)	16	23
All	All	1393/1832 (76%)	1317 (94%)	76 (6%)	20	29

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	152	GLU
1	D	508	MET
1	D	184	LEU
1	D	346	LYS
1	D	539	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	471	GLN
1	D	519	HIS
1	C	532	ASN
1	D	471	GLN
1	C	529	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLU	D	800	-	8,9,9	1.10	1 (12%)	8,11,11	0.85	0
2	GLU	A	800	-	8,9,9	1.19	1 (12%)	8,11,11	1.46	1 (12%)
2	GLU	C	800	-	8,9,9	1.06	1 (12%)	8,11,11	1.27	1 (12%)
2	GLU	B	800	-	8,9,9	1.13	1 (12%)	8,11,11	1.53	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	D	800	-	-	4/9/9/9	-
2	GLU	A	800	-	-	3/9/9/9	-
2	GLU	C	800	-	-	0/9/9/9	-
2	GLU	B	800	-	-	2/9/9/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	800	GLU	OE2-CD	-2.09	1.23	1.30
2	C	800	GLU	OE2-CD	-2.08	1.23	1.30
2	A	800	GLU	OXT-C	-2.06	1.24	1.30
2	B	800	GLU	OE2-CD	-2.05	1.24	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	GLU	CB-CA-C	-3.33	101.62	110.45
2	B	800	GLU	CB-CA-C	-2.88	102.83	110.45
2	C	800	GLU	CB-CA-C	-2.65	103.44	110.45

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	800	GLU	N-CA-CB-CG
2	D	800	GLU	OE1-CD-CG-CB
2	D	800	GLU	OE2-CD-CG-CB
2	A	800	GLU	OE2-CD-CG-CB
2	B	800	GLU	OE2-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	800	GLU	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	407/534 (76%)	0.55	30 (7%)	20 20	24, 39, 80, 96	2 (0%)
1	B	400/534 (74%)	0.54	34 (8%)	16 16	24, 39, 78, 89	1 (0%)
1	C	400/534 (74%)	0.56	27 (6%)	23 23	24, 39, 78, 92	6 (1%)
1	D	403/534 (75%)	0.55	33 (8%)	17 17	24, 39, 78, 89	3 (0%)
All	All	1610/2136 (75%)	0.55	124 (7%)	19 19	24, 39, 78, 96	12 (0%)

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	321	LEU	8.4
1	C	136	LEU	7.3
1	C	193	VAL	5.9
1	A	321	LEU	5.7
1	C	322	PHE	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLU	C	800	10/10	0.81	0.14	48,64,66,66	0
2	GLU	D	800	10/10	0.81	0.18	61,69,70,71	0
2	GLU	A	800	10/10	0.83	0.16	60,63,64,64	0
2	GLU	B	800	10/10	0.84	0.13	49,60,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.