



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:13 PM UTC

PDB ID : 4UNA / pdb_00004una
Title : THE CRYSTAL STRUCTURE OF I-DMOI IN COMPLEX WITH ITS TARGET DNA AT 2 DAYS INCUBATION IN 5MM MN (STATE 4)
Authors : Molina, R.; Stella, S.; Redondo, P.; Gomez, H.; Marcaida, M.J.; Orozco, M.; Prieto, J.; Montoya, G.
Deposited on : 2014-05-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	8% 71% 18% • • 8%
1	D	199	10% 79% 13% • • 8%
1	G	199	3% 67% 20% • 12%
2	B	25	76% 24%
2	E	25	84% 16%
2	H	25	4% 80% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	15	 87%13%
3	F	15	 67%33%
3	I	15	 87%13%
4	K	10	 100%
4	M	10	 100%
4	O	10	 60%40%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HOMING ENDONUCLEASE I-DMOI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	184	Total	C	N	O	S	0	0	0
			1516	977	275	261	3			
1	D	184	Total	C	N	O	S	0	1	0
			1523	984	275	261	3			
1	G	176	Total	C	N	O	S	0	4	0
			1481	958	267	253	3			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	expression tag	UNP P21505
A	189	ALA	-	expression tag	UNP P21505
A	190	ALA	-	expression tag	UNP P21505
A	191	ALA	-	expression tag	UNP P21505
A	192	LEU	-	expression tag	UNP P21505
A	193	GLU	-	expression tag	UNP P21505
A	194	HIS	-	expression tag	UNP P21505
A	195	HIS	-	expression tag	UNP P21505
A	196	HIS	-	expression tag	UNP P21505
A	197	HIS	-	expression tag	UNP P21505
A	198	HIS	-	expression tag	UNP P21505
A	199	HIS	-	expression tag	UNP P21505
D	1	ALA	-	expression tag	UNP P21505
D	189	ALA	-	expression tag	UNP P21505
D	190	ALA	-	expression tag	UNP P21505
D	191	ALA	-	expression tag	UNP P21505
D	192	LEU	-	expression tag	UNP P21505
D	193	GLU	-	expression tag	UNP P21505
D	194	HIS	-	expression tag	UNP P21505
D	195	HIS	-	expression tag	UNP P21505
D	196	HIS	-	expression tag	UNP P21505
D	197	HIS	-	expression tag	UNP P21505
D	198	HIS	-	expression tag	UNP P21505

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	199	HIS	-	expression tag	UNP P21505
G	1	ALA	-	expression tag	UNP P21505
G	189	ALA	-	expression tag	UNP P21505
G	190	ALA	-	expression tag	UNP P21505
G	191	ALA	-	expression tag	UNP P21505
G	192	LEU	-	expression tag	UNP P21505
G	193	GLU	-	expression tag	UNP P21505
G	194	HIS	-	expression tag	UNP P21505
G	195	HIS	-	expression tag	UNP P21505
G	196	HIS	-	expression tag	UNP P21505
G	197	HIS	-	expression tag	UNP P21505
G	198	HIS	-	expression tag	UNP P21505
G	199	HIS	-	expression tag	UNP P21505

- Molecule 2 is a DNA chain called 25MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	25	Total	C	N	O	P	0	0	0
			511	242	94	151	24			
2	E	25	Total	C	N	O	P	0	0	0
			511	242	94	151	24			
2	H	25	Total	C	N	O	P	0	0	0
			511	242	94	151	24			

- Molecule 3 is a DNA chain called 5'-D(*CP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*T
P*TP*AP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			
3	F	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			
3	I	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			

- Molecule 4 is a DNA chain called 5'-D(*CP*CP*GP*GP*CP*AP*AP*GP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	K	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			
4	M	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	O	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			

- Molecule 5 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Mn	0	0
			3	3		
5	D	3	Total	Mn	0	0
			3	3		
5	G	3	Total	Mn	0	0
			3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	35	Total	O	0	0
			35	35		
6	B	8	Total	O	0	0
			8	8		
6	C	2	Total	O	0	0
			2	2		
6	D	30	Total	O	0	0
			30	30		
6	E	5	Total	O	0	0
			5	5		
6	F	3	Total	O	0	0
			3	3		
6	G	30	Total	O	0	0
			30	30		
6	H	3	Total	O	0	0
			3	3		
6	I	2	Total	O	0	0
			2	2		
6	M	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HOMING ENDONUCLEASE I-DMOI




• Molecule 1: HOMING ENDONUCLEASE I-DMOI



• Molecule 1: HOMING ENDONUCLEASE I-DMOI




• Molecule 2: 25MER



- Molecule 2: 25MER

Chain E:  84% 16%



- Molecule 2: 25MER

Chain H:  4% 80% 20%



- Molecule 3: 5'-D(*CP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*AP*C)-3'

Chain C:  87% 13%



- Molecule 3: 5'-D(*CP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*AP*C)-3'

Chain F:  67% 33%




- Molecule 3: 5'-D(*CP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*AP*C)-3'

Chain I:  87% 13%



- Molecule 4: 5'-D(*CP*CP*GP*GP*CP*AP*AP*GP*GP*C)-3'

Chain K:  100%

There are no outlier residues recorded for this chain.


- Molecule 4: 5'-D(*CP*CP*GP*GP*CP*AP*AP*GP*GP*C)-3'

Chain M:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: 5'-D(*CP*CP*GP*GP*CP*AP*AP*GP*GP*C)-3'

Chain O:  60% 40%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.52Å 70.15Å 106.70Å 90.00° 119.83° 90.00°	Depositor
Resolution (Å)	46.28 – 2.30 46.28 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.3 (46.28-2.30) 96.7 (46.28-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.29Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.190 , 0.219 0.201 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	54.9	Xtriage
Anisotropy	0.490	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for -h-l,k,h 0.020 for l,k,-h-l 0.015 for h,-k,-h-l 0.017 for -h-l,-k,l 0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7708	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/1543	0.86	3/2076 (0.1%)
1	D	0.53	0/1554	0.84	2/2092 (0.1%)
1	G	0.58	1/1518 (0.1%)	0.81	0/2038
2	B	0.45	0/572	0.71	0/882
2	E	0.41	0/572	0.65	0/882
2	H	0.49	0/572	0.68	0/882
3	C	0.40	0/338	0.60	0/519
3	F	0.46	0/338	0.61	0/519
3	I	0.45	0/338	0.66	0/519
4	K	0.36	0/232	0.50	0/354
4	M	0.34	0/232	0.48	0/354
4	O	0.37	0/232	0.52	0/354
All	All	0.51	1/8041 (0.0%)	0.75	5/11471 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	18	ILE	CA-CB	5.41	1.60	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	5	GLU	N-CA-C	-7.07	104.27	113.17
1	A	187	PRO	N-CA-CB	5.94	109.53	103.00
1	A	186	PRO	N-CA-C	-5.40	104.11	110.70
1	D	77	ARG	CB-CG-CD	5.33	123.56	111.30
1	A	184	PRO	CA-C-O	-5.14	111.24	120.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1516	0	1586	22	1
1	D	1523	0	1603	18	0
1	G	1481	0	1568	26	0
2	B	511	0	282	8	0
2	E	511	0	282	4	1
2	H	511	0	282	3	0
3	C	302	0	169	1	0
3	F	302	0	168	3	0
3	I	302	0	169	2	0
4	K	207	0	111	0	0
4	M	207	0	111	0	0
4	O	207	0	111	3	0
5	A	3	0	0	0	0
5	D	3	0	0	0	0
5	G	3	0	0	0	0
6	A	35	0	0	1	0
6	B	8	0	0	2	0
6	C	2	0	0	0	0
6	D	30	0	0	0	0
6	E	5	0	0	0	0
6	F	3	0	0	0	0
6	G	30	0	0	2	0
6	H	3	0	0	0	0
6	I	2	0	0	0	0
6	M	1	0	0	0	0
All	All	7708	0	6442	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:5:DC:H2''	3:C:6:DC:H5''	1.58	0.83
1:D:4:ASN:HA	1:D:7:VAL:HG13	1.59	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:33:ARG:NH1	2:E:21:DG:O6	2.15	0.79
2:H:23:DG:H2''	2:H:24:DC:H5''	1.67	0.77
2:B:22:DC:P	1:G:97:ARG:HH22	2.11	0.73
1:A:75:ASP:HB3	1:A:76:THR:HG23	1.72	0.72
1:D:60[A]:ILE:HD11	1:D:87:LEU:HD22	1.71	0.72
1:A:19:ILE:HD11	1:A:95:LEU:HD13	1.71	0.71
2:H:24:DC:H2'	2:H:25:DG:C8	2.25	0.71
2:B:12:DT:O2	6:B:2003:HOH:O	2.08	0.71
1:G:172:LYS:O	1:G:176:THR:OG1	2.11	0.69
1:D:181:HIS:HA	1:D:185:LEU:HD11	1.74	0.69
1:A:41:THR:HB	1:A:77:ARG:HD2	1.75	0.67
3:I:7:DG:H2''	3:I:8:DG:H5''	1.78	0.65
1:A:180:SER:HB2	1:A:182:LEU:HD13	1.79	0.65
3:F:2:DG:H2''	3:F:3:DC:H5''	1.77	0.65
1:D:77:ARG:CG	1:D:77:ARG:HH11	2.13	0.61
1:G:152[B]:HIS:HD2	6:G:2026:HOH:O	1.84	0.60
1:A:77:ARG:NH2	2:B:15:DG:O6	2.34	0.60
1:D:77:ARG:HH11	1:D:77:ARG:HG2	1.66	0.59
1:D:30:LYS:O	1:D:33:ARG:NE	2.30	0.58
1:A:29:TYR:HB3	1:A:33:ARG:NH2	2.19	0.58
1:G:118:GLY:O	1:G:120:LYS:HE3	2.05	0.56
1:G:46:ASN:ND2	1:G:50:GLN:OE1	2.37	0.56
1:G:121:THR:HB	1:G:123:LYS:HE3	1.87	0.56
1:G:66:LYS:O	1:G:66:LYS:HG2	2.07	0.53
1:A:45:GLU:HB2	1:A:78:TYR:CE2	2.44	0.53
1:G:22:GLY:HA3	1:G:39:VAL:O	2.09	0.53
1:A:118:GLY:O	1:A:120:LYS:HE3	2.10	0.52
1:A:184:PRO:O	1:A:185:LEU:C	2.52	0.52
3:F:10:DA:H1'	3:F:11:DC:H5'	1.92	0.51
1:A:152:HIS:HB3	6:A:2031:HOH:O	2.11	0.50
2:E:19:DC:H2''	2:E:20:DG:H5'	1.93	0.50
2:B:4:DT:OP2	6:B:2001:HOH:O	2.19	0.49
1:G:19[A]:ILE:HD11	1:G:95:LEU:HD22	1.94	0.48
1:A:76:THR:OG1	2:B:14:DA:N7	2.45	0.48
1:G:132:LYS:O	1:G:136[B]:GLU:HG2	2.13	0.48
1:G:128:TRP:CZ3	4:O:16:DC:H2'	2.48	0.48
1:G:175:HIS:HA	1:G:179:SER:HB2	1.94	0.48
2:H:3:DC:H2'	2:H:4:DT:H71	1.96	0.47
1:D:73:LYS:HE2	1:D:78:TYR:CE2	2.50	0.47
1:A:107:ILE:HG23	1:A:177:ILE:HD13	1.95	0.47
1:G:104:ARG:NH1	6:G:2016:HOH:O	2.25	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:33:ARG:HD2	3:F:4:DG:O6	2.14	0.46
2:B:22:DC:OP2	1:G:97:ARG:NH2	2.48	0.46
1:D:155:ASP:HB3	1:D:160:VAL:HB	1.98	0.46
1:G:30:LYS:O	1:G:33:ARG:NE	2.43	0.46
2:E:20:DG:H2''	2:E:21:DG:O5'	2.16	0.45
1:D:22:GLY:HA3	1:D:39:VAL:O	2.17	0.45
1:D:76:THR:OG1	2:E:14:DA:N7	2.47	0.45
1:D:77:ARG:CG	1:D:77:ARG:NH1	2.77	0.45
1:A:180:SER:HB2	1:A:182:LEU:CD1	2.47	0.44
2:B:22:DC:P	1:G:97:ARG:NH2	2.87	0.44
1:A:41:THR:CB	1:A:77:ARG:HD2	2.45	0.44
1:D:111:LYS:HD2	1:D:186:PRO:HD3	2.00	0.44
3:I:7:DG:C2'	3:I:8:DG:H5''	2.46	0.44
1:G:155:ASP:HB3	1:G:160:VAL:HB	2.01	0.43
1:A:42:GLN:HB3	1:A:48:ILE:HG12	1.99	0.43
4:O:21:DA:H2''	4:O:22:DA:C8	2.53	0.43
4:O:20:DC:H2''	4:O:21:DA:C8	2.53	0.43
1:A:167:LEU:O	1:A:170:ARG:HB2	2.19	0.42
1:G:60:ILE:HG23	1:G:65:VAL:HB	2.02	0.42
1:A:115:VAL:O	1:A:120:LYS:HE2	2.20	0.42
1:A:43:LYS:NZ	2:B:13:DA:OP2	2.30	0.42
1:A:22:GLY:HA3	1:A:39:VAL:O	2.19	0.42
1:A:132:LYS:HD2	1:A:161:TYR:CZ	2.55	0.42
1:D:31:GLY:HA2	1:D:32:ASN:HA	1.87	0.42
1:A:111:LYS:O	1:A:115:VAL:HG23	2.20	0.41
1:G:42:GLN:HG2	1:G:47:LEU:HD23	2.03	0.41
1:G:153:LEU:HD23	1:G:156:HIS:CE1	2.55	0.41
1:G:87:LEU:HD12	1:G:87:LEU:HA	1.85	0.41
1:G:35:GLU:CD	1:G:37:ARG:HE	2.28	0.41
1:D:73:LYS:HE2	1:D:78:TYR:CZ	2.56	0.41
1:G:121:THR:C	1:G:123:LYS:H	2.28	0.41
1:G:167:LEU:HD23	1:G:167:LEU:HA	1.79	0.41
1:D:155:ASP:OD2	1:D:158:HIS:HD2	2.04	0.41
1:A:49:LYS:HG2	1:A:71:ILE:HD13	2.03	0.40
1:D:121:THR:C	1:D:123:LYS:H	2.29	0.40
1:G:163:LEU:C	1:G:163:LEU:HD23	2.46	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:ARG:NH2	2:E:19:DC:OP1[2_10410]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/199 (92%)	173 (95%)	7 (4%)	2 (1%)	11	13
1	D	183/199 (92%)	177 (97%)	6 (3%)	0	100	100
1	G	178/199 (89%)	172 (97%)	5 (3%)	1 (1%)	21	27
All	All	543/597 (91%)	522 (96%)	18 (3%)	3 (1%)	21	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	ASP
1	G	122	LEU
1	A	184	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/178 (93%)	153 (92%)	13 (8%)	11	16
1	D	168/178 (94%)	163 (97%)	5 (3%)	36	53
1	G	163/178 (92%)	156 (96%)	7 (4%)	26	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	497/534 (93%)	472 (95%)	25 (5%)	21	33

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ILE
1	A	27	LEU
1	A	30	LYS
1	A	64	ASN
1	A	66	LYS
1	A	68	LYS
1	A	73	LYS
1	A	77	ARG
1	A	98	ILE
1	A	120	LYS
1	A	129	ASN
1	A	182	LEU
1	A	186	PRO
1	D	7	VAL
1	D	48	ILE
1	D	66	LYS
1	D	77	ARG
1	D	165	ILE
1	G	5	GLU
1	G	48	ILE
1	G	104	ARG
1	G	120	LYS
1	G	129	ASN
1	G	165	ILE
1	G	176	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	D	42	GLN
1	D	152	HIS
1	D	158	HIS
1	G	156	HIS
1	G	158	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	184/199 (92%)	0.66	16 (8%)	16	17	39, 54, 79, 96	0
1	D	184/199 (92%)	0.71	19 (10%)	12	13	32, 56, 84, 100	1 (0%)
1	G	176/199 (88%)	0.33	6 (3%)	48	50	25, 47, 69, 82	4 (2%)
2	B	25/25 (100%)	0.03	0	100	100	46, 62, 71, 75	0
2	E	25/25 (100%)	0.06	0	100	100	46, 63, 75, 81	0
2	H	25/25 (100%)	-0.01	1 (4%)	42	44	47, 58, 73, 77	0
3	C	15/15 (100%)	0.03	0	100	100	45, 54, 78, 79	0
3	F	15/15 (100%)	-0.06	0	100	100	47, 56, 69, 72	0
3	I	15/15 (100%)	-0.14	0	100	100	44, 54, 75, 76	0
4	K	10/10 (100%)	0.17	0	100	100	51, 75, 78, 79	0
4	M	10/10 (100%)	0.46	0	100	100	52, 81, 90, 90	0
4	O	10/10 (100%)	0.17	0	100	100	47, 68, 84, 85	0
All	All	694/747 (92%)	0.46	42 (6%)	27	29	25, 55, 80, 100	5 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	HIS	4.9
1	A	187	PRO	4.8
1	A	182	LEU	4.3
1	D	187	PRO	4.2
1	G	31	GLY	4.2
1	D	182	LEU	4.0
1	D	31	GLY	3.9
1	D	181	HIS	3.9
1	D	7	VAL	3.8
1	A	186	PRO	3.7
1	A	183	ASN	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6	ASN	3.5
1	D	183	ASN	3.3
1	A	4	ASN	3.0
1	A	184	PRO	2.9
1	D	122	LEU	2.9
1	G	75	ASP	2.9
1	A	185	LEU	2.8
1	A	77	ARG	2.8
1	D	156	HIS	2.7
1	D	184	PRO	2.6
1	G	175	HIS	2.6
1	A	7	VAL	2.5
2	H	1	DG	2.5
1	D	185	LEU	2.5
1	D	186	PRO	2.5
1	G	179	SER	2.4
1	D	176	THR	2.4
1	G	66	LYS	2.4
1	D	180	SER	2.3
1	A	75	ASP	2.3
1	D	152	HIS	2.2
1	A	175	HIS	2.2
1	A	66	LYS	2.2
1	A	73	LYS	2.2
1	G	5	GLU	2.2
1	D	171	ILE	2.1
1	D	4	ASN	2.1
1	D	5	GLU	2.1
1	D	177	ILE	2.1
1	A	144	ASN	2.1
1	D	75	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MN	D	1191	1/1	0.97	0.07	54,54,54,54	1
5	MN	A	1190	1/1	0.98	0.05	53,53,53,53	1
5	MN	G	1182	1/1	0.98	0.09	54,54,54,54	1
5	MN	D	1189	1/1	0.99	0.05	48,48,48,48	0
5	MN	A	1189	1/1	0.99	0.04	48,48,48,48	0
5	MN	G	1180	1/1	0.99	0.04	42,42,42,42	0
5	MN	G	1181	1/1	0.99	0.03	43,43,43,43	0
5	MN	A	1188	1/1	0.99	0.05	53,53,53,53	0
5	MN	D	1190	1/1	1.00	0.07	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.