



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 02:59 PM UTC

PDB ID : 5UNY / pdb_00005uny
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with (RS)-3-[(2-amino-4-methylquinolin-7-yl)methoxy]-5-(2-(methylamino)propyl) benzonitrile
Authors : Li, H.; Poulos, T.L.
Deposited on : 2017-01-31
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	4% 86% 10% ..
1	B	422	3% 90% 7% .

i


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 3-[(2-amino-4-methylquinolin-7-yl)methoxy]-5-[(2R)-2-(methylamino)propyl]benzonitrile (CCD ID: M4R) (formula: $C_{22}H_{24}N_4O$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			27	22	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	1
			27	22	4	1		

- Molecule 5 is 3-[(2-amino-4-methylquinolin-7-yl)methoxy]-5-[(2S)-2-(methylamino)propyl]benzonitrile (CCD ID: 8FD) (formula: C₂₂H₂₄N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	1
			27	22	4	1		
5	B	1	Total	C	N	O	0	1
			27	22	4	1		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	175	Total	O	0	0
			175	175		
8	B	231	Total	O	0	0
			231	231		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.97Å 111.94Å 164.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.23 – 1.82 39.23 – 1.82	Depositor EDS
% Data completeness (in resolution range)	91.6 (39.23-1.82) 92.7 (39.23-1.82)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.193 , 0.238 0.196 , 0.236	Depositor DCC
R_{free} test set	4013 reflections (3.96%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 60.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7354	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: H4B, 8FD, M4R, ZN, HEM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/3456	0.79	0/4688
1	B	0.47	0/3459	0.77	0/4689
All	All	0.46	0/6915	0.78	0/9377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3354	0	3264	29	0
1	B	3357	0	3275	19	0
2	A	43	0	30	3	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	27	0	0	1	0
4	B	27	0	0	1	0
5	A	27	0	0	1	0
5	B	27	0	0	1	0
6	A	4	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	4	0	3	0	0
7	A	1	0	0	0	0
8	A	175	0	0	1	0
8	B	231	0	0	3	0
All	All	7354	0	6635	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:801:HEM:HBB2	2:A:801:HEM:HHC	1.69	0.74
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.69	0.72
2:B:801:HEM:HMC2	2:B:801:HEM:HBC2	1.74	0.69
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.75	0.69
1:A:516:ARG:NH2	8:A:901:HOH:O	2.25	0.67
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.78	0.66
1:B:706:TYR:OH	2:B:801:HEM:O2D	2.11	0.65
1:A:467:ASP:N	1:A:467:ASP:OD1	2.28	0.64
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.80	0.63
1:A:706:TYR:OH	2:A:801:HEM:O2D	2.17	0.61
1:B:667:ARG:NH1	1:B:668:CYS:HB3	2.20	0.56
1:B:516:ARG:NH1	8:B:907:HOH:O	2.40	0.54
1:A:491:SER:OG	1:A:492:THR:N	2.39	0.54
1:A:660:LYS:O	1:A:663[B]:GLU:HG3	2.08	0.52
1:B:607:LEU:HD13	1:B:626:LYS:HG2	1.93	0.50
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.94	0.50
1:B:595:VAL:O	1:B:599:CYS:HB2	2.12	0.49
1:A:330:ILE:HD11	1:B:696:LEU:HD22	1.94	0.49
1:B:600:ASP:OD2	8:B:901:HOH:O	2.20	0.49
1:B:567:VAL:HG21	4:B:803[A]:M4R:C07	2.43	0.48
1:B:508:GLN:NE2	8:B:910:HOH:O	2.47	0.47
1:B:322:LEU:HD13	1:B:699:ARG:NH2	2.30	0.46
1:A:359:LYS:HE2	1:A:381:LEU:HD21	1.98	0.46
1:A:595:VAL:O	1:A:599:CYS:HB2	2.16	0.46
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.98	0.45
1:A:496:PRO:HA	1:A:499:VAL:HG23	1.97	0.45
1:B:701:THR:HA	1:B:702:PRO:C	2.41	0.45
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.99	0.45
1:B:567:VAL:HG21	5:B:804[B]:8FD:C07	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:373:GLY:HA2	1:A:378:MET:HE3	1.99	0.44
1:B:323:GLU:O	1:B:699:ARG:HD3	2.17	0.44
1:A:553:TRP:HB2	1:A:613:LYS:NZ	2.33	0.44
1:A:546:ILE:HG22	1:A:554:PHE:HE2	1.83	0.44
1:A:567:VAL:HG21	4:A:803[A]:M4R:C07	2.48	0.44
1:B:336:MET:HE2	1:B:336:MET:HB2	1.85	0.44
1:A:567:VAL:HG21	5:A:804[B]:8FD:C07	2.48	0.43
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.53	0.43
1:A:360:GLU:O	1:A:364:GLN:HG2	2.18	0.43
1:A:475:TRP:CE2	1:A:710:PRO:HB2	2.53	0.43
1:B:373:GLY:HA2	1:B:378:MET:HE2	1.99	0.43
1:A:684:SER:HB3	1:A:687:ILE:HD11	2.00	0.43
1:B:366:TYR:HA	1:B:369:ILE:HG12	2.01	0.42
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.54	0.42
1:A:596:ARG:NH2	1:A:600:ASP:OD2	2.51	0.42
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.08	0.41
1:A:546:ILE:HG12	1:A:560:LYS:HA	2.03	0.41
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.87	0.41
1:A:467:ASP:C	1:A:469:LYS:H	2.29	0.41
1:A:328:GLU:H	1:A:328:GLU:HG3	1.62	0.41
1:A:537:PRO:HA	1:A:538:PRO:HD3	1.93	0.41
1:A:686:SER:HA	1:A:691:PHE:CG	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/422 (97%)	391 (96%)	18 (4%)	0	100	100
1	B	410/422 (97%)	399 (97%)	11 (3%)	0	100	100
All	All	819/844 (97%)	790 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/377 (98%)	360 (98%)	9 (2%)	43	27
1	B	369/377 (98%)	363 (98%)	6 (2%)	55	44
All	All	738/754 (98%)	723 (98%)	15 (2%)	48	35

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	337	LEU
1	A	348	VAL
1	A	381	LEU
1	A	467	ASP
1	A	503	GLU
1	A	523	LEU
1	A	527	ASN
1	A	613	LYS
1	A	715	VAL
1	B	321	THR
1	B	348	VAL
1	B	353	GLN
1	B	360	GLU
1	B	516	ARG
1	B	667	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	420	GLN
1	A	436	HIS
1	A	712	ASN
1	B	535	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	H4B	A	802	-	17,18,18	0.81	0	14,26,26	1.94	4 (28%)
2	HEM	A	801	1	50,50,50	2.10	10 (20%)	67,82,82	1.31	7 (10%)
5	8FD	A	804[B]	-	29,29,29	1.61	1 (3%)	38,40,40	1.24	4 (10%)
5	8FD	B	804[B]	-	29,29,29	1.65	1 (3%)	38,40,40	1.17	3 (7%)
6	ACT	B	805	-	3,3,3	0.88	0	3,3,3	0.56	0
3	H4B	B	802	-	17,18,18	0.85	0	14,26,26	1.89	4 (28%)
4	M4R	B	803[A]	-	29,29,29	1.65	2 (6%)	38,40,40	1.10	3 (7%)
4	M4R	A	803[A]	-	29,29,29	1.61	1 (3%)	38,40,40	1.17	3 (7%)
2	HEM	B	801	1	50,50,50	2.06	10 (20%)	67,82,82	1.37	6 (8%)
6	ACT	A	805	-	3,3,3	0.82	0	3,3,3	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
2	HEM	A	801	1	-	8/14/54/54	-
5	8FD	A	804[B]	-	-	1/13/13/13	0/3/3/3
5	8FD	B	804[B]	-	-	1/13/13/13	0/3/3/3
3	H4B	B	802	-	-	1/8/17/17	0/2/2/2
4	M4R	B	803[A]	-	-	1/13/13/13	0/3/3/3
4	M4R	A	803[A]	-	-	2/13/13/13	0/3/3/3
2	HEM	B	801	1	-	2/14/54/54	-

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	C3D-C2D	7.80	1.53	1.36
2	A	801	HEM	C3D-C2D	7.72	1.53	1.36
5	B	804[B]	8FD	C23-C27	-7.41	1.28	1.44
4	B	803[A]	M4R	C23-C27	-7.40	1.28	1.44
5	A	804[B]	8FD	C23-C27	-7.36	1.29	1.44
4	A	803[A]	M4R	C23-C27	-7.27	1.29	1.44
2	A	801	HEM	FE-NA	6.59	2.16	1.95
2	B	801	HEM	FE-NA	5.87	2.14	1.95
2	A	801	HEM	FE-NC	4.69	2.10	1.95
2	B	801	HEM	FE-NC	4.06	2.08	1.95
2	A	801	HEM	FE-NB	4.05	2.07	1.94
2	B	801	HEM	FE-ND	3.95	2.07	1.94
2	B	801	HEM	FE-NB	3.77	2.06	1.94
2	A	801	HEM	CAB-C3B	3.17	1.55	1.47
2	A	801	HEM	FE-ND	2.98	2.04	1.94
2	B	801	HEM	CAC-C3C	2.96	1.55	1.47
2	B	801	HEM	CAB-C3B	2.95	1.55	1.47
2	A	801	HEM	CAC-C3C	2.90	1.55	1.47
2	B	801	HEM	CMC-C2C	2.58	1.56	1.50
2	B	801	HEM	CMB-C2B	2.58	1.56	1.50
2	B	801	HEM	CMD-C2D	2.49	1.55	1.50
2	A	801	HEM	CMC-C2C	2.48	1.55	1.50
2	A	801	HEM	CMD-C2D	2.47	1.55	1.50
2	A	801	HEM	CMA-C3A	2.13	1.55	1.50
4	B	803[A]	M4R	C09-C10	-2.06	1.38	1.41

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	C4D-ND-C1D	5.08	111.23	105.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	H4B	C2-N1-C8A	4.49	121.28	113.36
2	A	801	HEM	C4D-ND-C1D	4.39	110.40	105.21
3	B	802	H4B	C2-N1-C8A	4.31	120.97	113.36
5	B	804[B]	8FD	C35-N34-C32	-3.75	109.91	114.39
5	A	804[B]	8FD	C35-N34-C32	-3.49	110.22	114.39
2	B	801	HEM	C2A-C1A-NA	-3.39	106.39	110.15
4	A	803[A]	M4R	C33-N32-C30	-3.34	110.40	114.39
4	B	803[A]	M4R	C33-N32-C30	-3.24	110.52	114.39
2	A	801	HEM	C3B-C2B-C1B	3.08	108.72	106.41
5	A	804[B]	8FD	C04-C05-C10	2.94	119.79	118.00
4	A	803[A]	M4R	C04-C05-C10	2.77	119.69	118.00
3	B	802	H4B	C2-N3-C4	-2.72	120.17	125.11
3	A	802	H4B	O4-C4-C4A	-2.72	120.70	127.26
5	A	804[B]	8FD	C05-C10-N01	-2.72	119.92	122.80
3	B	802	H4B	C4A-C4-N3	2.71	119.57	112.13
2	A	801	HEM	CHD-C1D-ND	2.71	127.34	124.42
3	A	802	H4B	C2-N3-C4	-2.63	120.35	125.11
3	A	802	H4B	C4A-C4-N3	2.61	119.30	112.13
2	B	801	HEM	CHD-C1D-ND	2.54	127.16	124.42
5	B	804[B]	8FD	C04-C05-C10	2.54	119.55	118.00
3	B	802	H4B	O4-C4-C4A	-2.54	121.14	127.26
2	B	801	HEM	CAD-C3D-C4D	2.52	129.09	124.70
5	A	804[B]	8FD	C12-O13-C21	2.49	123.54	117.62
2	A	801	HEM	CHA-C4D-ND	2.47	127.42	124.37
2	A	801	HEM	CMD-C2D-C1D	2.41	128.79	125.03
4	B	803[A]	M4R	C04-C05-C10	2.40	119.47	118.00
4	A	803[A]	M4R	C05-C10-N01	-2.33	120.33	122.80
5	B	804[B]	8FD	C05-C10-N01	-2.23	120.44	122.80
2	A	801	HEM	C2A-C1A-NA	-2.19	107.72	110.15
2	B	801	HEM	CAD-CBD-CGD	-2.15	107.97	113.67
4	B	803[A]	M4R	C05-C10-N01	-2.09	120.59	122.80
2	A	801	HEM	CAD-CBD-CGD	-2.03	108.28	113.67
2	B	801	HEM	CBA-CAA-C2A	-2.03	106.92	112.53

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	803[A]	M4R	C25-C29-C30-C31
4	B	803[A]	M4R	C25-C29-C30-C31
2	A	801	HEM	C2D-C3D-CAD-CBD
2	A	801	HEM	C2A-CAA-CBA-CGA

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	801	HEM	C4D-C3D-CAD-CBD
2	A	801	HEM	C1A-C2A-CAA-CBA
2	B	801	HEM	C2A-CAA-CBA-CGA
2	A	801	HEM	C4B-C3B-CAB-CBB
2	B	801	HEM	C4B-C3B-CAB-CBB
3	B	802	H4B	C7-C6-C9-C10
4	A	803[A]	M4R	C25-C29-C30-N32
2	A	801	HEM	C3A-C2A-CAA-CBA
2	A	801	HEM	CAD-CBD-CGD-O2D
2	A	801	HEM	CAD-CBD-CGD-O1D
5	A	804[B]	8FD	C22-C23-C27-N28
5	B	804[B]	8FD	C22-C23-C27-N28


There are no ring outliers.

6 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	HEM	3	0
5	A	804[B]	8FD	1	0
5	B	804[B]	8FD	1	0
4	B	803[A]	M4R	1	0
4	A	803[A]	M4R	1	0
2	B	801	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.








Ligand 8FD B 804 (B)



Ligand M4R B 803 (A)







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	410/422 (97%)	0.50	17 (4%)	41 47	30, 61, 115, 162	3 (0%)
1	B	411/422 (97%)	0.22	11 (2%)	56 61	29, 50, 87, 117	3 (0%)
All	All	821/844 (97%)	0.36	28 (3%)	48 55	29, 55, 108, 162	6 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	VAL	4.7
1	B	338	PRO	4.2
1	A	338	PRO	3.4
1	A	716	TRP	3.3
1	A	300	PHE	3.0
1	A	599	CYS	3.0
1	A	355	PHE	2.9
1	A	715	VAL	2.8
1	A	490	GLY	2.8
1	A	348	VAL	2.8
1	A	606	ILE	2.5
1	B	706	TYR	2.4
1	B	300	PHE	2.4
1	B	360	GLU	2.4
1	B	321	THR	2.4
1	A	595	VAL	2.4
1	B	337	LEU	2.3
1	A	604	TYR	2.3
1	A	504	ILE	2.2
1	A	515	GLY	2.2
1	B	599	CYS	2.2
1	A	711	TRP	2.2
1	B	604	TYR	2.1
1	B	594	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	322	LEU	2.1
1	A	706	TYR	2.1
1	B	336	MET	2.1
1	A	598	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	8FD	B	804[B]	27/27	0.84	0.17	32,52,82,84	27
6	ACT	A	805	4/4	0.85	0.15	56,62,63,63	0
4	M4R	B	803[A]	27/27	0.86	0.16	32,52,83,85	27
4	M4R	A	803[A]	27/27	0.88	0.15	32,56,84,88	27
3	H4B	B	802	17/17	0.90	0.10	36,45,52,53	0
5	8FD	A	804[B]	27/27	0.91	0.15	33,54,84,87	27
3	H4B	A	802	17/17	0.92	0.10	32,47,59,65	0
6	ACT	B	805	4/4	0.94	0.10	49,56,57,58	0
2	HEM	A	801	43/43	0.98	0.08	26,40,71,74	0
2	HEM	B	801	43/43	0.98	0.08	28,40,62,65	0
7	ZN	A	806	1/1	0.99	0.03	42,42,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 8FD B 804 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around M4R B 803 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around M4R A 803 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 8FD A 804 (B):**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.