



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 01:08 PM UTC

PDB ID : 6UN6 / pdb_00006un6
Title : Crystal structure of green fluorescent protein (GFP); S65T, Y66(3-NO₂Y); ih circular permutant (50-51)
Authors : Lin, C.-Y.; Boxer, S.G.
Deposited on : 2019-10-10
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	
1	B	252	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	18	0
			1881	1201	309	367	4			
1	B	231	Total	C	N	O	S	0	11	0
			1879	1193	318	364	4			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	initiating methionine	UNP A0A059PIQ0
A	-9	GLY	-	expression tag	UNP A0A059PIQ0
A	-8	HIS	-	expression tag	UNP A0A059PIQ0
A	-7	HIS	-	expression tag	UNP A0A059PIQ0
A	-6	HIS	-	expression tag	UNP A0A059PIQ0
A	-5	HIS	-	expression tag	UNP A0A059PIQ0
A	-4	HIS	-	expression tag	UNP A0A059PIQ0
A	-3	HIS	-	expression tag	UNP A0A059PIQ0
A	-2	SER	-	expression tag	UNP A0A059PIQ0
A	-1	SER	-	expression tag	UNP A0A059PIQ0
A	0	GLY	-	expression tag	UNP A0A059PIQ0
A	15	QCA	THR	chromophore	UNP A0A059PIQ0
A	?	-	TYR	chromophore	UNP A0A059PIQ0
A	?	-	GLY	chromophore	UNP A0A059PIQ0
A	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
A	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
A	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
A	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
A	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
A	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
A	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
A	155	THR	SER	engineered mutation	UNP A0A059PIQ0
A	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
A	188	GLY	-	linker	UNP A0A059PIQ0
A	189	GLY	-	linker	UNP A0A059PIQ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	THR	-	linker	UNP A0A059PIQ0
A	191	GLY	-	linker	UNP A0A059PIQ0
A	192	GLY	-	linker	UNP A0A059PIQ0
A	193	SER	-	linker	UNP A0A059PIQ0
A	194	ALA	-	linker	UNP A0A059PIQ0
A	195	SER	-	linker	UNP A0A059PIQ0
A	196	GLN	-	linker	UNP A0A059PIQ0
A	223	ARG	SER	engineered mutation	UNP P42212
A	232	ILE	TYR	engineered mutation	UNP P42212
A	241	SER	CYS	engineered mutation	UNP P42212
B	-10	MET	-	initiating methionine	UNP A0A059PIQ0
B	-9	GLY	-	expression tag	UNP A0A059PIQ0
B	-8	HIS	-	expression tag	UNP A0A059PIQ0
B	-7	HIS	-	expression tag	UNP A0A059PIQ0
B	-6	HIS	-	expression tag	UNP A0A059PIQ0
B	-5	HIS	-	expression tag	UNP A0A059PIQ0
B	-4	HIS	-	expression tag	UNP A0A059PIQ0
B	-3	HIS	-	expression tag	UNP A0A059PIQ0
B	-2	SER	-	expression tag	UNP A0A059PIQ0
B	-1	SER	-	expression tag	UNP A0A059PIQ0
B	0	GLY	-	expression tag	UNP A0A059PIQ0
B	15	QCA	THR	chromophore	UNP A0A059PIQ0
B	?	-	TYR	chromophore	UNP A0A059PIQ0
B	?	-	GLY	chromophore	UNP A0A059PIQ0
B	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
B	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
B	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
B	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
B	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
B	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
B	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
B	155	THR	SER	engineered mutation	UNP A0A059PIQ0
B	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
B	188	GLY	-	linker	UNP A0A059PIQ0
B	189	GLY	-	linker	UNP A0A059PIQ0
B	190	THR	-	linker	UNP A0A059PIQ0
B	191	GLY	-	linker	UNP A0A059PIQ0
B	192	GLY	-	linker	UNP A0A059PIQ0
B	193	SER	-	linker	UNP A0A059PIQ0
B	194	ALA	-	linker	UNP A0A059PIQ0
B	195	SER	-	linker	UNP A0A059PIQ0
B	196	GLN	-	linker	UNP A0A059PIQ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	223	ARG	SER	engineered mutation	UNP P42212
B	232	ILE	TYR	engineered mutation	UNP P42212
B	241	SER	CYS	engineered mutation	UNP P42212


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	139	Total 139	O 139	0	0
2	B	124	Total 124	O 124	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.95Å 67.72Å 58.36Å 90.00° 102.28° 90.00°	Depositor
Resolution (Å)	38.53 – 1.50 38.53 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.0 (38.53-1.50) 92.3 (38.53-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.89 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.13RC2_2986	Depositor
R, R_{free}	0.159 , 0.181 0.162 , 0.182	Depositor DCC
R_{free} test set	2862 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4023	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	2/1948 (0.1%)	0.51	0/2636
1	B	0.27	0/1928	0.50	0/2607
All	All	0.33	2/3876 (0.1%)	0.50	0/5243

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	GLU	C-O	-5.80	1.16	1.23
1	A	40	GLU	C-O	-5.56	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1881	0	1851	20	0
1	B	1879	0	1844	15	0
2	A	139	0	0	4	0
2	B	124	0	0	4	0
All	All	4023	0	3695	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:238:LYS:HE2	1:B:240[B]:ILE:HD11	1.54	0.89
1:A:156[B]:VAL:HG23	1:B:156:VAL:HG13	1.66	0.78
1:B:51[B]:LYS:HG3	1:B:128:LEU:HB2	1.74	0.68
1:A:161:PRO:HG3	1:B:232:ILE:HG21	1.80	0.63
1:A:195:SER:OG	1:A:198:GLU:HG3	2.01	0.60
1:B:192:GLY:N	2:B:404:HOH:O	2.33	0.60
1:A:156[A]:VAL:HG13	1:B:156:VAL:HG13	1.83	0.60
1:B:29:LYS:HE3	2:B:465:HOH:O	2.02	0.58
1:A:28:MET:HE3	1:A:179:ILE:HD12	1.87	0.55
1:A:112:LYS:HE2	1:A:132:TYR:OH	2.07	0.54
1:A:156[B]:VAL:HG11	1:B:173:PHE:HD2	1.73	0.53
1:A:55:LYS:HE3	1:A:79:ASP:OD2	2.09	0.53
1:B:231:THR:HG23	2:B:401:HOH:O	2.11	0.50
1:A:107[B]:GLN:HG3	1:A:108:LYS:N	2.27	0.49
1:A:236:THR:HG23	2:A:426:HOH:O	2.12	0.48
1:A:70[B]:VAL:HG23	2:A:465:HOH:O	2.13	0.47
1:A:156[B]:VAL:HG11	1:B:173:PHE:CD2	2.48	0.47
1:A:156[A]:VAL:HG12	1:B:154:GLN:HB3	1.98	0.46
1:B:196:GLN:HG2	2:B:432:HOH:O	2.16	0.45
1:A:51:LYS:HD3	1:A:51:LYS:HA	1.72	0.45
1:B:27:HIS:HB2	1:B:182:GLY:HA2	2.00	0.44
1:A:154:GLN:HB3	1:B:156:VAL:HG12	2.01	0.42
1:A:76:LYS:HA	1:A:76:LYS:HD3	1.92	0.42
1:A:23:ARG:HD2	1:A:231:THR:O	2.20	0.42
1:A:198:GLU:CG	2:A:414:HOH:O	2.67	0.42
1:A:223:ARG:HH12	1:A:225[A]:GLU:CD	2.28	0.41
1:B:196:GLN:H	1:B:196:GLN:CD	2.29	0.41
1:A:198:GLU:HG2	2:A:414:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/252 (94%)	235 (99%)	2 (1%)	0	100	100
1	B	236/252 (94%)	232 (98%)	3 (1%)	1 (0%)	30	12
All	All	473/504 (94%)	467 (99%)	5 (1%)	1 (0%)	43	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	181	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/215 (100%)	214 (100%)	0	100	100
1	B	210/215 (98%)	210 (100%)	0	100	100
All	All	424/430 (99%)	424 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	96	ASN
1	B	114	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	QCA	A	15	1	25,26,27	2.28	5 (20%)	31,37,39	2.17	11 (35%)
1	QCA	B	15	1	25,26,27	2.50	5 (20%)	31,37,39	2.20	13 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	QCA	A	15	1	-	0/14/35/36	0/2/2/2
1	QCA	B	15	1	-	0/14/35/36	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	15	QCA	CB2-CA2	9.98	1.44	1.35
1	A	15	QCA	CB2-CA2	8.59	1.43	1.35
1	B	15	QCA	CB1-CA1	-3.46	1.40	1.53
1	A	15	QCA	OG1-CB1	-3.40	1.34	1.43
1	A	15	QCA	CB1-CA1	-3.37	1.40	1.53
1	B	15	QCA	CA1-N1	-3.18	1.38	1.47
1	B	15	QCA	OG1-CB1	-3.04	1.35	1.43
1	A	15	QCA	CA1-N1	-2.55	1.40	1.47
1	A	15	QCA	C1-N3	2.24	1.41	1.37
1	B	15	QCA	C1-N2	2.09	1.35	1.32

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	QCA	O2-C2-CA2	-6.11	127.12	131.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	15	QCA	O2-C2-CA2	-6.05	127.16	131.02
1	B	15	QCA	C2-N3-C1	4.26	110.04	108.07
1	A	15	QCA	CG2-CD1-CE1	-4.24	116.60	120.22
1	A	15	QCA	CA2-N2-C1	4.02	108.94	105.80
1	B	15	QCA	CG2-CD1-CE1	-3.58	117.17	120.22
1	B	15	QCA	CA2-N2-C1	3.44	108.49	105.80
1	B	15	QCA	OG1-CB1-CA1	3.36	116.15	109.00
1	A	15	QCA	OG1-CB1-CA1	3.08	115.55	109.00
1	A	15	QCA	C2-N3-C1	3.04	109.47	108.07
1	A	15	QCA	CD2-CG2-CD1	2.86	122.32	118.72
1	A	15	QCA	CG2-CB2-CA2	-2.81	126.52	129.87
1	B	15	QCA	CD2-CG2-CD1	2.71	122.13	118.72
1	B	15	QCA	CG2-CB2-CA2	-2.49	126.90	129.87
1	A	15	QCA	C2-CA2-N2	-2.45	107.20	108.95
1	A	15	QCA	CD2-CE2-CZ	-2.40	118.10	120.50
1	B	15	QCA	C2-CA2-N2	-2.39	107.24	108.95
1	B	15	QCA	CE2-CZ-CE1	2.37	121.91	118.94
1	A	15	QCA	CE2-CZ-CE1	2.36	121.88	118.94
1	B	15	QCA	CD2-CG2-CB2	-2.20	113.64	121.22
1	B	15	QCA	O2-C2-N3	2.11	128.71	124.31
1	B	15	QCA	C3-CA3-N3	2.07	117.14	112.43
1	B	15	QCA	CD2-CE2-CZ	-2.07	118.43	120.50
1	A	15	QCA	O2-C2-N3	2.04	128.57	124.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	224/252 (88%)	0.05	7 (3%)	51	55	11, 25, 53, 86	18 (8%)
1	B	230/252 (91%)	0.16	8 (3%)	47	51	13, 27, 60, 100	11 (4%)
All	All	454/504 (90%)	0.11	15 (3%)	49	53	11, 26, 59, 100	29 (6%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	182	GLY	4.8
1	A	179	ILE	4.4
1	A	195	SER	3.3
1	B	180	THR	3.1
1	A	242	THR	3.1
1	A	243	THR	3.1
1	B	181	HIS	2.8
1	B	243	THR	2.7
1	B	140	ASP	2.5
1	B	196	GLN	2.4
1	A	161	PRO	2.3
1	B	194	ALA	2.2
1	A	162	ASN	2.2
1	B	107[A]	GLN	2.1
1	A	203	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	QCA	A	15	25/26	0.95	0.08	14,19,26,48	0
1	QCA	B	15	25/26	0.96	0.07	16,19,32,46	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.