



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:31 AM UTC

PDB ID : 3UOX / pdb_00003uox
Title : Crystal Structure of OTEMO (FAD bound form 2)
Authors : Shi, R.; Matte, A.; Cygler, M.; Lau, P.
Deposited on : 2011-11-17
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	545	
1	B	545	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OTEMO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	537	Total	C	N	O	S	0	1	0
			4272	2715	736	807	14			
1	B	533	Total	C	N	O	S	0	1	0
			4245	2699	732	800	14			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	206	Total 206	O 206	0	0
3	B	199	Total 199	O 199	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: OTEMO



• Molecule 1: OTEMO



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.32Å 92.54Å 93.35Å 90.00° 103.27° 90.00°	Depositor
Resolution (Å)	48.16 – 1.96 48.16 – 1.96	Depositor EDS
% Data completeness (in resolution range)	98.2 (48.16-1.96) 98.1 (48.16-1.96)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.190 , 0.230 0.192 , 0.234	Depositor DCC
R_{free} test set	3973 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9028	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/4385 (0.0%)	0.96	5/5957 (0.1%)
1	B	0.85	0/4357	0.96	9/5916 (0.2%)
All	All	0.87	1/8742 (0.0%)	0.96	14/11873 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	ILE	CA-CB	5.10	1.57	1.54

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241	TYR	CA-C-N	-6.35	112.30	119.28
1	B	241	TYR	C-N-CA	-6.35	112.30	119.28
1	B	513	VAL	N-CA-C	6.17	117.08	107.28
1	A	513	VAL	N-CA-C	5.73	116.11	107.75
1	B	331	HIS	N-CA-C	5.62	117.34	108.23
1	B	142	ALA	N-CA-C	-5.52	102.16	110.06
1	B	86	GLN	CA-C-N	-5.29	113.58	119.19
1	B	86	GLN	C-N-CA	-5.29	113.58	119.19
1	A	222	THR	CA-C-N	-5.16	114.60	119.76
1	A	222	THR	C-N-CA	-5.16	114.60	119.76
1	B	508	LYS	CA-C-N	5.05	126.16	119.84
1	B	508	LYS	C-N-CA	5.05	126.16	119.84
1	A	241	TYR	CA-C-N	-5.04	113.73	119.28
1	A	241	TYR	C-N-CA	-5.04	113.73	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4272	0	4126	25	0
1	B	4245	0	4100	33	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	206	0	0	1	0
3	B	199	0	0	6	0
All	All	9028	0	8288	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:GLU:HG2	1:A:109:ASN:HD21	1.39	0.85
1:A:94:ASN:HD22	1:A:103:ARG:HH21	1.25	0.82
1:B:94:ASN:HD22	1:B:103:ARG:HH21	1.34	0.74
1:A:220:TRP:O	1:A:339:PRO:HD2	1.95	0.66
1:A:481:ARG:HG3	3:A:727:HOH:O	1.94	0.66
1:B:131:GLU:HG2	1:B:132:GLU:N	2.14	0.63
1:B:131:GLU:HG2	1:B:132:GLU:H	1.64	0.63
1:B:51:ASN:O	1:B:86:GLN:HG3	1.99	0.62
1:B:487:TYR:OH	1:B:515:ARG:HD3	1.99	0.61
1:A:94:ASN:HD22	1:A:103:ARG:NH2	1.99	0.60
1:B:501:TRP:CD1	1:B:501:TRP:H	2.19	0.58
1:A:24:GLN:HE21	1:A:434:LEU:HD11	1.69	0.58
1:A:51:ASN:O	1:A:86:GLN:HG3	2.03	0.58
1:B:24:GLN:HE21	1:B:434:LEU:HD11	1.70	0.57
1:B:485:GLU:OE2	1:B:530:ARG:NH1	2.37	0.56
1:B:487:TYR:CZ	1:B:515:ARG:HD3	2.40	0.56
1:B:54:PRO:HG3	1:B:181:VAL:HG12	1.88	0.55
1:B:30:GLN:NE2	3:B:651:HOH:O	2.38	0.55
1:A:424:GLN:HE21	1:A:474:PRO:HD3	1.71	0.55
1:A:495:LEU:HD12	1:A:521:VAL:HG11	1.88	0.55
1:A:202:PRO:HG3	1:A:344:TYR:HA	1.88	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:507:THR:HA	1:B:512:SER:O	2.08	0.54
1:A:94:ASN:ND2	1:A:103:ARG:HH21	2.01	0.52
1:B:417:PRO:HD3	1:B:518:LEU:HD23	1.92	0.52
1:B:82:ASN:HD21	1:B:233:LYS:NZ	2.08	0.51
1:B:42:GLU:HG2	1:B:109:ASN:HD21	1.76	0.51
1:A:424:GLN:HG2	1:A:431:PHE:O	2.13	0.48
1:B:220:TRP:O	1:B:339:PRO:HD2	2.13	0.47
1:B:485:GLU:CD	1:B:530:ARG:NH1	2.72	0.47
1:A:532:GLU:OE2	1:A:532:GLU:HA	2.14	0.47
1:B:196:THR:HG23	3:B:685:HOH:O	2.15	0.47
1:A:42:GLU:HG2	1:A:109:ASN:ND2	2.19	0.47
1:B:282:GLN:HG2	3:B:555:HOH:O	2.14	0.47
1:A:318:LYS:HD3	1:A:318:LYS:HA	1.61	0.46
1:B:370:GLY:HA3	1:B:378:TYR:O	2.15	0.46
1:A:127:THR:CG2	1:A:130:ASN:HA	2.45	0.46
1:B:501:TRP:H	1:B:501:TRP:HD1	1.63	0.45
1:B:367:THR:HB	1:B:368:PRO:HD2	1.98	0.45
1:B:424:GLN:HE21	1:B:474:PRO:HD3	1.81	0.45
1:B:485:GLU:OE1	1:B:530:ARG:NH1	2.42	0.45
1:B:508:LYS:HB2	1:B:508:LYS:NZ	2.31	0.45
1:B:119:GLU:HG3	3:B:661:HOH:O	2.17	0.44
1:A:54:PRO:HG3	1:A:181:VAL:HG12	1.99	0.44
1:B:490:PHE:CE2	1:B:496:ALA:HA	2.52	0.44
1:A:131:GLU:HG3	1:A:132:GLU:H	1.83	0.43
1:A:372:LYS:HD3	1:A:373:THR:O	2.19	0.42
1:A:483:THR:CG2	1:A:516:ARG:HG2	2.50	0.42
1:B:181:VAL:HG22	3:B:650:HOH:O	2.19	0.41
1:B:495:LEU:HD12	1:B:521:VAL:HG11	2.02	0.41
1:B:255:PHE:CE1	1:B:285:TYR:HD2	2.39	0.41
1:B:487:TYR:CZ	1:B:515:ARG:CD	3.04	0.41
1:B:107:ARG:NH2	3:B:670:HOH:O	2.48	0.41
1:A:123:LEU:C	1:A:123:LEU:HD12	2.46	0.41
1:A:62:SER:HB3	1:A:79:TRP:CD2	2.56	0.41
1:A:194:GLY:O	1:A:198:VAL:HG13	2.21	0.41
1:A:82:ASN:HD21	1:A:233:LYS:NZ	2.19	0.40
1:A:331:HIS:HA	1:A:332:PRO:HD3	1.95	0.40
1:B:10:LEU:HD23	1:B:34:LYS:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/545 (98%)	517 (97%)	13 (2%)	4 (1%)	18	10
1	B	528/545 (97%)	515 (98%)	11 (2%)	2 (0%)	30	21
All	All	1062/1090 (97%)	1032 (97%)	24 (2%)	6 (1%)	21	12

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	509	PRO
1	A	130	ASN
1	A	509	PRO
1	A	435	VAL
1	B	435	VAL
1	A	511	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/454 (98%)	430 (96%)	17 (4%)	29	19
1	B	444/454 (98%)	435 (98%)	9 (2%)	48	43
All	All	891/908 (98%)	865 (97%)	26 (3%)	37	29

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	181	VAL
1	A	198	VAL
1	A	240	ARG
1	A	285	TYR
1	A	318	LYS
1	A	359	ARG
1	A	360	GLU
1	A	369	GLU
1	A	372	LYS
1	A	398	ARG
1	A	410	ILE
1	A	444	CYS
1	A	504	LYS
1	A	507	THR
1	A	508	LYS
1	A	514	VAL
1	B	131	GLU
1	B	153	ASP
1	B	196	THR
1	B	270	SER
1	B	329	LYS
1	B	364	GLN
1	B	444	CYS
1	B	508	LYS
1	B	530	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	29	ASN
1	A	51	ASN
1	A	82	ASN
1	A	86	GLN
1	A	94	ASN
1	A	109	ASN
1	A	130	ASN
1	A	315	GLN
1	A	424	GLN
1	A	467	ASN
1	A	499	ASN
1	B	24	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	51	ASN
1	B	82	ASN
1	B	94	ASN
1	B	109	ASN
1	B	315	GLN
1	B	406	ASN
1	B	424	GLN
1	B	467	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	551	-	58,58,58	1.12	5 (8%)	85,89,89	1.56	14 (16%)
2	FAD	A	551	-	58,58,58	1.25	6 (10%)	85,89,89	1.73	19 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	551	-	-	3/34/50/50	0/6/6/6
2	FAD	A	551	-	-	3/34/50/50	0/6/6/6

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	551	FAD	C4X-N5	4.59	1.40	1.30
2	B	551	FAD	C4X-N5	3.51	1.38	1.30
2	B	551	FAD	C2A-N1A	3.13	1.39	1.33
2	B	551	FAD	C2A-N3A	2.97	1.39	1.33
2	A	551	FAD	C2A-N3A	2.80	1.38	1.33
2	A	551	FAD	C2A-N1A	2.70	1.38	1.33
2	A	551	FAD	P-O3P	2.47	1.62	1.59
2	B	551	FAD	C10-N1	2.42	1.38	1.33
2	A	551	FAD	C10-N1	2.37	1.38	1.33
2	A	551	FAD	C8A-N7A	2.33	1.36	1.31
2	B	551	FAD	C5A-N7A	-2.18	1.35	1.39

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	551	FAD	N3A-C2A-N1A	-7.14	117.77	128.58
2	A	551	FAD	N3A-C2A-N1A	-6.08	119.38	128.58
2	A	551	FAD	N9A-C8A-N7A	-4.63	107.36	113.94
2	A	551	FAD	C5A-C4A-N3A	-4.24	120.87	126.72
2	B	551	FAD	N9A-C8A-N7A	-3.95	108.33	113.94
2	A	551	FAD	C5A-N7A-C8A	3.79	109.41	103.45
2	A	551	FAD	C2A-N3A-C4A	3.67	120.80	111.83
2	A	551	FAD	C4-N3-C2	-3.55	119.34	125.64
2	A	551	FAD	C5X-C9A-N10	3.34	120.99	117.97
2	B	551	FAD	C2A-N3A-C4A	3.30	119.88	111.83
2	B	551	FAD	C5A-C4A-N3A	-3.16	122.36	126.72
2	A	551	FAD	C9A-C5X-N5	-3.14	119.12	122.45
2	A	551	FAD	C4X-C4-N3	3.14	121.23	113.25
2	A	551	FAD	N3A-C4A-N9A	3.07	132.38	127.17
2	B	551	FAD	C4X-C10-N10	2.85	120.56	116.48
2	B	551	FAD	C5A-N7A-C8A	2.82	107.88	103.45
2	A	551	FAD	C4A-N9A-C8A	2.75	108.62	105.74
2	A	551	FAD	O4-C4-C4X	-2.74	119.30	126.53
2	B	551	FAD	C4X-C4-N3	2.68	120.06	113.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	551	FAD	C4-N3-C2	-2.64	120.96	125.64
2	B	551	FAD	O4-C4-C4X	-2.60	119.66	126.53
2	B	551	FAD	C4A-N9A-C8A	2.45	108.31	105.74
2	B	551	FAD	C2A-N1A-C6A	2.44	122.75	118.73
2	A	551	FAD	C4A-C5A-N7A	-2.38	107.86	110.58
2	A	551	FAD	O3P-PA-O1A	-2.34	103.66	110.70
2	B	551	FAD	N3A-C4A-N9A	2.33	131.14	127.17
2	B	551	FAD	C1'-C2'-C3'	2.30	115.91	109.66
2	A	551	FAD	C4-C4X-N5	2.26	121.33	118.21
2	A	551	FAD	C10-C4X-N5	-2.19	120.34	124.81
2	A	551	FAD	C2B-C3B-C4B	2.14	106.75	102.61
2	A	551	FAD	C4X-C10-N10	2.13	119.53	116.48
2	B	551	FAD	C10-C4X-N5	-2.11	120.49	124.81
2	A	551	FAD	C1'-C2'-C3'	2.06	115.25	109.66

There are no chirality outliers.


All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	551	FAD	N10-C1'-C2'-O2'
2	A	551	FAD	N10-C1'-C2'-C3'
2	A	551	FAD	PA-O3P-P-O5'
2	B	551	FAD	N10-C1'-C2'-O2'
2	B	551	FAD	N10-C1'-C2'-C3'
2	B	551	FAD	PA-O3P-P-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	537/545 (98%)	0.08	24 (4%)	38 44	11, 21, 40, 64	1 (0%)
1	B	533/545 (97%)	0.10	21 (3%)	43 50	11, 22, 39, 59	1 (0%)
All	All	1070/1090 (98%)	0.09	45 (4%)	40 47	11, 22, 39, 64	2 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	501	TRP	5.8
1	A	509	PRO	4.9
1	B	152	PRO	4.9
1	A	148	ALA	4.7
1	B	512	SER	4.6
1	A	176	GLY	4.2
1	A	508	LYS	4.0
1	A	511	GLY	4.0
1	A	389	PHE	4.0
1	B	148	ALA	4.0
1	A	514	VAL	3.8
1	A	152	PRO	3.8
1	B	394	GLY	3.8
1	B	389	PHE	3.7
1	B	509	PRO	3.6
1	A	501	TRP	3.5
1	B	388	GLY	3.5
1	B	507	THR	3.4
1	A	177	ALA	3.3
1	B	513	VAL	3.3
1	A	512	SER	3.0
1	B	398	ARG	2.9
1	B	506	THR	2.9
1	B	511	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	398	ARG	2.7
1	A	392	VAL	2.7
1	A	391	ALA	2.7
1	A	500	ALA	2.6
1	B	508	LYS	2.6
1	A	174	ALA	2.6
1	A	175	GLU	2.6
1	A	130	ASN	2.6
1	B	154	ILE	2.6
1	A	330	ASP	2.5
1	B	514	VAL	2.5
1	A	449	CYS	2.5
1	A	513	VAL	2.4
1	A	390	ASP	2.4
1	B	174	ALA	2.3
1	A	507	THR	2.2
1	B	387	THR	2.2
1	B	147	SER	2.1
1	B	265	THR	2.1
1	B	155	LYS	2.1
1	A	510	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FAD	A	551	53/53	0.98	0.05	12,16,22,24	0
2	FAD	B	551	53/53	0.98	0.04	13,16,24,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.