



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:02 AM UTC

PDB ID : 4UO8 / pdb_00004uo8
Title : Structure of the A_Canine_Colorado_17864_06 H3 haemagglutinin in complex with 6SO4-3SLN
Authors : Vachieri, S.G.; Collins, P.J.; Haire, L.F.; Ogradowicz, R.W.; Martin, S.R.; Walker, P.A.; Xiong, X.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2014-05-31
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	84% 11% ..
2	B	175	87% 13%
3	C	2	100%
4	D	5	20% 80%
4	E	5	60% 40%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	5	 20% 80%
6	G	3	 33% 33% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	D	1	X	-	-	-
4	MAN	D	5	X	-	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 4190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2468	1541	434	477	16			


- Molecule 2 is a protein called HA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	175	Total	C	N	O	S	0	0	0
			1425	886	250	283	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	173	SER	-	expression tag	UNP E0UVR5
B	174	GLY	-	expression tag	UNP E0UVR5
B	175	ARG	-	expression tag	UNP E0UVR5
B	131	GLU	ASP	conflict	UNP E0UVR5

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	5	Total	C	N	O	0	0	0
			61	34	2	25			
4	E	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-2)-alpha-D-mannopyranos e-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-ace tamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galacto pyranose-(1-4)-2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	3	Total	C	N	O	S	0	0	0
			50	25	2	22	1			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		
8	A	1	Total	O	S	0	0
			5	4	1		
8	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	4	Total 4	O 4	0	0
9	B	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HA1



- Molecule 2: HA2




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  60% 40%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 5: beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  20% 80%

MAG1
MAG2
BMA3
MAN4
BMA5

- Molecule 6: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyranose

Chain G:  33% 33% 33%

NCS1
GAL2
SIA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	97.29Å 97.29Å 348.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.69 – 3.00 48.69 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.69-3.00) 99.3 (48.69-3.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.207 , 0.251 0.216 , 0.207	Depositor DCC
R_{free} test set	1050 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	64.2	Xtriage
Anisotropy	0.868	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4190	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SIA, GAL, NGS, MAN, SO4, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2519	0.75	0/3418
2	B	0.51	0/1450	0.83	1/1949 (0.1%)
All	All	0.51	0/3969	0.78	1/5367 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	12	ASN	N-CA-C	5.04	115.96	108.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2468	0	2412	16	0
2	B	1425	0	1344	7	0
3	C	28	0	25	0	0
4	D	61	0	52	0	0
4	E	61	0	52	2	0
5	F	61	0	52	0	0
6	G	50	0	35	1	0
7	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	10	0	0	0	0
8	B	5	0	0	0	0
9	A	4	0	0	0	0
9	B	3	0	0	0	0
All	All	4190	0	3985	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:ASN:OD1	1:A:304:LYS:O	2.16	0.64
1:A:15:LEU:HD22	2:B:119:PHE:HA	1.83	0.60
6:G:3:SIA:H113	6:G:3:SIA:O7	2.06	0.55
1:A:27:LYS:NZ	2:B:97:GLU:OE2	2.40	0.54
1:A:34:ILE:HD11	1:A:321:ARG:HD2	1.92	0.51
1:A:180:TRP:CE2	1:A:204:VAL:HG21	2.46	0.50
4:E:3:BMA:H61	4:E:5:MAN:H5	1.94	0.49
2:B:19:ASP:HB2	2:B:36:ALA:HB2	1.94	0.48
1:A:74:PRO:HG2	1:A:139:CYS:HB2	1.97	0.47
1:A:295:GLN:HB3	1:A:306:PRO:HB2	1.97	0.47
2:B:158:ASP:HB3	2:B:161:ILE:HD12	1.97	0.46
1:A:272:VAL:HG13	1:A:273:PRO:HD2	1.98	0.45
4:E:3:BMA:C6	4:E:5:MAN:H5	2.47	0.45
2:B:6:ILE:O	2:B:7:ALA:HB3	2.16	0.44
1:A:197:GLN:NE2	1:A:248:ASN:O	2.49	0.44
1:A:97:CYS:HA	1:A:139:CYS:HB3	2.00	0.44
1:A:67:ILE:O	1:A:70:MET:HG3	2.17	0.43
1:A:170:ASN:OD1	1:A:170:ASN:C	2.61	0.43
2:B:78:GLN:HE21	2:B:82:LYS:NZ	2.16	0.42
1:A:54:LYS:O	1:A:55:SER:C	2.62	0.42
1:A:288:ILE:HD12	1:A:295:GLN:HG3	2.03	0.41
2:B:5:ALA:HB3	2:B:112:ASP:OD1	2.20	0.41
1:A:248:ASN:HD22	1:A:248:ASN:H	1.68	0.40
1:A:296:ASN:CG	1:A:296:ASN:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/328 (97%)	296 (93%)	20 (6%)	1 (0%)	36	70
2	B	173/175 (99%)	162 (94%)	10 (6%)	1 (1%)	21	56
All	All	490/503 (97%)	458 (94%)	30 (6%)	2 (0%)	30	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	172	GLN
1	A	163	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/289 (96%)	256 (92%)	23 (8%)	10	37
2	B	148/148 (100%)	139 (94%)	9 (6%)	17	49
All	All	427/437 (98%)	395 (92%)	32 (8%)	12	41

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	40	THR
1	A	45	SER
1	A	54	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	SER
1	A	59	LEU
1	A	70	MET
1	A	126	THR
1	A	131	THR
1	A	139	CYS
1	A	155	THR
1	A	156	LYS
1	A	173	ASN
1	A	210	GLN
1	A	223	VAL
1	A	230	ILE
1	A	242	ILE
1	A	248	ASN
1	A	260	LEU
1	A	264	LYS
1	A	295	GLN
1	A	311	GLN
1	A	326	LYS
2	B	17	MET
2	B	38	LEU
2	B	48	ILE
2	B	59	THR
2	B	60	ASN
2	B	102	LEU
2	B	110	LEU
2	B	126	LEU
2	B	131	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	173	ASN
1	A	183	HIS
1	A	211	GLN
1	A	226	GLN
1	A	248	ASN
1	A	261	ASN
1	A	311	GLN
2	B	34	GLN
2	B	53	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	60	ASN
2	B	78	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.83	0	17,19,21	1.48	2 (11%)
3	NAG	C	2	3	14,14,15	0.44	0	17,19,21	1.65	1 (5%)
4	NAG	D	1	1,4	14,14,15	0.52	0	17,19,21	0.89	0
4	NAG	D	2	4	14,14,15	0.56	0	17,19,21	1.07	1 (5%)
4	BMA	D	3	4	11,11,12	0.44	0	15,15,17	1.03	1 (6%)
4	MAN	D	4	4	11,11,12	0.54	0	15,15,17	1.05	1 (6%)
4	MAN	D	5	4	11,11,12	0.61	0	15,15,17	1.30	3 (20%)
4	NAG	E	1	1,4	14,14,15	0.51	0	17,19,21	1.87	3 (17%)
4	NAG	E	2	4	14,14,15	0.56	0	17,19,21	0.86	1 (5%)
4	BMA	E	3	4	11,11,12	0.52	0	15,15,17	1.21	1 (6%)
4	MAN	E	4	4	11,11,12	0.87	0	15,15,17	1.86	3 (20%)
4	MAN	E	5	4	11,11,12	0.64	0	15,15,17	1.37	2 (13%)
5	NAG	F	1	1,5	14,14,15	0.73	0	17,19,21	0.97	1 (5%)
5	NAG	F	2	5	14,14,15	0.54	0	17,19,21	0.69	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BMA	F	3	5	11,11,12	0.42	0	15,15,17	1.03	1 (6%)
5	MAN	F	4	5	11,11,12	0.87	1 (9%)	15,15,17	1.77	4 (26%)
5	BMA	F	5	5	11,11,12	0.59	0	15,15,17	1.76	2 (13%)
6	NGS	G	1	6	19,19,19	0.74	1 (5%)	27,28,28	1.49	3 (11%)
6	GAL	G	2	6	11,11,12	0.53	0	15,15,17	0.77	0
6	SIA	G	3	6	20,20,21	0.54	0	21,28,31	1.64	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	NAG	D	1	1,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	2/2/19/22	0/1/1/1
4	MAN	D	4	4	-	0/2/19/22	1/1/1/1
4	MAN	D	5	4	1/1/4/5	2/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1
4	MAN	E	4	4	-	2/2/19/22	0/1/1/1
4	MAN	E	5	4	-	0/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	0/2/19/22	0/1/1/1
5	BMA	F	5	5	-	2/2/19/22	0/1/1/1
6	NGS	G	1	6	-	8/10/30/30	0/1/1/1
6	GAL	G	2	6	-	0/2/19/22	0/1/1/1
6	SIA	G	3	6	-	2/18/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	1	NGS	O8-S	2.25	1.64	1.50
5	F	4	MAN	C2-C3	2.08	1.55	1.52

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C1-O5-C5	6.14	120.41	112.19
4	E	1	NAG	C1-O5-C5	5.36	119.37	112.19
4	E	4	MAN	C1-C2-C3	5.30	117.36	109.64
5	F	5	BMA	C1-O5-C5	4.90	118.75	112.19
3	C	1	NAG	C4-C3-C2	4.28	117.30	111.02
5	F	4	MAN	C3-C4-C5	4.13	117.72	110.23
5	F	5	BMA	C1-C2-C3	3.73	115.07	109.64
6	G	1	NGS	C8-C7-N2	3.51	121.94	116.12
4	D	4	MAN	C1-O5-C5	3.46	116.82	112.19
4	E	3	BMA	C1-C2-C3	3.39	114.57	109.64
4	E	4	MAN	C2-C3-C4	3.36	116.76	110.86
6	G	3	SIA	O1B-C1-C2	3.34	121.40	112.71
6	G	3	SIA	C11-C10-N5	3.22	121.46	116.12
4	D	5	MAN	O5-C1-C2	3.19	118.41	110.79
5	F	4	MAN	C2-C3-C4	3.15	116.40	110.86
4	E	5	MAN	C3-C4-C5	3.08	115.82	110.23
6	G	3	SIA	O6-C2-C3	-3.05	106.45	110.56
4	D	2	NAG	C4-C3-C2	2.91	115.29	111.02
4	E	1	NAG	C8-C7-N2	2.78	120.72	116.12
5	F	3	BMA	C1-C2-C3	2.59	113.41	109.64
3	C	1	NAG	O5-C5-C4	-2.59	104.54	110.83
6	G	1	NGS	O6-C6-C5	2.50	112.02	107.57
4	D	5	MAN	C1-C2-C3	2.50	113.28	109.64
6	G	1	NGS	C1-C2-N2	-2.45	107.88	110.73
4	E	1	NAG	O5-C1-C2	-2.43	107.53	111.29
5	F	4	MAN	C1-C2-C3	2.35	113.06	109.64
4	D	3	BMA	C3-C4-C5	2.34	114.48	110.23
4	E	2	NAG	C4-C3-C2	-2.34	107.59	111.02
6	G	3	SIA	C5-N5-C10	2.31	128.51	123.11
4	E	5	MAN	C1-O5-C5	2.20	115.14	112.19
4	E	4	MAN	C3-C4-C5	2.20	114.22	110.23
4	D	5	MAN	C1-O5-C5	2.16	115.08	112.19
6	G	3	SIA	O1A-C1-C2	-2.11	118.29	122.85
5	F	4	MAN	O2-C2-C3	2.11	114.51	110.15
5	F	1	NAG	C4-C3-C2	2.07	114.05	111.02

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	1	NAG	C1
4	D	5	MAN	C1

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	1	NGS	C4-C5-C6-O6
6	G	1	NGS	O5-C5-C6-O6
4	E	1	NAG	O5-C5-C6-O6
5	F	5	BMA	O5-C5-C6-O6
4	D	5	MAN	O5-C5-C6-O6
4	E	4	MAN	C4-C5-C6-O6
4	E	4	MAN	O5-C5-C6-O6
4	E	3	BMA	O5-C5-C6-O6
4	E	3	BMA	C4-C5-C6-O6
5	F	5	BMA	C4-C5-C6-O6
4	D	3	BMA	C4-C5-C6-O6
4	E	1	NAG	C8-C7-N2-C2
4	E	1	NAG	O7-C7-N2-C2
6	G	1	NGS	O7-C7-N2-C2
6	G	1	NGS	C8-C7-N2-C2
6	G	3	SIA	C11-C10-N5-C5
6	G	3	SIA	O10-C10-N5-C5
4	D	3	BMA	O5-C5-C6-O6
4	E	1	NAG	C4-C5-C6-O6
4	D	5	MAN	C4-C5-C6-O6
6	G	1	NGS	C6-O6-S-O7A
6	G	1	NGS	C6-O6-S-O9
6	G	1	NGS	C1-C2-N2-C7
6	G	1	NGS	C3-C2-N2-C7


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	4	MAN	C1-C2-C3-C4-C5-O5


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	5	MAN	2	0
4	E	3	BMA	2	0
6	G	3	SIA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




Oligosaccharide Chain E



Oligosaccharide Chain F





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	SO4	A	1328	-	4,4,4	0.48	0	6,6,6	0.07	0
8	SO4	A	1327	-	4,4,4	0.44	0	6,6,6	0.09	0
7	NAG	A	601	1	14,14,15	0.53	0	17,19,21	1.43	3 (17%)
8	SO4	B	1176	-	4,4,4	0.43	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	601	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	601	NAG	C2-N2-C7	2.93	126.83	122.90
7	A	601	NAG	C8-C7-N2	2.91	120.94	116.12
7	A	601	NAG	O5-C5-C6	2.18	111.91	107.66

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	601	NAG	C8-C7-N2-C2
7	A	601	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/328 (97%)	-0.11	3 (0%) 81 61	57, 80, 119, 144	0
2	B	175/175 (100%)	-0.29	2 (1%) 78 57	56, 75, 96, 129	0
All	All	494/503 (98%)	-0.17	5 (1%) 79 59	56, 79, 115, 144	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	THR	3.3
2	B	173	SER	2.6
1	A	217	ILE	2.1
2	B	1	GLY	2.1
1	A	9	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	D	3	11/12	0.36	0.12	141,152,158,158	0
4	MAN	E	5	11/12	0.43	0.16	153,164,167,167	0
4	NAG	D	2	14/15	0.48	0.17	127,137,145,153	0
5	BMA	F	3	11/12	0.48	0.15	154,163,167,168	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MAN	D	4	11/12	0.62	0.10	142,155,158,158	0
5	MAN	F	4	11/12	0.63	0.13	140,148,151,154	0
4	MAN	E	4	11/12	0.66	0.15	130,139,141,142	0
4	MAN	D	5	11/12	0.70	0.12	144,153,156,156	0
3	NAG	C	2	14/15	0.72	0.11	138,144,152,154	0
5	BMA	F	5	11/12	0.73	0.13	118,127,134,138	0
5	NAG	F	2	14/15	0.76	0.15	123,140,146,153	0
6	NGS	G	1	19/19	0.80	0.10	127,134,145,148	0
4	BMA	E	3	11/12	0.82	0.11	124,131,137,138	0
3	NAG	C	1	14/15	0.83	0.12	110,117,129,136	0
4	NAG	E	1	14/15	0.84	0.13	110,113,116,118	0
4	NAG	D	1	14/15	0.85	0.12	92,103,117,130	0
5	NAG	F	1	14/15	0.86	0.13	105,113,119,128	0
6	GAL	G	2	11/12	0.89	0.09	108,115,124,124	0
4	NAG	E	2	14/15	0.93	0.09	100,112,119,119	0
6	SIA	G	3	20/21	0.93	0.09	93,101,106,108	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NAG	A	601	14/15	0.71	0.17	104,118,122,123	0
8	SO4	A	1327	5/5	0.73	0.08	139,141,145,146	0
8	SO4	A	1328	5/5	0.76	0.15	123,132,135,135	0
8	SO4	B	1176	5/5	0.93	0.12	81,84,92,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.