



Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 12:53 PM UTC

PDB ID : 6UOX / pdb_00006uox
EMDB ID : EMD-20834
Title : Structure of itraconazole-bound NPC1
Authors : Long, T.; Li, X.
Deposited on : 2019-10-15
Resolution : 4.02 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6727 (3.52 - 4.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1286	14% 88% 10%
2	B	2	100%
3	C	3	33% 67%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NPC intracellular cholesterol transporter 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1161	8909	5745	1448	1648	68	2	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1279	TRP	-	expression tag	UNP O15118
A	1280	SER	-	expression tag	UNP O15118
A	1281	HIS	-	expression tag	UNP O15118
A	1282	PRO	-	expression tag	UNP O15118
A	1283	GLN	-	expression tag	UNP O15118
A	1284	PHE	-	expression tag	UNP O15118
A	1285	GLU	-	expression tag	UNP O15118
A	1286	LYS	-	expression tag	UNP O15118

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	B	2	28	16	2	10	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	


- Molecule 5 is 4-(3-bromo-4-{4-[4-({(2R,4S)-2-(2,4-dichlorophenyl)-2-[(1H-1,2,4-triazol-1-yl)methyl]-1,3-dioxolan-4-yl}methoxy)phenyl]piperazin-1-yl}phenyl)-2-[(2S)-butan-2-yl]-2,4-dihydro-3H-1,2,4-triazol-3-one (CCD ID: QDG) (formula: $C_{35}H_{37}BrCl_2N_8O_4$).



Mol	Chain	Residues	Atoms						AltConf
			Total	Br	C	Cl	N	O	
5	A	1	50	1	35	2	8	4	0

MAG1
MAG2

- Molecule 3: α -D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 

MAG1
MAG2
MAN3

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	209612	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	80	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.033	Depositor
Minimum map value	-0.015	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0069	Depositor
Map size (Å)	231.00002, 231.00002, 231.00002	wwPDB
Map dimensions	350, 350, 350	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.66, 0.66, 0.66	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, QDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/9142	0.80	3/12460 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	PRO	N-CA-CB	6.86	110.45	103.25
1	A	433	PRO	N-CA-C	6.18	118.24	110.70
1	A	552	ASP	N-CA-C	5.23	121.94	110.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	253	PRO	Peptide
1	A	376	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8909	0	8577	14	0
2	B	28	0	25	0	0
3	C	39	0	34	0	0
4	A	112	0	104	0	0
5	A	50	0	0	1	0
All	All	9138	0	8740	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:624:VAL:HG11	5:A:1314:QDG:C05	2.28	0.64
1:A:1065:VAL:HG23	1:A:1078:VAL:HG21	1.89	0.55
1:A:1011:CYS:SG	1:A:1012:GLY:N	2.81	0.53
1:A:546:VAL:HG13	1:A:547:LEU:HG	1.93	0.49
1:A:1130:THR:HA	1:A:1133:MET:HG2	1.98	0.45
1:A:173:LEU:O	1:A:177:CYS:N	2.47	0.44
1:A:741:SER:HA	1:A:1112:VAL:HG11	1.99	0.44
1:A:1023:VAL:HG12	1:A:1036:THR:HG22	2.00	0.44
1:A:958:ARG:HB2	1:A:967:CYS:HB2	2.00	0.43
1:A:893:LEU:HD11	1:A:1069:MET:HE1	2.02	0.42
1:A:119:GLN:HB3	1:A:218:GLU:HB3	2.02	0.41
1:A:1114:MET:HE1	1:A:1121:LEU:HA	2.02	0.41
1:A:369:VAL:HA	1:A:372:ARG:HD2	2.04	0.40
1:A:699:VAL:HG11	1:A:1204:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1155/1286 (90%)	1061 (92%)	94 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	968/1117 (87%)	968 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
1	A	119	GLN
1	A	154	ASN
1	A	376	ASN
1	A	490	ASN
1	A	614	ASN
1	A	641	HIS
1	A	707	GLN
1	A	722	GLN
1	A	925	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.32	0	17,19,21	0.73	0
2	NAG	B	2	2	14,14,15	0.54	0	17,19,21	0.76	0
3	NAG	C	1	1,3	14,14,15	0.32	0	17,19,21	0.71	0
3	NAG	C	2	3	14,14,15	0.44	0	17,19,21	1.20	3 (17%)
3	MAN	C	3	3	11,11,12	0.42	0	15,15,17	1.31	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
3	NAG	C	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	MAN	C	3	3	-	1/2/19/22	1/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	MAN	C1-O5-C5	4.51	118.23	112.19
3	C	2	NAG	C3-C4-C5	2.42	114.63	110.23
3	C	2	NAG	C4-C3-C2	2.20	114.25	111.02
3	C	2	NAG	O5-C1-C2	-2.00	108.20	111.29

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	O5-C5-C6-O6
3	C	3	MAN	O5-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	3	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1312	1	14,14,15	0.30	0	17,19,21	0.81	0
5	QDG	A	1314	-	55,56,56	2.78	12 (21%)	77,81,81	2.07	23 (29%)
4	NAG	A	1311	1	14,14,15	0.33	0	17,19,21	0.72	0
4	NAG	A	1305	1	14,14,15	0.30	0	17,19,21	0.73	0
4	NAG	A	1310	1	14,14,15	0.28	0	17,19,21	0.75	0
4	NAG	A	1301	1	14,14,15	0.27	0	17,19,21	0.75	0
4	NAG	A	1309	1	14,14,15	0.25	0	17,19,21	0.69	0
4	NAG	A	1313	1	14,14,15	0.46	0	17,19,21	0.90	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1302	1	14,14,15	0.37	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1312	1	-	2/6/23/26	0/1/1/1
5	QDG	A	1314	-	-	12/34/55/55	0/7/7/7
4	NAG	A	1311	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1305	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1310	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1301	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1309	1	-	1/6/23/26	0/1/1/1
4	NAG	A	1313	1	-	1/6/23/26	0/1/1/1
4	NAG	A	1302	1	-	2/6/23/26	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1314	QDG	C49-C17	-9.91	1.21	1.51
5	A	1314	QDG	C45-N44	9.06	1.45	1.30
5	A	1314	QDG	O16-C09	-8.19	1.20	1.42
5	A	1314	QDG	C09-C03	6.79	1.61	1.53
5	A	1314	QDG	C38-N39	5.64	1.47	1.36
5	A	1314	QDG	C45-N37	3.99	1.43	1.37
5	A	1314	QDG	C38-N37	3.85	1.44	1.39
5	A	1314	QDG	O16-C17	3.73	1.56	1.45
5	A	1314	QDG	N39-N44	3.71	1.43	1.37
5	A	1314	QDG	C02-CL01	2.22	1.78	1.73
5	A	1314	QDG	C23-N26	2.21	1.44	1.38
5	A	1314	QDG	BR32-C31	2.16	1.94	1.89

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1314	QDG	C38-N39-N44	-7.75	106.78	112.87
5	A	1314	QDG	C45-N44-N39	5.76	109.37	103.68
5	A	1314	QDG	C10-C09-C03	4.44	115.83	109.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1314	QDG	N37-C38-N39	4.30	107.73	102.88
5	A	1314	QDG	C22-C23-N26	-3.75	116.18	121.39
5	A	1314	QDG	C10-N11-N12	3.72	126.67	121.25
5	A	1314	QDG	C03-C02-CL01	3.18	125.55	121.70
5	A	1314	QDG	C47-N29-C28	2.83	117.93	111.57
5	A	1314	QDG	C27-C28-N29	2.80	116.67	110.78
5	A	1314	QDG	C18-O19-C20	-2.77	111.75	117.85
5	A	1314	QDG	C08-C02-CL01	-2.70	114.03	118.45
5	A	1314	QDG	BR32-C31-C30	2.63	123.74	120.53
5	A	1314	QDG	O50-C49-C17	2.61	108.08	103.26
4	A	1313	NAG	O5-C1-C2	-2.56	107.33	111.29
5	A	1314	QDG	N12-C13-N14	-2.49	107.81	114.90
5	A	1314	QDG	C13-N12-N11	2.44	108.23	102.60
5	A	1314	QDG	C34-C30-N29	-2.28	118.85	122.42
5	A	1314	QDG	O50-C09-C03	-2.26	107.49	111.37
5	A	1314	QDG	O50-C09-O16	-2.19	101.98	106.02
5	A	1314	QDG	C33-C31-C30	-2.19	118.96	121.34
5	A	1314	QDG	C04-C05-C06	2.18	121.44	119.24
5	A	1314	QDG	N37-C45-N44	-2.16	107.34	111.42
4	A	1302	NAG	C1-O5-C5	2.04	114.92	112.19
5	A	1314	QDG	C45-N37-C38	2.02	108.77	107.27
5	A	1314	QDG	O46-C38-N37	-2.01	125.12	128.10

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1314	QDG	C03-C09-C10-N11
5	A	1314	QDG	O16-C09-C10-N11
5	A	1314	QDG	C42-C40-N39-C38
5	A	1314	QDG	C41-C40-N39-N44
4	A	1312	NAG	O5-C5-C6-O6
4	A	1302	NAG	O5-C5-C6-O6
4	A	1311	NAG	O5-C5-C6-O6
4	A	1311	NAG	C4-C5-C6-O6
4	A	1312	NAG	C4-C5-C6-O6
5	A	1314	QDG	C25-C20-O19-C18
5	A	1314	QDG	C21-C20-O19-C18
4	A	1302	NAG	C4-C5-C6-O6
4	A	1305	NAG	O5-C5-C6-O6
5	A	1314	QDG	C41-C40-C42-C43
5	A	1314	QDG	N39-C40-C42-C43

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	1314	QDG	C24-C23-N26-C48
4	A	1309	NAG	O5-C5-C6-O6
4	A	1313	NAG	O5-C5-C6-O6
5	A	1314	QDG	C17-C18-O19-C20
5	A	1314	QDG	C22-C23-N26-C48
4	A	1301	NAG	C4-C5-C6-O6
4	A	1305	NAG	C4-C5-C6-O6
5	A	1314	QDG	C24-C23-N26-C27
4	A	1301	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1314	QDG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20834. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 175



Y Index: 175




Z Index: 175


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 156



Y Index: 196




Z Index: 149


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0069. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 56 nm^3 ; this corresponds to an approximate mass of 51 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.249 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-20834 and PDB model 6UOX. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.0069 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0069) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6250	0.3720
A	0.6250	0.3720
B	0.5710	0.3500
C	0.6410	0.4230

