



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:15 AM EDT

PDB ID : 5UQD / pdb_00005uqd
Title : DPY-21 in complex with Fe(II) and alpha-Ketoglutarate
Authors : Brejc, K.; Bian, Q.; Uzawa, S.; Wheeler, B.S.; Anderson, E.C.; King, D.S.;
Kranzusch, P.J.; Preston, C.G.; Meyer, B.J.
Deposited on : 2017-02-07
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DumPY: shorter than wild-type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	324	2703	1711	490	478	24	0	11	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1176	MET	-	initiating methionine	UNP Q9GRZ3
A	1177	LYS	-	expression tag	UNP Q9GRZ3
A	1178	SER	-	expression tag	UNP Q9GRZ3
A	1179	SER	-	expression tag	UNP Q9GRZ3
A	1180	TRP	-	expression tag	UNP Q9GRZ3
A	1181	SER	-	expression tag	UNP Q9GRZ3
A	1182	HIS	-	expression tag	UNP Q9GRZ3
A	1183	PRO	-	expression tag	UNP Q9GRZ3
A	1184	GLN	-	expression tag	UNP Q9GRZ3
A	1185	PHE	-	expression tag	UNP Q9GRZ3
A	1186	GLU	-	expression tag	UNP Q9GRZ3
A	1187	LYS	-	expression tag	UNP Q9GRZ3
A	1188	GLY	-	expression tag	UNP Q9GRZ3
A	1189	ALA	-	expression tag	UNP Q9GRZ3
A	1190	MET	-	expression tag	UNP Q9GRZ3
A	1191	THR	-	expression tag	UNP Q9GRZ3
A	1192	GLY	-	expression tag	UNP Q9GRZ3
A	1193	TRP	-	expression tag	UNP Q9GRZ3
A	1194	SER	-	expression tag	UNP Q9GRZ3
A	1195	HIS	-	expression tag	UNP Q9GRZ3
A	1196	PRO	-	expression tag	UNP Q9GRZ3
A	1197	GLN	-	expression tag	UNP Q9GRZ3
A	1198	PHE	-	expression tag	UNP Q9GRZ3
A	1199	GLU	-	expression tag	UNP Q9GRZ3
A	1200	LYS	-	expression tag	UNP Q9GRZ3
A	1201	GLU	-	expression tag	UNP Q9GRZ3
A	1202	ASN	-	expression tag	UNP Q9GRZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1203	LEU	-	expression tag	UNP Q9GRZ3
A	1204	TYR	-	expression tag	UNP Q9GRZ3
A	1205	PHE	-	expression tag	UNP Q9GRZ3
A	1206	GLN	-	expression tag	UNP Q9GRZ3
A	1207	SER	-	expression tag	UNP Q9GRZ3
A	1208	ASN	-	expression tag	UNP Q9GRZ3
A	1209	ALA	-	expression tag	UNP Q9GRZ3

- Molecule 2 is 2-OXOGLUTARIC ACID (CCD ID: AKG) (formula: C₅H₆O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is 2-(2-METHOXYETHOXY)ETHANOL (CCD ID: PG0) (formula: C₅H₁₂O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	5	3		
4	A	1	Total	C	O	0	0
			8	5	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	254	Total	O	0	1
			255	255		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- [illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	87.62Å 97.88Å 98.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.86 – 1.80 43.86 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (43.86-1.80) 99.4 (43.86-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.163 , 0.193 0.165 , 0.194	Depositor DCC
R_{free} test set	1978 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2985	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG0, FE2, AKG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/2777 (0.0%)	0.70	0/3743

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1293	MET	SD-CE	-10.99	1.52	1.79

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2703	0	2662	14	0
2	A	10	0	4	0	0
3	A	1	0	0	0	0
4	A	16	0	24	1	0
5	A	255	0	0	0	0
All	All	2985	0	2690	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1448[B]:ARG:HG2	1:A:1558:TRP:CH2	2.27	0.70
1:A:1446:ASN:HB3	1:A:1448[B]:ARG:HD3	1.78	0.65
1:A:1539:VAL:HG13	1:A:1567:MET:HE1	1.90	0.54
1:A:1211:MET:HE2	1:A:1236:LYS:HB2	1.90	0.52
1:A:1358[B]:MET:HE1	1:A:1373:MET:SD	2.53	0.48
1:A:1448[B]:ARG:HG2	1:A:1558:TRP:CZ3	2.49	0.47
1:A:1319:GLY:O	1:A:1351:GLY:HA3	2.17	0.45
1:A:1418:VAL:HG22	1:A:1443:THR:O	2.18	0.44
1:A:1448[A]:ARG:HB3	1:A:1558:TRP:CZ3	2.53	0.44
1:A:1369:ASN:C	1:A:1371:LYS:H	2.28	0.41
1:A:1285:VAL:HG11	1:A:1534[B]:HIS:CG	2.56	0.41
1:A:1369:ASN:C	1:A:1371:LYS:N	2.76	0.41
1:A:1269:HIS:HA	4:A:1704:PG0:H12	2.02	0.40
1:A:1528[B]:VAL:HG13	1:A:1577:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/442 (73%)	320 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/389 (76%)	290 (99%)	4 (1%)	59	52

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1267	MET
1	A	1293	MET
1	A	1378	ASN
1	A	1404	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1303	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AKG	A	1701	3	9,9,9	2.06	3 (33%)	11,11,11	2.72	7 (63%)
4	PG0	A	1703	-	7,7,7	0.45	0	6,6,6	0.30	0
4	PG0	A	1704	-	7,7,7	0.48	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AKG	A	1701	3	-	1/9/9/9	-
4	PG0	A	1703	-	-	0/5/5/5	-
4	PG0	A	1704	-	-	2/5/5/5	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1701	AKG	O1-C1	3.44	1.31	1.22
2	A	1701	AKG	O4-C5	-2.92	1.21	1.30
2	A	1701	AKG	C3-C2	2.83	1.54	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1701	AKG	C4-C3-C2	-4.20	105.04	112.91
2	A	1701	AKG	C3-C2-C1	4.01	122.68	115.86
2	A	1701	AKG	O5-C2-C3	-3.44	113.79	121.17
2	A	1701	AKG	C3-C4-C5	-3.35	104.78	113.67
2	A	1701	AKG	O2-C1-O1	-2.84	117.13	123.90
2	A	1701	AKG	O5-C2-C1	2.50	123.24	119.67
2	A	1701	AKG	O2-C1-C2	2.14	119.51	113.59

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1704	PG0	O1-C3-C4-O2
4	A	1704	PG0	C4-C3-O1-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1701	AKG	C3-C4-C5-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1704	PG0	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/442 (73%)	-0.01	15 (4%) 37 36	7, 23, 50, 66	11 (3%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1369	ASN	5.3
1	A	1395	VAL	3.7
1	A	1232	SER	3.2
1	A	1548	TYR	3.2
1	A	1402	THR	3.1
1	A	1326	PHE	3.1
1	A	1404	ASP	2.6
1	A	1370	PHE	2.5
1	A	1360	GLY	2.5
1	A	1457	ARG	2.4
1	A	1403	SER	2.4
1	A	1509	PRO	2.3
1	A	1325	GLN	2.2
1	A	1610	LEU	2.2
1	A	1233	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PG0	A	1704	8/8	0.75	0.24	72,75,76,76	0
2	AKG	A	1701	10/10	0.86	0.13	23,29,32,41	0
4	PG0	A	1703	8/8	0.91	0.12	45,46,57,59	0
3	FE2	A	1702	1/1	1.00	0.02	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.