



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:35 AM UTC

PDB ID : 8UQ8 / pdb_00008uq8
Title : Crystal structure of RNF168 (RING)-UbcH5c fused to H2A-H2B via a 2-residue linker
Authors : Hu, Q.; Botuyan, M.V.; Mer, G.
Deposited on : 2023-10-23
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	
1	a	437	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13815 atoms, of which 6850 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase RNF168, Ubiquitin-conjugating enzyme E2 D3, Histone H2B type 2-E, Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	426	Total	C	H	N	O	S	263	2	0
			6782	2124	3415	613	609	21			
1	a	423	Total	C	H	N	O	S	137	1	0
			6743	2112	3397	608	605	21			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q8IYW5
A	0	HIS	-	expression tag	UNP Q8IYW5
A	95	SER	-	linker	UNP Q8IYW5
A	96	GLY	-	linker	UNP Q8IYW5
A	97	SER	-	linker	UNP Q8IYW5
A	98	GLY	-	linker	UNP Q8IYW5
A	99	SER	-	linker	UNP Q8IYW5
A	100	GLY	-	linker	UNP Q8IYW5
A	101	SER	-	linker	UNP Q8IYW5
A	1148	GLY	-	linker	UNP P61077
A	1149	SER	-	linker	UNP P61077
A	3011	SER	-	linker	UNP Q16778
a	-1	GLY	-	expression tag	UNP Q8IYW5
a	0	HIS	-	expression tag	UNP Q8IYW5
a	95	SER	-	linker	UNP Q8IYW5
a	96	GLY	-	linker	UNP Q8IYW5
a	97	SER	-	linker	UNP Q8IYW5
a	98	GLY	-	linker	UNP Q8IYW5
a	99	SER	-	linker	UNP Q8IYW5
a	100	GLY	-	linker	UNP Q8IYW5
a	101	SER	-	linker	UNP Q8IYW5
a	1148	GLY	-	linker	UNP P61077
a	1149	SER	-	linker	UNP P61077
a	3011	SER	-	linker	UNP Q16778

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	7	Total	Cl	0	0
			7	7		
2	a	14	Total	Cl	0	0
			14	14		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			13	3	7	3		
3	a	1	Total	C	H	O	0	0
			13	3	7	3		
3	a	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	a	2	Total	Zn	0	0
			2	2		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Na	0	0
			3	3		
5	a	2	Total	Na	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	94	Total	O	0	0
			94	94		
6	a	98	Total	O	0	0
			98	98		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 ubiquitin-protein ligase RNF168, Ubiquitin-conjugating enzyme E2 D3, Histone H2B type 2-E, Histone H2A type 1-B/E



- Molecule 1: E3 ubiquitin-protein ligase RNF168, Ubiquitin-conjugating enzyme E2 D3, Histone H2B type 2-E, Histone H2A type 1-B/E



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	107.54Å 107.54Å 113.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.06 – 2.34 36.06 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.06-2.34) 99.4 (36.06-2.34)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.18 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.196 , 0.236 (Not available) , 0.209	Depositor DCC
R_{free} test set	1992 reflections (3.21%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.730	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.488 for -h,-k,l 0.042 for h,-h-k,-l 0.042 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13815	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/3448	0.33	0/4669
1	a	0.16	0/3423	0.31	0/4636
All	All	0.16	0/6871	0.32	0/9305

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3367	3415	3416	19	0
1	a	3346	3397	3398	8	0
2	A	7	0	0	0	0
2	a	14	0	0	2	0
3	A	18	23	24	1	0
3	a	12	15	16	0	0
4	A	2	0	0	0	0
4	a	2	0	0	0	0
5	A	3	0	0	0	0
5	a	2	0	0	0	0
6	A	94	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	a	98	0	0	1	0
All	All	6965	6850	6854	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3073:ASN:O	1:A:3073:ASN:ND2	2.12	0.82
1:A:3073:ASN:OD1	1:A:3081:ARG:NH1	2.29	0.65
1:A:2118:VAL:O	1:A:2122:THR:HG23	1.99	0.61
1:a:3073:ASN:OD1	1:a:3075:LYS:HG3	2.08	0.53
2:a:4012:CL:CL	6:a:5035:HOH:O	2.57	0.51
1:A:1026:VAL:HG12	1:A:1034:GLN:HB2	1.94	0.49
1:A:36:CYS:SG	1:A:91:ARG:NH2	2.84	0.48
1:A:1042:ASP:O	1:A:1042:ASP:OD1	2.32	0.48
1:a:3073:ASN:ND2	1:a:3081:ARG:HH21	2.11	0.48
1:A:24:VAL:HG11	1:A:91:ARG:HG2	1.96	0.48
1:a:35:LEU:HD23	1:a:35:LEU:N	2.29	0.47
1:a:42:SER:OG	1:a:99:SER:O	2.30	0.46
1:A:1032:HIS:CE1	1:A:1055[A]:HIS:CD2	3.04	0.46
1:A:3066:ALA:CB	1:A:3083:LEU:HD12	2.45	0.46
1:A:34:THR:O	3:A:4006:GOL:H31	2.16	0.46
1:a:5:LYS:HG3	2:a:4003:CL:CL	2.53	0.45
1:A:1030:MET:HA	1:A:1030:MET:HE2	1.99	0.44
1:a:3073:ASN:OD1	1:a:3073:ASN:O	2.36	0.44
1:A:3073:ASN:O	1:A:3073:ASN:CG	2.60	0.43
1:A:3065:LEU:HB3	1:A:3086:ALA:HB1	2.00	0.43
1:A:1066:LYS:HA	1:A:1066:LYS:HE3	1.99	0.43
1:a:67:ARG:HG3	1:a:67:ARG:NH1	2.33	0.43
1:A:35:LEU:N	1:A:35:LEU:HD23	2.34	0.42
1:A:3065:LEU:CB	1:A:3086:ALA:HB1	2.50	0.42
1:a:29:LEU:HB3	1:a:30:PRO:HD2	2.02	0.41
1:A:1088:ILE:CG1	1:A:1097:LEU:HD13	2.50	0.41
1:A:2070:PHE:CD1	1:A:2070:PHE:C	2.99	0.40
1:A:24:VAL:CG1	1:A:91:ARG:HG2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/437 (97%)	410 (97%)	14 (3%)	0	100	100
1	a	420/437 (96%)	408 (97%)	12 (3%)	0	100	100
All	All	844/874 (97%)	818 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/377 (99%)	373 (100%)	0	100	100
1	a	371/377 (98%)	371 (100%)	0	100	100
All	All	744/754 (99%)	744 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1032	HIS
1	A	2047	GLN
1	A	2095	GLN
1	A	3024	GLN
1	a	32	ASN
1	a	1032	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	a	2082	HIS
1	a	2084	ASN
1	a	2095	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 35 ligands modelled in this entry, 30 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	4005	-	5,5,5	0.83	0	5,5,5	0.98	0
3	GOL	a	4009	-	5,5,5	1.04	0	5,5,5	0.89	0
3	GOL	A	4006	-	5,5,5	0.95	0	5,5,5	0.75	0
3	GOL	A	4010	-	5,5,5	0.91	0	5,5,5	1.13	1 (20%)
3	GOL	a	4008	-	5,5,5	0.95	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	4005	-	-	0/4/4/4	-
3	GOL	a	4009	-	-	2/4/4/4	-
3	GOL	A	4006	-	-	4/4/4/4	-
3	GOL	A	4010	-	-	2/4/4/4	-
3	GOL	a	4008	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	4010	GOL	C3-C2-C1	-2.06	104.24	111.80

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	4006	GOL	C1-C2-C3-O3
3	A	4010	GOL	C1-C2-C3-O3
3	a	4009	GOL	C1-C2-C3-O3
3	a	4009	GOL	O2-C2-C3-O3
3	A	4006	GOL	O1-C1-C2-C3
3	A	4006	GOL	O2-C2-C3-O3
3	A	4010	GOL	O2-C2-C3-O3
3	A	4006	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	4006	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/437 (97%)	0.38	34 (7%) 18 22	6, 45, 77, 105	27 (6%)
1	a	423/437 (96%)	0.35	33 (7%) 19 23	18, 45, 76, 103	14 (3%)
All	All	849/874 (97%)	0.37	67 (7%) 18 22	6, 45, 76, 105	41 (4%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	a	0	HIS	6.7
1	A	3099	ARG	6.7
1	A	100	GLY	5.5
1	A	98	GLY	5.0
1	A	99	SER	4.8
1	A	91	ARG	4.8
1	a	100	GLY	4.7
1	a	3099	ARG	4.4
1	A	0	HIS	4.0
1	a	1020	GLN	3.7
1	a	3073	ASN	3.6
1	A	3038	ASN	3.5
1	A	-1	GLY	3.4
1	A	2085	LYS	3.3
1	a	47	ALA	3.2
1	A	1015	ARG	3.2
1	A	3077	ARG	3.2
1	a	1015	ARG	3.2
1	a	3079	ILE	3.1
1	A	1016	ASP	3.1
1	A	3085	LEU	3.1
1	a	3038	ASN	3.0
1	a	3072	ASP	3.0
1	a	2086	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	a	99	SER	2.9
1	a	3085	LEU	2.8
1	a	1018	PRO	2.7
1	A	3076	THR	2.6
1	a	1019	ALA	2.6
1	a	2082	HIS	2.6
1	A	3073	ASN	2.6
1	a	3077	ARG	2.5
1	A	3074	LYS	2.5
1	A	1042	ASP	2.4
1	a	1016	ASP	2.4
1	A	3078	ILE	2.4
1	A	1019	ALA	2.4
1	a	2	ALA	2.4
1	a	46	LYS	2.4
1	a	3074	LYS	2.4
1	A	2057	LYS	2.4
1	a	1145	TYR	2.3
1	a	1	MET	2.3
1	A	2050	PRO	2.3
1	a	3076	THR	2.3
1	A	3079	ILE	2.3
1	A	97	SER	2.3
1	a	1021	CYS	2.3
1	a	3075	LYS	2.3
1	A	1055[A]	HIS	2.2
1	A	1037	ILE	2.2
1	a	3071	ARG	2.2
1	A	1145	TYR	2.2
1	a	90	LEU	2.2
1	A	1028	ASP	2.2
1	A	1032	HIS	2.2
1	a	3081	ARG	2.2
1	A	1018	PRO	2.1
1	A	1020	GLN	2.1
1	A	2086	ARG	2.1
1	a	2083	TYR	2.1
1	A	3072	ASP	2.1
1	a	1042	ASP	2.1
1	A	3071	ARG	2.1
1	A	1021	CYS	2.1
1	a	3098	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	a	1014	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	a	4005	1/1	0.77	0.28	81,81,81,81	0
3	GOL	a	4008	6/6	0.78	0.20	57,73,90,98	0
2	CL	a	4012	1/1	0.79	0.40	95,95,95,95	0
3	GOL	A	4006	6/6	0.80	0.22	40,58,75,75	0
2	CL	a	4019	1/1	0.81	0.20	82,82,82,82	0
2	CL	A	4015	1/1	0.82	0.26	82,82,82,82	0
3	GOL	a	4009	6/6	0.83	0.19	39,53,71,72	0
2	CL	A	4003	1/1	0.84	0.18	73,73,73,73	0
2	CL	a	4007	1/1	0.87	0.21	75,75,75,75	0
2	CL	a	4004	1/1	0.87	0.20	74,74,74,74	0
2	CL	A	4004	1/1	0.87	0.17	67,67,67,67	0
2	CL	a	4015	1/1	0.88	0.18	67,67,67,67	0
3	GOL	A	4010	6/6	0.90	0.13	48,69,83,93	0
3	GOL	A	4005	6/6	0.90	0.15	52,67,85,85	0
2	CL	a	4020	1/1	0.90	0.11	77,77,77,77	0
2	CL	a	4013	1/1	0.91	0.29	59,59,59,59	0
5	NA	A	4009	1/1	0.91	0.11	33,33,33,33	0
5	NA	a	4016	1/1	0.91	0.17	50,50,50,50	0
2	CL	a	4006	1/1	0.92	0.13	76,76,76,76	0
5	NA	a	4017	1/1	0.93	0.08	54,54,54,54	0
2	CL	A	4014	1/1	0.94	0.10	54,54,54,54	0
5	NA	A	4011	1/1	0.94	0.12	48,48,48,48	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	a	4018	1/1	0.95	0.11	48,48,48,48	0
2	CL	a	4003	1/1	0.95	0.11	50,50,50,50	0
5	NA	A	4012	1/1	0.96	0.09	49,49,49,49	0
2	CL	A	4013	1/1	0.97	0.05	36,36,36,36	0
2	CL	a	4002	1/1	0.98	0.06	43,43,43,43	0
2	CL	a	4014	1/1	0.98	0.06	34,34,34,34	0
2	CL	a	4001	1/1	0.99	0.04	37,37,37,37	0
2	CL	A	4002	1/1	0.99	0.05	43,43,43,43	0
2	CL	A	4001	1/1	0.99	0.03	37,37,37,37	0
4	ZN	A	4008	1/1	1.00	0.01	30,30,30,30	0
4	ZN	a	4010	1/1	1.00	0.01	32,32,32,32	0
4	ZN	a	4011	1/1	1.00	0.03	32,32,32,32	0
4	ZN	A	4007	1/1	1.00	0.03	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 4008:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN a 4010:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN a 4011:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 4007:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.