



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:46 PM UTC

PDB ID : 3UR5 / pdb_00003ur5
Title : Crystal Structure of PTE mutant K185R/I274N
Authors : Tsai, P.; Fox, N.G.; Li, Y.; Barondeau, D.P.; Raushel, F.M.
Deposited on : 2011-11-21
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	13% 90% 9% .
1	B	327	8% 91% 7% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Parathion hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2504	1580	447	470	7			
1	B	323	Total	C	N	O	S	0	0	0
			2471	1561	442	461	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	ARG	LYS	engineered mutation	UNP P0A434
A	274	ASN	ILE	engineered mutation	UNP P0A434
B	185	ARG	LYS	engineered mutation	UNP P0A434
B	274	ASN	ILE	engineered mutation	UNP P0A434

- Molecule 2 is DIETHYL HYDROGEN PHOSPHATE (CCD ID: DPF) (formula: $C_4H_{11}O_4P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			9	4	4	1		
2	B	1	Total	C	O	P	0	0
			9	4	4	1		

- Molecule 3 is COBALT (II) ION (CCD ID: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Co	0	0
			2	2		
3	B	3	Total	Co	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	303	Total	O	0	0
			303	303		
4	B	305	Total	O	0	0
			305	305		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Parathion hydrolase



• Molecule 1: Parathion hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.48Å 85.98Å 88.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 50.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-1.60) 100.0 (50.00-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.85 (at 1.60Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.209 , 0.228 0.208 , 0.223	Depositor DCC
R_{free} test set	4329 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 26.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for -h,l,k 0.012 for -l,-k,-h 0.013 for k,h,-l 0.001 for k,l,h 0.001 for l,h,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5606	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DPF, KCX, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	2/2539 (0.1%)	1.11	15/3447 (0.4%)
1	B	0.70	3/2505 (0.1%)	1.08	14/3399 (0.4%)
All	All	0.70	5/5044 (0.1%)	1.09	29/6846 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	255	ILE	CA-CB	9.33	1.58	1.54
1	B	328	ILE	CA-CB	9.17	1.59	1.54
1	B	255	ILE	CA-CB	7.10	1.57	1.54
1	B	333	ILE	CA-CB	6.86	1.58	1.54
1	A	333	ILE	CA-CB	5.11	1.60	1.54

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	106	ILE	N-CA-C	-8.51	104.40	112.83
1	B	54	THR	N-CA-C	8.46	123.42	113.18
1	A	106	ILE	N-CA-C	-8.28	104.51	112.29
1	A	54	THR	N-CA-C	8.25	123.16	113.18
1	A	313	ILE	N-CA-C	8.17	119.94	110.62
1	A	58	ILE	N-CA-C	-7.87	103.24	111.58
1	B	36	ARG	N-CA-C	7.33	121.39	109.59
1	B	132	PHE	N-CA-C	6.95	120.71	112.93
1	B	203	ALA	N-CA-C	-6.44	94.41	107.41
1	B	58	ILE	N-CA-C	-6.32	104.12	111.00
1	A	201	HIS	N-CA-C	-6.31	99.43	109.59
1	A	203	ALA	N-CA-C	-6.30	94.69	107.41
1	A	311	THR	N-CA-C	6.24	118.08	111.28
1	A	132	PHE	N-CA-C	6.22	119.89	112.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	200	THR	N-CA-C	6.12	119.52	109.85
1	B	311	THR	N-CA-C	6.11	121.39	111.37
1	A	200	THR	N-CA-C	5.99	119.31	109.85
1	B	171	ALA	N-CA-C	5.75	118.01	108.99
1	A	253	ASP	N-CA-C	5.64	120.85	113.30
1	A	139	ARG	N-CA-C	5.60	119.73	113.01
1	A	226	VAL	N-CA-C	5.58	116.16	108.12
1	B	226	VAL	N-CA-C	5.58	116.16	108.12
1	B	155	GLN	N-CA-C	5.53	118.49	111.69
1	A	46	ILE	N-CA-C	5.33	116.70	110.62
1	B	253	ASP	N-CA-C	5.22	120.29	113.30
1	B	201	HIS	N-CA-C	-5.15	100.85	109.24
1	A	252	LEU	N-CA-C	-5.10	102.51	109.71
1	A	171	ALA	N-CA-C	5.04	116.90	108.99
1	B	139	ARG	N-CA-C	5.03	119.04	113.01

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2504	0	2511	5	0
1	B	2471	0	2483	5	0
2	A	9	0	10	3	0
2	B	9	0	10	4	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
4	A	303	0	0	2	0
4	B	305	0	0	2	0
All	All	5606	0	5014	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:901:DPF:H21	4:B:1445:HOH:O	1.81	0.79
2:B:901:DPF:H31	4:B:1593:HOH:O	1.89	0.72
1:A:60:GLY:HA3	2:A:902:DPF:H41	1.96	0.47
2:A:902:DPF:H43	4:A:1007:HOH:O	2.14	0.47
2:A:902:DPF:H32	4:A:1313:HOH:O	2.15	0.45
1:B:333:ILE:HG23	1:B:346:LEU:HD13	1.98	0.45
1:A:131:TRP:CG	1:A:132:PHE:H	2.35	0.44
1:B:254:HIS:NE2	2:B:901:DPF:H32	2.33	0.43
1:B:131:TRP:CG	1:B:132:PHE:H	2.37	0.42
1:A:80:ALA:O	1:A:84:VAL:HG23	2.20	0.42
1:B:106:ILE:O	1:B:106:ILE:HG22	2.20	0.42
1:A:106:ILE:HG22	1:A:106:ILE:O	2.21	0.41
1:A:180:GLN:O	1:A:184:LEU:HD13	2.20	0.41
1:B:131:TRP:CZ2	2:B:901:DPF:H12	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/327 (99%)	314 (97%)	10 (3%)	0	100	100
1	B	318/327 (97%)	306 (96%)	12 (4%)	0	100	100
All	All	642/654 (98%)	620 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/262 (100%)	252 (96%)	10 (4%)	29	9
1	B	258/262 (98%)	254 (98%)	4 (2%)	55	33
All	All	520/524 (99%)	506 (97%)	14 (3%)	39	16

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	37	ILE
1	A	46	ILE
1	A	140	LEU
1	A	196	VAL
1	A	262	LEU
1	A	274	ASN
1	A	299	SER
1	A	303	LEU
1	A	344	GLU
1	B	178	PRO
1	B	196	VAL
1	B	299	SER
1	B	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	GLN
1	B	148	GLN
1	B	206	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	169	3,1	10,11,12	0.88	0	6,12,14	1.93	1 (16%)
1	KCX	B	169	3,1	10,11,12	0.90	1 (10%)	6,12,14	1.91	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	169	3,1	-	0/9/10/12	-
1	KCX	B	169	3,1	-	0/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	169	KCX	OQ1-CX	2.05	1.25	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	KCX	OQ1-CX-NZ	-4.61	117.92	124.92
1	B	169	KCX	OQ1-CX-NZ	-4.55	118.01	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DPF	A	902	3	8,8,8	0.77	0	9,10,10	0.50	0
2	DPF	B	901	3	8,8,8	0.76	0	9,10,10	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DPF	A	902	3	-	3/8/8/8	-
2	DPF	B	901	3	-	3/8/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	902	DPF	C1-O3-P1-O4
2	A	902	DPF	C1-O3-P1-O1
2	B	901	DPF	C2-O1-P1-O3
2	B	901	DPF	C3-C2-O1-P1
2	A	902	DPF	C1-O3-P1-O2
2	B	901	DPF	C2-O1-P1-O2

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	902	DPF	3	0
2	B	901	DPF	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/327 (99%)	0.87	43 (13%) 7 7	15, 26, 44, 56	0
1	B	322/327 (98%)	0.51	27 (8%) 17 17	13, 21, 43, 62	1 (0%)
All	All	648/654 (99%)	0.69	70 (10%) 11 10	13, 24, 43, 62	1 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	261	GLY	7.4
1	B	266	ALA	5.5
1	A	340	GLY	4.6
1	A	37	ILE	4.5
1	B	270	ALA	4.1
1	A	341	VAL	4.1
1	B	260	ILE	4.0
1	A	293	MET	3.9
1	B	267	SER	3.9
1	B	271	LEU	3.8
1	A	360	PRO	3.7
1	A	361	THR	3.7
1	A	44	ILE	3.6
1	B	338	GLU	3.6
1	B	268	ALA	3.5
1	A	335	PHE	3.5
1	A	342	PRO	3.5
1	A	332	VAL	3.4
1	A	336	LEU	3.3
1	A	274	ASN	3.3
1	B	269	SER	3.3
1	B	272	LEU	3.3
1	A	46	ILE	3.2
1	A	328	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	340	GLY	3.1
1	B	35	ASP	3.1
1	A	262	LEU	3.1
1	A	329	PRO	3.0
1	A	333	ILE	3.0
1	B	274	ASN	2.9
1	A	344	GLU	2.9
1	A	338	GLU	2.8
1	A	292	TYR	2.7
1	B	288	ILE	2.7
1	B	254	HIS	2.7
1	B	259	ALA	2.7
1	B	132	PHE	2.7
1	A	334	PRO	2.7
1	A	330	LEU	2.7
1	B	341	VAL	2.6
1	A	260	ILE	2.6
1	A	339	LYS	2.6
1	B	293	MET	2.6
1	A	281	ALA	2.6
1	B	312	ASN	2.6
1	A	288	ILE	2.5
1	B	292	TYR	2.5
1	B	286	ALA	2.4
1	A	45	THR	2.4
1	A	296	ILE	2.4
1	A	254	HIS	2.4
1	A	89	ARG	2.3
1	A	92	ALA	2.3
1	B	46	ILE	2.3
1	A	277	TRP	2.2
1	A	156	TYR	2.2
1	B	316	VAL	2.2
1	A	51	PHE	2.2
1	A	356	ARG	2.2
1	B	320	VAL	2.2
1	A	331	ARG	2.1
1	B	36	ARG	2.1
1	A	350	THR	2.1
1	A	343	GLN	2.1
1	B	273	GLY	2.1
1	A	36	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	282	LEU	2.1
1	A	351	VAL	2.0
1	A	345	THR	2.0
1	A	320	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	169	12/13	0.94	0.08	17,19,20,20	0
1	KCX	B	169	12/13	0.96	0.06	13,14,16,17	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DPF	A	902	9/9	0.63	0.25	40,44,46,46	0
2	DPF	B	901	9/9	0.83	0.18	21,31,37,37	0
3	CO	B	805	1/1	0.92	0.19	50,50,50,50	0
3	CO	A	801	1/1	0.98	0.04	20,20,20,20	0
3	CO	B	804	1/1	0.99	0.02	17,17,17,17	0
3	CO	A	802	1/1	1.00	0.02	20,20,20,20	0
3	CO	B	803	1/1	1.00	0.01	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.