



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 01:40 PM UTC

PDB ID : 3URI / pdb_00003uri
Title : Endothiapepsin-DB5 complex.
Authors : Bailey, D.; Sanz-Aparicio, J.; Albert, A.; Cooper, J.B.
Deposited on : 2011-11-22
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	 67% 31% .
2	B	8	 25% 25% 12% 25% 12%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endothiapepsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2388	1514	366	506	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	SUI	ASP	SEE REMARK 999	UNP P11838
A	54	SUI	GLY	SEE REMARK 999	UNP P11838

- Molecule 2 is a protein called DB5 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	S	0	0	0
			65	43	12	9	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	283	Total	O	0	0
			283	283		
3	B	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endothiapepsin



• Molecule 2: DB5 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.02Å 75.56Å 42.80Å 90.00° 97.02° 90.00°	Depositor
Resolution (Å)	10.00 – 2.10 10.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.10) 58.5 (10.00-2.10)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.148 , 0.253 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	5.5	Xtriage
Anisotropy	0.503	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 77.4	EDS
L-test for twinning ¹	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.17$	Xtriage
Estimated twinning fraction	0.127 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2742	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SUI, 13E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2432	1.32	15/3326 (0.5%)
2	B	1.92	2/48 (4.2%)	2.26	2/61 (3.3%)
All	All	0.47	2/2480 (0.1%)	1.34	17/3387 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402	HIS	CD2-NE2	-6.83	1.30	1.37
2	B	408	HIS	CD2-NE2	-6.04	1.31	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	TYR	CA-C-N	7.25	133.12	122.39
1	A	201	TYR	C-N-CA	7.25	133.12	122.39
1	A	156	PHE	CA-CB-CG	6.76	120.56	113.80
1	A	78	TYR	CA-C-N	6.05	127.75	120.13
1	A	78	TYR	C-N-CA	6.05	127.75	120.13
1	A	290	PHE	CA-C-N	6.04	126.47	120.31
1	A	290	PHE	C-N-CA	6.04	126.47	120.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	ASP	CA-CB-CG	5.82	118.42	112.60
1	A	170	PHE	CA-CB-CG	5.56	119.36	113.80
2	B	408	HIS	CB-CG-CD2	-5.40	124.18	131.20
1	A	277	ILE	CA-C-N	-5.22	113.72	121.24
1	A	277	ILE	C-N-CA	-5.22	113.72	121.24
2	B	407	ALA	N-CA-C	-5.21	96.43	111.00
1	A	258	LEU	CA-C-N	5.18	125.12	119.78
1	A	258	LEU	C-N-CA	5.18	125.12	119.78
1	A	11	ASP	CA-C-N	5.02	127.51	120.28
1	A	11	ASP	C-N-CA	5.02	127.51	120.28

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	54	SUI	Mainchain
2	B	405	13E	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2275	66	0
2	B	65	0	56	3	0
3	A	283	0	0	11	0
3	B	6	0	0	1	0
All	All	2742	0	2331	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:TYR:HB3	1:A:296:SER:HB3	1.59	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ALA:HB3	1:A:262:THR:HB	1.60	0.83
1:A:155:VAL:HG22	1:A:317:ASN:HA	1.58	0.82
1:A:265:VAL:HG13	1:A:268:ALA:HB3	1.64	0.80
1:A:190:LYS:HB2	3:A:539:HOH:O	1.86	0.75
1:A:315:VAL:O	1:A:324:LEU:HD12	1.87	0.74
1:A:41:LEU:HD12	1:A:125:LEU:HG	1.71	0.72
1:A:41:LEU:O	1:A:106:GLU:HG3	1.89	0.71
1:A:100:VAL:HG13	3:A:567:HOH:O	1.94	0.68
1:A:268:ALA:O	1:A:269:ARG:HD3	1.93	0.67
1:A:76:ILE:HG13	1:A:134:THR:HG21	1.77	0.66
1:A:230:VAL:O	1:A:233:ALA:HB3	1.94	0.66
1:A:44:PHE:HB3	1:A:57:ILE:HG22	1.80	0.64
1:A:141:LYS:HE3	3:A:540:HOH:O	1.97	0.63
1:A:191:GLN:HB3	1:A:193:PHE:HD1	1.63	0.63
1:A:201:TYR:HA	1:A:262:THR:O	2.00	0.62
1:A:191:GLN:HB3	1:A:193:PHE:CD1	2.35	0.62
1:A:265:VAL:O	1:A:268:ALA:HB3	2.01	0.60
1:A:86:GLY:HA2	1:A:109:LYS:HG3	1.84	0.58
1:A:113:SER:O	1:A:117:GLU:HG3	2.03	0.58
1:A:199:THR:OG1	1:A:264:GLY:HA3	2.08	0.54
1:A:86:GLY:CA	1:A:109:LYS:HG3	2.38	0.53
1:A:74:TRP:O	1:A:85:SER:HB2	2.09	0.53
1:A:222:THR:HG22	1:A:223:LEU:HD23	1.91	0.53
1:A:54:SUI:OD	1:A:119:SER:HB3	2.09	0.53
1:A:322:PRO:HD2	3:A:574:HOH:O	2.08	0.53
1:A:213:ILE:O	1:A:213:ILE:HG22	2.11	0.51
1:A:175:THR:HA	1:A:178:TYR:CE1	2.46	0.51
1:A:13:LEU:O	1:A:14:ASP:HB2	2.12	0.50
1:A:76:ILE:O	1:A:83:SER:HB2	2.11	0.50
1:A:252:PHE:CZ	1:A:258:LEU:HD11	2.48	0.49
1:A:273:PRO:HG2	1:A:276:TYR:CE1	2.48	0.48
1:A:130:SER:OG	1:A:145:ASP:OD1	2.30	0.48
1:A:32:LEU:HD22	1:A:43:VAL:HG12	1.95	0.48
1:A:1:SER:HB2	3:A:607:HOH:O	2.14	0.48
1:A:222:THR:HB	3:B:501:HOH:O	2.12	0.48
1:A:185:THR:HA	3:A:605:HOH:O	2.12	0.47
1:A:129:PHE:CE1	1:A:189:THR:HG22	2.49	0.47
1:A:53:VAL:HG12	1:A:54:SUI:O1	2.14	0.47
1:A:191:GLN:HE21	1:A:195:GLU:CD	2.23	0.47
1:A:4:SER:O	1:A:96:GLY:HA3	2.14	0.46
1:A:324:LEU:HD12	1:A:324:LEU:HA	1.79	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:VAL:HA	3:A:567:HOH:O	2.15	0.46
1:A:157:THR:HG22	1:A:315:VAL:HG22	1.97	0.46
1:A:6:THR:O	1:A:19:THR:HG23	2.16	0.45
1:A:10:ILE:HG22	1:A:18:ILE:HG12	1.98	0.45
1:A:273:PRO:HD2	1:A:276:TYR:CG	2.51	0.45
1:A:157:THR:OG1	1:A:169:ASN:HB2	2.18	0.44
1:A:221:THR:HG21	1:A:303:ILE:HG21	1.99	0.44
1:A:290:PHE:HE2	2:B:402:HIS:HE1	1.65	0.44
1:A:100:VAL:HG11	1:A:146:ASN:HB3	1.99	0.44
1:A:146:ASN:ND2	3:A:540:HOH:O	2.50	0.44
1:A:19:THR:HB	1:A:34:PHE:CE2	2.53	0.43
1:A:154:PRO:HG2	3:A:557:HOH:O	2.17	0.43
1:A:258:LEU:HA	1:A:259:PRO:HD2	1.67	0.43
1:A:282:ILE:CD1	1:A:288:SER:HB2	2.48	0.43
1:A:118:ASP:CB	1:A:121:ILE:HD12	2.48	0.43
1:A:20:PRO:HB3	3:A:453:HOH:O	2.19	0.43
1:A:79:GLY:HA3	2:B:405:13E:H22	2.01	0.42
1:A:39:SER:HB2	3:A:406:HOH:O	2.19	0.42
1:A:265:VAL:HG13	1:A:265:VAL:O	2.19	0.42
1:A:44:PHE:CE1	1:A:55:GLN:HG2	2.54	0.42
2:B:403:LEU:HD23	2:B:405:13E:C6	2.49	0.42
1:A:222:THR:O	1:A:307:VAL:HG13	2.19	0.42
1:A:21:VAL:HG21	1:A:32:LEU:HD11	2.03	0.41
1:A:268:ALA:C	1:A:269:ARG:HD3	2.46	0.41
1:A:225:TYR:HB3	1:A:296:SER:CB	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/329 (99%)	317 (97%)	8 (2%)	1 (0%)	36	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	4/8 (50%)	2 (50%)	1 (25%)	1 (25%)	0	0
All	All	330/337 (98%)	319 (97%)	9 (3%)	2 (1%)	21	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	402	HIS
1	A	194	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/262 (100%)	239 (91%)	23 (9%)	9	7
2	B	5/6 (83%)	3 (60%)	2 (40%)	0	0
All	All	267/268 (100%)	242 (91%)	25 (9%)	8	6

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	A	12	SER
1	A	17	TYR
1	A	21	VAL
1	A	40	ASP
1	A	87	ASP
1	A	94	SER
1	A	118	ASP
1	A	120	THR
1	A	124	LEU
1	A	131	THR
1	A	137	PRO
1	A	138	THR
1	A	146	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	SER
1	A	153	SER
1	A	179	THR
1	A	181	SER
1	A	232	SER
1	A	246	SER
1	A	282	ILE
1	A	284	THR
1	A	301	ILE
2	B	402	HIS
2	B	408	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN
1	A	146	ASN
1	A	191	GLN
2	B	402	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SUI	A	54	1	10,11,12	1.27	1 (10%)	12,15,17	2.14	3 (25%)
2	13E	B	405	2	17,18,19	0.73	0	15,21,23	1.19	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SUI	A	54	1	-	1/2/19/20	0/1/1/1
2	13E	B	405	2	-	5/14/15/17	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54	SUI	CB-CG	2.33	1.54	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	SUI	C1-N2-CG	5.32	116.18	113.07
1	A	54	SUI	OD-CG-N2	2.99	127.14	123.94
2	B	405	13E	C3-C2-CA	2.56	118.16	113.23
1	A	54	SUI	C2A-N2-C1	-2.41	120.45	123.07
2	B	405	13E	O-C-C10	-2.03	119.55	124.77

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	405	13E	C10-C11-C12-SD
2	B	405	13E	CA-C2-C3-C5
2	B	405	13E	CA-C2-C3-C4
2	B	405	13E	CA-C9-N2-C10
1	A	54	SUI	C-C2A-N2-CG
2	B	405	13E	C11-C12-SD-C13

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	54	SUI	2	0
2	B	405	13E	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/329 (99%)	-0.78	0 100 100	0, 8, 18, 34	0
2	B	6/8 (75%)	0.13	0 100 100	9, 20, 53, 75	0
All	All	334/337 (99%)	-0.76	0 100 100	0, 8, 19, 75	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	13E	B	405	18/19	0.96	0.05	0,7,20,33	0
1	SUI	A	54	11/12	0.98	0.04	0,9,13,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.