



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:29 PM UTC

PDB ID : 4UR7 / pdb\_00004ur7  
Title : Crystal structure of keto-deoxy-D-galactarate dehydratase complexed with pyruvate  
Authors : Taberman, H.; Parkkinen, T.; Hakulinen, N.; Rouvinen, J.  
Deposited on : 2014-06-26  
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	 89% 8% .
1	B	312	 2% 92% 8%
1	C	312	 2% 91% 8% .
1	D	312	 2% 88% 8% .

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 10972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KETO-DEOXY-D-GALACTARATE DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	303	2316	1482	390	431	13	0	5	0
1	B	310	2374	1521	400	440	13	0	3	0
1	C	307	2348	1504	396	435	13	0	4	0
1	D	301	2303	1474	386	430	13	0	5	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ASP	ASN	cloning artifact	UNP Q8UB77
A	166	KPI	LYS	microheterogeneity	UNP Q8UB77
A	304	TRP	-	expression tag	UNP Q8UB77
A	305	SER	-	expression tag	UNP Q8UB77
A	306	HIS	-	expression tag	UNP Q8UB77
A	307	PRO	-	expression tag	UNP Q8UB77
A	308	GLN	-	expression tag	UNP Q8UB77
A	309	PHE	-	expression tag	UNP Q8UB77
A	310	GLU	-	expression tag	UNP Q8UB77
A	311	LYS	-	expression tag	UNP Q8UB77
B	2	ASP	ASN	cloning artifact	UNP Q8UB77
B	166	KPI	LYS	microheterogeneity	UNP Q8UB77
B	304	TRP	-	expression tag	UNP Q8UB77
B	305	SER	-	expression tag	UNP Q8UB77
B	306	HIS	-	expression tag	UNP Q8UB77
B	307	PRO	-	expression tag	UNP Q8UB77
B	308	GLN	-	expression tag	UNP Q8UB77
B	309	PHE	-	expression tag	UNP Q8UB77
B	310	GLU	-	expression tag	UNP Q8UB77
B	311	LYS	-	expression tag	UNP Q8UB77
C	2	ASP	ASN	cloning artifact	UNP Q8UB77

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	166	KPI	LYS	microheterogeneity	UNP Q8UB77
C	304	TRP	-	expression tag	UNP Q8UB77
C	305	SER	-	expression tag	UNP Q8UB77
C	306	HIS	-	expression tag	UNP Q8UB77
C	307	PRO	-	expression tag	UNP Q8UB77
C	308	GLN	-	expression tag	UNP Q8UB77
C	309	PHE	-	expression tag	UNP Q8UB77
C	310	GLU	-	expression tag	UNP Q8UB77
C	311	LYS	-	expression tag	UNP Q8UB77
D	2	ASP	ASN	cloning artifact	UNP Q8UB77
D	166	KPI	LYS	microheterogeneity	UNP Q8UB77
D	304	TRP	-	expression tag	UNP Q8UB77
D	305	SER	-	expression tag	UNP Q8UB77
D	306	HIS	-	expression tag	UNP Q8UB77
D	307	PRO	-	expression tag	UNP Q8UB77
D	308	GLN	-	expression tag	UNP Q8UB77
D	309	PHE	-	expression tag	UNP Q8UB77
D	310	GLU	-	expression tag	UNP Q8UB77
D	311	LYS	-	expression tag	UNP Q8UB77

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	C	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	468	Total	O	0	0
			468	468		
4	B	406	Total	O	0	0
			406	406		
4	C	325	Total	O	0	0
			325	325		
4	D	360	Total	O	0	0
			360	360		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: KETO-DEOXY-D-GALACTARATE DEHYDRATASE




- Molecule 1: KETO-DEOXY-D-GALACTARATE DEHYDRATASE



- Molecule 1: KETO-DEOXY-D-GALACTARATE DEHYDRATASE



- Molecule 1: KETO-DEOXY-D-GALACTARATE DEHYDRATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.80Å 120.14Å 74.07Å 90.00° 112.01° 90.00°	Depositor
Resolution (Å)	48.09 – 1.50 48.09 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.09-1.50) 97.8 (48.09-1.50)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.163 , 0.182 (Not available) , 0.188	Depositor DCC
$R_{free}$ test set	10750 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.3	Xtrriage
Anisotropy	0.536	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 33.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10972	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, GOL, KPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/2360	0.73	0/3193
1	B	0.46	0/2417	0.73	0/3272
1	C	0.38	0/2393	0.72	0/3240
1	D	0.40	0/2347	0.71	1/3176 (0.0%)
All	All	0.43	0/9517	0.72	1/12881 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	13	GLY	N-CA-C	5.05	117.02	110.45

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2316	0	2316	21	0
1	B	2374	0	2357	16	0
1	C	2348	0	2339	13	0
1	D	2303	0	2298	19	0
2	A	24	0	32	0	0
2	B	18	0	24	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	6	0	8	0	0
2	D	12	0	16	0	0
3	A	3	0	1	0	0
3	B	3	0	1	0	0
3	C	3	0	1	0	0
3	D	3	0	1	0	0
4	A	468	0	0	7	1
4	B	406	0	0	6	1
4	C	325	0	0	1	0
4	D	360	0	0	5	1
All	All	10972	0	9394	63	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:123:TYR:OH	4:D:2201:HOH:O	2.05	0.75
1:D:233[B]:GLU:OE1	4:D:2291:HOH:O	2.07	0.71
1:A:289:ILE:HD11	4:A:2421:HOH:O	1.91	0.70
1:A:282:LYS:NZ	4:A:2437:HOH:O	2.25	0.68
1:C:233:GLU:OE1	4:C:2247:HOH:O	2.11	0.68

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:2125:HOH:O	4:D:2127:HOH:O[2_658]	1.95	0.25
4:A:2148:HOH:O	4:B:2124:HOH:O[2_557]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/312 (97%)	300 (99%)	4 (1%)	0	100	100
1	B	309/312 (99%)	305 (99%)	4 (1%)	0	100	100
1	C	307/312 (98%)	302 (98%)	5 (2%)	0	100	100
1	D	302/312 (97%)	296 (98%)	6 (2%)	0	100	100
All	All	1222/1248 (98%)	1203 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/238 (98%)	233 (100%)	1 (0%)	84	71
1	B	239/238 (100%)	236 (99%)	3 (1%)	61	35
1	C	237/238 (100%)	235 (99%)	2 (1%)	73	54
1	D	233/238 (98%)	232 (100%)	1 (0%)	84	71
All	All	943/952 (99%)	936 (99%)	7 (1%)	73	57

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	310	GLU
1	C	44	LYS
1	D	138	MET
1	C	138	MET
1	B	138	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	306	HIS

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	C	306	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KPI	C	166[A]	-	11,13,14	3.03	2 (18%)	9,15,17	3.43	4 (44%)
1	KPI	B	166[A]	-	11,13,14	2.94	2 (18%)	9,15,17	2.96	5 (55%)
1	KPI	A	166[A]	-	11,13,14	3.00	2 (18%)	9,15,17	2.35	6 (66%)
1	KPI	D	166[A]	-	11,13,14	3.16	3 (27%)	9,15,17	2.05	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KPI	C	166[A]	-	-	3/13/14/16	-
1	KPI	B	166[A]	-	-	2/13/14/16	-
1	KPI	A	166[A]	-	-	2/13/14/16	-
1	KPI	D	166[A]	-	-	2/13/14/16	-

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	166[A]	KPI	CX2-CX1	-9.40	1.38	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	166[A]	KPI	CX2-CX1	-9.11	1.39	1.49
1	A	166[A]	KPI	CX2-CX1	-9.10	1.39	1.49
1	B	166[A]	KPI	CX2-CX1	-8.87	1.39	1.49
1	D	166[A]	KPI	O2-CX2	3.52	1.31	1.22

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	166[A]	KPI	CE-NZ-CX1	7.23	141.46	121.58
1	B	166[A]	KPI	CE-NZ-CX1	5.86	137.67	121.58
1	C	166[A]	KPI	CX2-CX1-NZ	5.57	127.99	114.88
1	B	166[A]	KPI	CX2-CX1-NZ	4.50	125.47	114.88
1	D	166[A]	KPI	CE-NZ-CX1	4.17	133.03	121.58

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	166[A]	KPI	NZ-CX1-CX2-O1
1	D	166[A]	KPI	NZ-CX1-CX2-O2
1	A	166[A]	KPI	NZ-CX1-CX2-O2
1	B	166[A]	KPI	NZ-CX1-CX2-O1
1	B	166[A]	KPI	NZ-CX1-CX2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FMT	C	1310	-	2,2,2	0.71	0	1,1,1	0.20	0
2	GOL	A	1309	-	5,5,5	0.37	0	5,5,5	0.35	0
3	FMT	B	1315	-	2,2,2	0.68	0	1,1,1	0.28	0
2	GOL	C	1309	-	5,5,5	0.38	0	5,5,5	0.21	0
2	GOL	B	1313	-	5,5,5	0.37	0	5,5,5	0.14	0
3	FMT	A	1306	-	2,2,2	0.71	0	1,1,1	0.19	0
2	GOL	A	1305	-	5,5,5	0.30	0	5,5,5	0.15	0
2	GOL	A	1308	-	5,5,5	0.36	0	5,5,5	0.45	0
2	GOL	D	1303	-	5,5,5	0.40	0	5,5,5	0.13	0
2	GOL	B	1314	-	5,5,5	0.28	0	5,5,5	0.39	0
2	GOL	B	1312	-	5,5,5	0.32	0	5,5,5	0.38	0
2	GOL	D	1305	-	5,5,5	0.42	0	5,5,5	0.64	0
3	FMT	D	1304	-	2,2,2	0.73	0	1,1,1	0.23	0
2	GOL	A	1307	-	5,5,5	0.27	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1309	-	-	0/4/4/4	-
2	GOL	C	1309	-	-	0/4/4/4	-
2	GOL	B	1313	-	-	0/4/4/4	-
2	GOL	A	1305	-	-	0/4/4/4	-
2	GOL	A	1308	-	-	2/4/4/4	-
2	GOL	D	1303	-	-	0/4/4/4	-
2	GOL	B	1314	-	-	0/4/4/4	-
2	GOL	B	1312	-	-	0/4/4/4	-
2	GOL	D	1305	-	-	2/4/4/4	-
2	GOL	A	1307	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1308	GOL	C1-C2-C3-O3
2	D	1305	GOL	O1-C1-C2-C3
2	A	1308	GOL	O2-C2-C3-O3
2	D	1305	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	302/312 (96%)	-0.31	0 <b>100</b> <b>100</b>	10, 17, 26, 42	4 (1%)
1	B	309/312 (99%)	-0.16	7 (2%) 61 65	10, 18, 28, 55	2 (0%)
1	C	306/312 (98%)	0.45	5 (1%) 70 74	15, 24, 36, 53	3 (0%)
1	D	300/312 (96%)	0.19	5 (1%) 69 72	14, 22, 34, 53	4 (1%)
All	All	1217/1248 (97%)	0.04	17 (1%) 73 77	10, 21, 33, 55	13 (1%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	309	PHE	4.5
1	C	306	HIS	3.4
1	C	24	ALA	3.4
1	B	78	GLU	3.2
1	B	24	ALA	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KPI	B	166[A]	14/15	0.92	0.10	14,16,22,23	14
1	KPI	A	166[A]	14/15	0.93	0.09	13,14,17,21	14
1	KPI	C	166[A]	14/15	0.93	0.10	17,19,27,28	14
1	KPI	D	166[A]	14/15	0.94	0.09	15,18,24,27	14

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	A	1309	6/6	0.77	0.18	28,40,45,47	0
3	FMT	D	1304	3/3	0.82	0.14	33,33,37,45	0
3	FMT	A	1306	3/3	0.83	0.15	32,32,32,38	0
3	FMT	C	1310	3/3	0.87	0.13	34,34,40,45	0
2	GOL	D	1305	6/6	0.89	0.13	25,31,37,40	0
2	GOL	B	1314	6/6	0.89	0.11	24,26,32,32	0
2	GOL	A	1307	6/6	0.91	0.11	22,24,30,34	0
3	FMT	B	1315	3/3	0.93	0.11	28,28,31,41	0
2	GOL	A	1308	6/6	0.94	0.09	17,20,22,28	0
2	GOL	C	1309	6/6	0.96	0.07	20,23,26,28	0
2	GOL	D	1303	6/6	0.97	0.07	17,19,22,23	0
2	GOL	B	1313	6/6	0.98	0.05	16,17,17,20	0
2	GOL	A	1305	6/6	0.98	0.06	15,16,16,19	0
2	GOL	B	1312	6/6	0.98	0.05	15,16,18,18	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.