



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:45 PM UTC

PDB ID : 4USV / pdb_00004usv
Title : Crystal structure of human soluble Adenylyl Cyclase with pyrophosphate resulting from soaking with ATP and Calcium
Authors : Kleinboelting, S.; Steegborn, C.
Deposited on : 2014-07-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	5% 77% 16% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	1471	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3843 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADENYLATE CYCLASE TYPE 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	458	3629	2351	586	659	33	0	5	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	470	HIS	-	expression tag	UNP Q96PN6
A	471	HIS	-	expression tag	UNP Q96PN6
A	472	HIS	-	expression tag	UNP Q96PN6
A	473	HIS	-	expression tag	UNP Q96PN6
A	474	HIS	-	expression tag	UNP Q96PN6
A	475	HIS	-	expression tag	UNP Q96PN6

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			9	7	2		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	185	Total 185	O 185	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADENYLATE CYCLASE TYPE 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	100.87Å 100.87Å 97.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	87.36 – 2.00 87.36 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (87.36-2.00) 100.0 (87.36-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.160 , 0.209 0.171 , 0.216	Depositor DCC
R_{free} test set	1929 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3843	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CME, EDO, GOL, CA, CL, POP, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	13/3720 (0.3%)	1.21	14/5042 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	PRO	CA-C	-7.63	1.47	1.52
1	A	285	TYR	C-O	-6.57	1.16	1.24
1	A	236	PHE	C-O	-6.11	1.15	1.23
1	A	236	PHE	N-CA	-6.01	1.38	1.46
1	A	220	PRO	C-O	-5.93	1.19	1.24
1	A	32	GLU	CA-C	5.77	1.59	1.52
1	A	106	ARG	C-O	-5.61	1.17	1.24
1	A	173	ASP	N-CA	5.54	1.53	1.46
1	A	369	ILE	N-CA	-5.45	1.40	1.46
1	A	179	GLN	CA-C	5.28	1.59	1.52
1	A	405	THR	CA-C	5.28	1.59	1.52
1	A	287	SER	CA-C	-5.04	1.46	1.52
1	A	83	VAL	C-O	-5.00	1.18	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	377	HIS	N-CA-C	9.16	120.87	111.07
1	A	220	PRO	O-C-N	8.02	124.92	121.15
1	A	355	VAL	CA-C-N	-7.93	112.67	120.52
1	A	355	VAL	C-N-CA	-7.93	112.67	120.52
1	A	222	PRO	CA-C-N	-7.31	110.96	122.65
1	A	222	PRO	C-N-CA	-7.31	110.96	122.65
1	A	48	ILE	CB-CA-C	-6.12	103.48	111.25
1	A	327	ILE	N-CA-C	5.99	116.73	110.62
1	A	166	LEU	CB-CA-C	-5.95	97.66	110.45
1	A	220	PRO	CA-C-O	-5.80	116.14	120.73
1	A	126	ILE	CB-CA-C	-5.61	104.67	112.02
1	A	201	MET	CB-CA-C	-5.26	100.24	109.07
1	A	246	LYS	N-CA-C	5.09	118.98	112.87
1	A	141	ILE	N-CA-C	5.05	115.18	108.11

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	355	VAL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3629	0	3580	65	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	8	0	12	4	0
5	A	4	0	3	4	0
6	A	9	0	0	0	0
7	A	6	0	8	3	0
8	A	185	0	0	12	0
All	All	3843	0	3603	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:GLU:HG3	4:A:1469:EDO:O1	1.70	0.91
1:A:409:GLN:HG2	8:A:2081:HOH:O	1.81	0.80
1:A:166:LEU:CD2	1:A:336:PHE:HA	2.16	0.76
1:A:9:GLN:HB2	8:A:2005:HOH:O	1.84	0.76
1:A:350:PHE:HB3	1:A:351:PRO:HD2	1.69	0.72
1:A:201:MET:HE1	1:A:224:PHE:CE1	2.27	0.69
1:A:270[B]:MET:HE1	1:A:404:TYR:CE2	2.27	0.69
1:A:28:HIS:HE1	8:A:2023:HOH:O	1.76	0.67
1:A:339:ASP:CG	1:A:419:MET:HG2	2.20	0.67
1:A:58:PHE:CD2	1:A:71[A]:LEU:HD13	2.30	0.67
1:A:245:HIS:HE1	8:A:2112:HOH:O	1.77	0.66
1:A:58:PHE:CE2	1:A:71[A]:LEU:HD13	2.35	0.62
1:A:270[B]:MET:HE1	1:A:404:TYR:CD2	2.33	0.62
1:A:324:VAL:HG12	1:A:365:CYS:SG	2.41	0.61
1:A:166:LEU:HD23	1:A:336:PHE:HA	1.81	0.61
1:A:356:PRO:HA	8:A:2155:HOH:O	2.02	0.59
1:A:54:MET:HB3	1:A:58:PHE:HE2	1.69	0.58
1:A:350:PHE:CB	1:A:351:PRO:HD2	2.34	0.58
1:A:160:GLU:O	1:A:267:LYS:HE3	2.05	0.56
1:A:335:VAL:H	5:A:1471:ACT:H2	1.70	0.56
1:A:331:GLN:HG3	1:A:353:GLU:OE1	2.08	0.54
1:A:350:PHE:O	1:A:351:PRO:C	2.51	0.52
1:A:75:LEU:HD11	1:A:79:ILE:CD1	2.39	0.51
1:A:183:GLN:O	1:A:184:MET:C	2.51	0.51
1:A:9:GLN:N	8:A:2005:HOH:O	2.44	0.51
1:A:28:HIS:CE1	8:A:2023:HOH:O	2.59	0.50
1:A:95:LYS:HE2	5:A:1471:ACT:H1	1.94	0.49
1:A:83:VAL:HG13	1:A:93:ILE:CD1	2.42	0.49
1:A:75:LEU:HD11	1:A:79:ILE:HD11	1.95	0.49
1:A:51:PHE:HB2	1:A:99:ASP:OD2	2.13	0.48
1:A:8:PHE:CE2	1:A:10[B]:ASP:HB2	2.48	0.48
1:A:75:LEU:HG	1:A:79:ILE:HD12	1.95	0.47
1:A:251:LEU:HD22	4:A:1470:EDO:H22	1.94	0.47
1:A:54:MET:HB3	1:A:58:PHE:CE2	2.48	0.47
1:A:345:LEU:C	1:A:345:LEU:HD23	2.40	0.47
1:A:373:CYS:HA	1:A:376:VAL:HG13	1.96	0.47
1:A:357:ASP:HA	8:A:2156:HOH:O	2.15	0.47
1:A:54:MET:O	1:A:58:PHE:HD2	1.99	0.46
1:A:54:MET:O	1:A:58:PHE:CD2	2.69	0.46
1:A:27:GLY:O	1:A:28:HIS:C	2.53	0.45
1:A:48:ILE:HD11	1:A:82:ILE:CD1	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:VAL:HG11	1:A:115:ILE:HG13	1.97	0.45
1:A:350:PHE:HB3	1:A:351:PRO:CD	2.37	0.45
1:A:31:PRO:HB3	7:A:1473:GOL:H2	1.99	0.44
7:A:1473:GOL:C3	8:A:2124:HOH:O	2.64	0.44
1:A:36:MET:HE3	1:A:308:GLU:HA	2.00	0.44
1:A:158:GLY:HA2	1:A:318:TYR:CE2	2.52	0.44
7:A:1473:GOL:H31	8:A:2124:HOH:O	2.17	0.44
1:A:95:LYS:CE	5:A:1471:ACT:H1	2.48	0.43
1:A:251:LEU:HD21	4:A:1470:EDO:H11	1.99	0.43
1:A:270[B]:MET:CG	8:A:2128:HOH:O	2.66	0.43
1:A:288:GLU:HG3	4:A:1469:EDO:HO1	1.76	0.43
1:A:203:GLU:HG2	1:A:217:PHE:CZ	2.54	0.43
1:A:324:VAL:CG1	1:A:365:CYS:SG	3.07	0.43
1:A:456:SER:C	1:A:457:GLY:O	2.61	0.42
1:A:417:MET:HE1	1:A:427:CYS:HA	2.01	0.42
1:A:205:GLU:HB3	1:A:216:ASN:HD21	1.85	0.41
1:A:297:VAL:O	1:A:343:SER:HA	2.21	0.41
1:A:334:LYS:HA	5:A:1471:ACT:H2	2.02	0.41
1:A:28:HIS:HB2	8:A:2024:HOH:O	2.20	0.41
1:A:58:PHE:CE2	1:A:71[A]:LEU:CD1	3.04	0.41
1:A:260:GLU:HA	1:A:263:MET:HE3	2.03	0.41
1:A:270[A]:MET:HE3	1:A:270[A]:MET:HB3	1.86	0.41
1:A:368:ASP:OD1	1:A:368:ASP:C	2.64	0.41
1:A:160:GLU:O	1:A:267:LYS:CE	2.69	0.40
1:A:166:LEU:HD21	1:A:336:PHE:HA	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/475 (96%)	449 (98%)	8 (2%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	457	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/418 (95%)	377 (95%)	19 (5%)	23	21

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	54	MET
1	A	56	GLU
1	A	108	GLU
1	A	140	ASP
1	A	201	MET
1	A	222	PRO
1	A	257	PRO
1	A	270[A]	MET
1	A	270[B]	MET
1	A	312	PRO
1	A	324	VAL
1	A	334	LYS
1	A	350	PHE
1	A	355	VAL
1	A	376	VAL
1	A	402	HIS
1	A	418	MET
1	A	466	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	78	HIS
1	A	111	GLN
1	A	114	ASN
1	A	195	GLN
1	A	216	ASN
1	A	238	HIS
1	A	276	GLN
1	A	281	GLN
1	A	331	GLN
1	A	433	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	253	1	8,9,10	0.84	0	6,9,11	2.24	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	253	1	-	0/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	253	CME	OH-CZ-CE	-4.71	92.45	110.82
1	A	253	CME	CB-SG-SD	-2.49	97.41	103.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	POP	A	1472	3	6,8,8	0.70	0	12,13,13	1.44	2 (16%)
5	ACT	A	1471	-	3,3,3	1.71	1 (33%)	3,3,3	0.56	0
7	GOL	A	1473	-	5,5,5	1.05	0	5,5,5	0.77	0
4	EDO	A	1469	-	3,3,3	0.45	0	2,2,2	0.54	0
4	EDO	A	1470	-	3,3,3	0.72	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	1473	-	-	4/4/4/4	-
6	POP	A	1472	3	-	0/6/6/6	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1469	-	-	1/1/1/1	-
4	EDO	A	1470	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1471	ACT	CH3-C	-2.50	1.39	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1472	POP	O3-P1-O2	3.24	119.96	107.80
6	A	1472	POP	O-P2-O4	-2.12	99.86	111.04

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1473	GOL	C1-C2-C3-O3
7	A	1473	GOL	O2-C2-C3-O3
7	A	1473	GOL	O1-C1-C2-C3
7	A	1473	GOL	O1-C1-C2-O2
4	A	1469	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1471	ACT	4	0
7	A	1473	GOL	3	0
4	A	1469	EDO	2	0
4	A	1470	EDO	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/475 (96%)	0.04	23 (5%) 34 33	18, 40, 77, 98	6 (1%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	LEU	3.9
1	A	350	PHE	3.6
1	A	351	PRO	3.6
1	A	466	THR	3.4
1	A	355	VAL	3.2
1	A	339	ASP	3.0
1	A	356	PRO	2.9
1	A	352	GLY	2.8
1	A	8	PHE	2.7
1	A	285	TYR	2.7
1	A	141	ILE	2.6
1	A	377	HIS	2.6
1	A	449	VAL	2.6
1	A	130	PHE	2.5
1	A	340	LYS	2.4
1	A	349	GLY	2.3
1	A	51	PHE	2.2
1	A	338	PHE	2.2
1	A	436	ASN	2.2
1	A	54	MET	2.1
1	A	353	GLU	2.1
1	A	456	SER	2.1
1	A	185	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	253	10/11	0.97	0.08	26,30,41,44	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	POP	A	1472	9/9	0.75	0.09	84,87,96,99	0
4	EDO	A	1469	4/4	0.84	0.15	59,71,77,77	0
7	GOL	A	1473	6/6	0.86	0.14	50,63,68,74	0
4	EDO	A	1470	4/4	0.89	0.21	48,50,57,59	0
3	CA	A	1468	1/1	0.90	0.11	81,81,81,81	0
5	ACT	A	1471	4/4	0.90	0.21	20,31,58,74	0
2	CL	A	1467	1/1	0.99	0.03	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.