



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 10:20 AM UTC

PDB ID : 8USX / pdb_00008usx
EMDB ID : EMD-42522
Title : Glycine-bound GluN1a-3A NMDA receptor
Authors : Michalski, K.; Furukawa, H.
Deposited on : 2023-10-30
Resolution : 4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	847	6% 76% 11% 13%
1	C	847	12% 76% 11% 13%
2	B	939	35% 70% 9% 21%
2	D	939	40% 67% 12% 21%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 17926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor ionotropic, NMDA 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	741	Total	C	N	O	S	0	0
			4477	2756	834	871	16		
1	C	738	Total	C	N	O	S	0	0
			4495	2765	843	870	17		

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q05586
A	2	SER	-	expression tag	UNP Q05586
A	3	THR	-	expression tag	UNP Q05586
A	4	MET	-	expression tag	UNP Q05586
A	5	HIS	-	expression tag	UNP Q05586
A	6	LEU	-	expression tag	UNP Q05586
A	7	LEU	-	expression tag	UNP Q05586
A	8	THR	-	expression tag	UNP Q05586
A	9	PHE	-	expression tag	UNP Q05586
A	10	ALA	-	expression tag	UNP Q05586
A	11	LEU	-	expression tag	UNP Q05586
A	12	LEU	-	expression tag	UNP Q05586
A	13	PHE	-	expression tag	UNP Q05586
A	14	SER	-	expression tag	UNP Q05586
A	15	CYS	-	expression tag	UNP Q05586
A	16	SER	-	expression tag	UNP Q05586
A	17	PHE	-	expression tag	UNP Q05586
A	18	ALA	-	expression tag	UNP Q05586
A	19	ARG	-	expression tag	UNP Q05586
A	20	ALA	-	expression tag	UNP Q05586
A	21	ALA	-	expression tag	UNP Q05586
A	22	SER	-	expression tag	UNP Q05586
A	415	MET	LEU	conflict	UNP Q05586
A	810	CYS	PHE	conflict	UNP Q05586
A	844	ASN	-	expression tag	UNP Q05586
A	845	GLY	-	expression tag	UNP Q05586

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	846	ALA	-	expression tag	UNP Q05586
A	847	GLN	-	expression tag	UNP Q05586
C	1	MET	-	initiating methionine	UNP Q05586
C	2	SER	-	expression tag	UNP Q05586
C	3	THR	-	expression tag	UNP Q05586
C	4	MET	-	expression tag	UNP Q05586
C	5	HIS	-	expression tag	UNP Q05586
C	6	LEU	-	expression tag	UNP Q05586
C	7	LEU	-	expression tag	UNP Q05586
C	8	THR	-	expression tag	UNP Q05586
C	9	PHE	-	expression tag	UNP Q05586
C	10	ALA	-	expression tag	UNP Q05586
C	11	LEU	-	expression tag	UNP Q05586
C	12	LEU	-	expression tag	UNP Q05586
C	13	PHE	-	expression tag	UNP Q05586
C	14	SER	-	expression tag	UNP Q05586
C	15	CYS	-	expression tag	UNP Q05586
C	16	SER	-	expression tag	UNP Q05586
C	17	PHE	-	expression tag	UNP Q05586
C	18	ALA	-	expression tag	UNP Q05586
C	19	ARG	-	expression tag	UNP Q05586
C	20	ALA	-	expression tag	UNP Q05586
C	21	ALA	-	expression tag	UNP Q05586
C	22	SER	-	expression tag	UNP Q05586
C	415	MET	LEU	conflict	UNP Q05586
C	810	CYS	PHE	conflict	UNP Q05586
C	844	ASN	-	expression tag	UNP Q05586
C	845	GLY	-	expression tag	UNP Q05586
C	846	ALA	-	expression tag	UNP Q05586
C	847	GLN	-	expression tag	UNP Q05586


- Molecule 2 is a protein called Glutamate receptor ionotropic, NMDA 3A.


Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	745	Total	C	N	O	S	0	0
			4500	2779	825	882	14		
2	D	743	Total	C	N	O	S	0	0
			4454	2748	823	870	13		

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	676	CYS	THR	conflict	UNP Q8TCU5
B	968	THR	-	expression tag	UNP Q8TCU5
B	969	GLU	-	expression tag	UNP Q8TCU5
B	970	THR	-	expression tag	UNP Q8TCU5
B	971	SER	-	expression tag	UNP Q8TCU5
B	972	GLN	-	expression tag	UNP Q8TCU5
B	973	VAL	-	expression tag	UNP Q8TCU5
B	974	ALA	-	expression tag	UNP Q8TCU5
B	975	PRO	-	expression tag	UNP Q8TCU5
B	976	ALA	-	expression tag	UNP Q8TCU5
D	676	CYS	THR	conflict	UNP Q8TCU5
D	968	THR	-	expression tag	UNP Q8TCU5
D	969	GLU	-	expression tag	UNP Q8TCU5
D	970	THR	-	expression tag	UNP Q8TCU5
D	971	SER	-	expression tag	UNP Q8TCU5
D	972	GLN	-	expression tag	UNP Q8TCU5
D	973	VAL	-	expression tag	UNP Q8TCU5
D	974	ALA	-	expression tag	UNP Q8TCU5
D	975	PRO	-	expression tag	UNP Q8TCU5
D	976	ALA	-	expression tag	UNP Q8TCU5

- Molecule 2: Glutamate receptor ionotropic, NMDA 3A





• Molecule 2: Glutamate receptor ionotropic, NMDA 3A



M900	D901	C913	GLY	LYS	ARG	SER	PHE	ALA	VAL	THR	GLU	THR	LEU	GLN	M926	S945	V955	TYR	ARG	LEU	LEU	LEU	PRO	ARG	ILE	LYS	ASN	LYS	SER	THR	GLU	THR	SER	GLN	VAL	ALA	ALA														
P788	R793	F794	E799	E803	P811	E812	M813	H814	E815	Y816	M817	R818	R819	Y820	N821	T825	F826	D827	G828	V829	L832	D835	P836	E837	F842	K846	L849	E852	V853	S854	I855	K860	L861	I870	E871	G872	Y873	G874	I875	L883	N886	S896	H897								
THR	ALA	VAL	PHE	LEU	THR	LEU	TYR	GLU	TRP	LYS	SER	PRO	PHE	GLY	LEU	THR	PRO	LYS	GLY	ARG	THR	VAL	ALA	ILE	CYS	TYR	ALA	LEU	PHE	GLY	ARG	THR	VAL	ALA	ILE	LYS	PRO	PRO	LYS	CYS	W740	V769	L777	H786	H787						
D602	G606	H612	W613	T614	G615	L616	T624	M627	A628	V629	T630	S631	F632	S633	I634	N635	R638	F644	S652	I655	L656	T659	R660	ASP	THR	ALA	ALA	PRO	ILE	GLY	PHE	PHE	MET	TRP	PRO	PRO	HIS	TRP	CYS	MET	LYS	TRP	LEU	GLY	ILE	PHE	VAL	ALA	LEU	HIS	ILE

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	225900	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.139	Depositor
Minimum map value	-1.308	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.037	Depositor
Recommended contour level	0.149	Depositor
Map size ($\text{\AA}$)	342.4, 342.4, 342.4	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.856, 0.856, 0.856	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/4526	0.34	0/6214
1	C	0.14	0/4542	0.38	1/6223 (0.0%)
2	B	0.15	0/4556	0.39	2/6260 (0.0%)
2	D	0.13	0/4508	0.38	0/6196
All	All	0.14	0/18132	0.37	3/24893 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	447	PRO	N-CA-CB	7.30	110.91	103.25
2	B	793	ARG	CA-C-N	6.15	133.29	121.54
2	B	793	ARG	C-N-CA	6.15	133.29	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	523	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4477	0	3168	68	0
1	C	4495	0	3224	69	0
2	B	4500	0	3129	60	0
2	D	4454	0	3072	92	0
All	All	17926	0	12593	280	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 280 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:622:ARG:HB3	2:B:622:ARG:NH1	1.81	0.96
2:B:877:LEU:HD12	2:B:878:PRO:HD2	1.54	0.87
1:C:533:PHE:HB3	1:C:776:ILE:HD11	1.67	0.76
2:B:529:ARG:O	2:B:575:CYS:HB2	1.87	0.75
2:D:629:VAL:HG12	2:D:875:ILE:HD12	1.68	0.74

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	735/847 (87%)	679 (92%)	52 (7%)	4 (0%)	24 61
1	C	732/847 (86%)	669 (91%)	58 (8%)	5 (1%)	18 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	735/939 (78%)	690 (94%)	42 (6%)	3 (0%)	30	65
2	D	733/939 (78%)	680 (93%)	49 (7%)	4 (0%)	24	61
All	All	2935/3572 (82%)	2718 (93%)	201 (7%)	16 (0%)	26	61

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	660	PRO
1	A	698	GLU
1	A	724	ASN
2	B	636	THR
1	C	444	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/728 (33%)	237 (100%)	0	100	100
1	C	243/728 (33%)	243 (100%)	0	100	100
2	B	246/816 (30%)	244 (99%)	2 (1%)	73	77
2	D	236/816 (29%)	236 (100%)	0	100	100
All	All	962/3088 (31%)	960 (100%)	2 (0%)	85	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	622	ARG
2	B	830	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	796	GLN
2	D	549	ASN
2	D	790	GLN
2	B	880	ASN
2	B	904	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-42522. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 187




Y Index: 201




Z Index: 208


6.3.2 Raw map



X Index: 201



Y Index: 200




Z Index: 210


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.149. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 404 nm³; this corresponds to an approximate mass of 365 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.58	7.06	4.66

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.58 differs from the reported value 4.1 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-42522 and PDB model 8USX. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.149 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.149).

9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.149) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7440	0.2720
A	0.9210	0.2930
B	0.6350	0.2610
C	0.8600	0.3130
D	0.5610	0.2210

