



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:05 PM UTC

PDB ID : 1UTB / pdb_00001utb
Title : DntR from Burkholderia sp. strain DNT
Authors : Smirnova, I.A.; Dian, C.; Leonard, G.A.; McSweeney, S.; Birse, D.; Brzezinski, P.
Deposited on : 2003-12-05
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	3% 51% 14% • 32%
2	B	315	6% 49% 20% • 28%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1304	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSR-TYPE REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1683	1081	300	291	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	SER	THR	engineered mutation	UNP Q7WT50

- Molecule 2 is a protein called LYSR-TYPE REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	0	0
			1764	1129	313	311	11			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	139	SER	LYS	conflict	UNP Q7WT50
B	192	SER	THR	engineered mutation	UNP Q7WT50
B	247	GLU	GLN	conflict	UNP Q7WT50

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total 59	O 59	0	1
5	B	54	Total 54	O 54	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	107.15Å 107.15Å 296.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	91.29 – 2.59 91.29 – 2.59	Depositor EDS
% Data completeness (in resolution range)	97.2 (91.29-2.59) 97.3 (91.29-2.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.215 , 0.261 0.220 , 0.263	Depositor DCC
R_{free} test set	1630 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3598	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	2/1729 (0.1%)	0.95	1/2346 (0.0%)
2	B	0.78	1/1810 (0.1%)	0.93	1/2458 (0.0%)
All	All	0.80	3/3539 (0.1%)	0.94	2/4804 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	103	MET	SD-CE	-10.73	1.52	1.79
2	B	142	MET	SD-CE	-6.63	1.62	1.79
1	A	174	MET	SD-CE	-5.76	1.65	1.79

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	VAL	CB-CA-C	-6.11	102.61	110.98
2	B	209	VAL	N-CA-C	5.34	116.06	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1683	0	1660	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1764	0	1726	37	0
3	A	12	0	16	6	0
3	B	18	0	24	5	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	59	0	0	0	0
5	B	54	0	0	1	0
All	All	3598	0	3432	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:TRP:HE1	1:A:282:ASN:HD22	1.37	0.72
1:A:138:LEU:HG	1:A:142:MET:HE2	1.72	0.71
1:A:241:LEU:HD13	3:A:1304:GOL:H2	1.73	0.70
1:A:103:MET:HE1	1:A:112:MET:SD	2.32	0.69
2:B:264:HIS:ND1	2:B:265:PRO:HD2	2.07	0.69
2:B:169:HIS:HB2	2:B:271:ILE:HG23	1.80	0.63
2:B:205:GLY:HA3	2:B:271:ILE:HD12	1.81	0.63
2:B:191:PHE:CE1	2:B:264:HIS:HE1	2.18	0.60
2:B:244:THR:O	3:B:1304:GOL:O1	2.20	0.57
2:B:264:HIS:CD2	2:B:268:LEU:HD12	2.40	0.56
2:B:206:HIS:HB3	3:B:1304:GOL:H12	1.86	0.56
1:A:190:GLN:HG2	1:A:194:LEU:HD22	1.89	0.53
1:A:110:TYR:CE1	1:A:248:ARG:HG2	2.44	0.53
1:A:171:TYR:OH	1:A:206:HIS:HA	2.08	0.53
2:B:277:TRP:CE3	2:B:291:ARG:HD3	2.44	0.53
1:A:106:ILE:HD12	1:A:249:PHE:HB2	1.91	0.53
1:A:142:MET:HE3	1:A:161:PHE:CE2	2.43	0.53
1:A:142:MET:HE1	1:A:157:LEU:CD1	2.39	0.52
1:A:223:ARG:NH2	2:B:123:ALA:O	2.43	0.52
2:B:109:MET:HE1	3:B:1302:GOL:O2	2.10	0.51
1:A:110:TYR:CZ	1:A:248:ARG:HD3	2.46	0.51
2:B:196:HIS:CE1	2:B:242:ILE:HD11	2.47	0.50
1:A:109:MET:HE3	3:B:1302:GOL:H32	1.94	0.50
1:A:224:LEU:HD21	2:B:112:MET:HE2	1.95	0.49
1:A:197:VAL:HG13	1:A:224:LEU:HB3	1.96	0.48
1:A:101:LEU:HD22	1:A:128:ILE:HG23	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:277:TRP:CD2	2:B:291:ARG:HD3	2.49	0.47
2:B:113:PRO:HB2	2:B:114:PRO:HD3	1.96	0.47
1:A:142:MET:HE1	1:A:157:LEU:HD13	1.96	0.47
2:B:244:THR:HG23	2:B:264:HIS:CD2	2.49	0.47
2:B:199:VAL:HA	2:B:226:VAL:O	2.15	0.46
1:A:223:ARG:HD2	3:A:1304:GOL:O2	2.16	0.46
2:B:274:ASN:HB2	2:B:276:PHE:CE1	2.51	0.46
1:A:255:VAL:HB	1:A:256:PRO:HD3	1.97	0.46
2:B:111:PHE:O	2:B:114:PRO:HD2	2.16	0.46
2:B:199:VAL:HG13	3:B:1304:GOL:H2	1.97	0.45
3:A:1304:GOL:H32	2:B:120:ALA:HB2	1.99	0.45
2:B:192:SER:OG	2:B:219:LYS:O	2.25	0.45
2:B:200:VAL:O	2:B:207:GLY:HA2	2.17	0.45
2:B:255:VAL:HB	2:B:256:PRO:HD3	1.98	0.45
1:A:105:ASP:HB2	2:B:230:ILE:HD13	2.00	0.44
3:A:1303:GOL:H2	2:B:228:HIS:CD2	2.52	0.44
2:B:248:ARG:NH1	5:B:2052:HOH:O	2.49	0.44
2:B:191:PHE:CE1	2:B:264:HIS:CE1	3.03	0.44
1:A:110:TYR:O	1:A:252:ARG:NH1	2.50	0.43
2:B:98:THR:HA	2:B:127:GLN:O	2.19	0.43
2:B:119:LEU:HD22	2:B:126:ILE:HD11	2.01	0.43
2:B:148:ASP:O	2:B:149:LEU:HD23	2.17	0.43
1:A:149:LEU:HD23	1:A:277:TRP:HB3	2.01	0.42
2:B:110:TYR:O	2:B:252:ARG:NH1	2.52	0.42
1:A:165:ARG:NH1	1:A:168:ARG:HD3	2.34	0.42
2:B:191:PHE:CZ	2:B:264:HIS:HE1	2.37	0.42
2:B:176:ARG:HA	2:B:259:LEU:HD12	2.01	0.41
1:A:262:SER:OG	1:A:263:PRO:O	2.38	0.41
2:B:173:CYS:O	2:B:261:THR:HA	2.21	0.41
1:A:103:MET:HE3	1:A:108:GLU:HA	2.02	0.41
1:A:241:LEU:CD1	3:A:1304:GOL:H2	2.48	0.41
2:B:244:THR:CG2	2:B:264:HIS:HE2	2.33	0.41
2:B:153:LEU:O	2:B:154:LEU:C	2.64	0.41
3:A:1303:GOL:H2	2:B:228:HIS:CG	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/315 (67%)	207 (98%)	5 (2%)	0	100	100
2	B	225/315 (71%)	220 (98%)	5 (2%)	0	100	100
All	All	437/630 (69%)	427 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/272 (65%)	148 (84%)	28 (16%)	2	4
2	B	183/272 (67%)	152 (83%)	31 (17%)	2	4
All	All	359/544 (66%)	300 (84%)	59 (16%)	2	4

All (59) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	THR
1	A	97	ARG
1	A	100	ASN
1	A	101	LEU
1	A	131	LEU
1	A	143	GLU
1	A	156	GLU
1	A	166	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	169	HIS
1	A	172	VAL
1	A	174	MET
1	A	183	LYS
1	A	194	LEU
1	A	197	VAL
1	A	199	VAL
1	A	204	THR
1	A	208	GLU
1	A	220	ARG
1	A	223	ARG
1	A	224	LEU
1	A	241	LEU
1	A	245	VAL
1	A	248	ARG
1	A	251	VAL
1	A	259	LEU
1	A	262	SER
1	A	290	LEU
1	A	293	LEU
2	B	87	ARG
2	B	88	ASP
2	B	96	THR
2	B	97	ARG
2	B	101	LEU
2	B	109	MET
2	B	129	SER
2	B	142	MET
2	B	143	GLU
2	B	154	LEU
2	B	163	GLN
2	B	165	ARG
2	B	166	LEU
2	B	172	VAL
2	B	177	LYS
2	B	183	LYS
2	B	192	SER
2	B	194	LEU
2	B	199	VAL
2	B	210	ASP
2	B	220	ARG
2	B	223	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	251	VAL
2	B	252	ARG
2	B	259	LEU
2	B	260	THR
2	B	262	SER
2	B	267	LYS
2	B	271	ILE
2	B	290	LEU
2	B	293	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	282	ASN
2	B	137	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	A	1305	-	3,3,3	0.85	0	3,3,3	0.88	0
3	GOL	B	1302	-	5,5,5	0.44	0	5,5,5	0.37	0
4	ACT	B	1305	-	3,3,3	0.81	0	3,3,3	1.12	0
3	GOL	A	1303	-	5,5,5	0.49	0	5,5,5	0.61	0
3	GOL	A	1304	-	5,5,5	0.43	0	5,5,5	0.53	0
3	GOL	B	1303	-	5,5,5	0.39	0	5,5,5	0.31	0
3	GOL	B	1304	-	5,5,5	0.51	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1302	-	-	2/4/4/4	-
3	GOL	A	1303	-	-	3/4/4/4	-
3	GOL	A	1304	-	-	3/4/4/4	-
3	GOL	B	1303	-	-	2/4/4/4	-
3	GOL	B	1304	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1303	GOL	O1-C1-C2-C3
3	A	1304	GOL	C1-C2-C3-O3
3	B	1304	GOL	C1-C2-C3-O3
3	B	1304	GOL	O2-C2-C3-O3
3	B	1302	GOL	O1-C1-C2-O2
3	A	1303	GOL	C1-C2-C3-O3
3	A	1304	GOL	O1-C1-C2-C3
3	B	1302	GOL	O1-C1-C2-C3
3	B	1303	GOL	C1-C2-C3-O3
3	A	1304	GOL	O2-C2-C3-O3
3	B	1303	GOL	O2-C2-C3-O3
3	A	1303	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1302	GOL	2	0
3	A	1303	GOL	2	0
3	A	1304	GOL	4	0
3	B	1304	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/315 (67%)	0.35	11 (5%) 33 28	33, 45, 61, 70	5 (2%)
2	B	227/315 (72%)	0.60	20 (8%) 15 12	35, 49, 83, 85	4 (1%)
All	All	441/630 (70%)	0.48	31 (7%) 22 18	33, 47, 65, 85	9 (2%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	82	THR	7.1
2	B	301	ALA	6.7
2	B	75	TYR	5.8
1	A	89	SER	5.6
2	B	78	ASN	5.1
2	B	77	LEU	4.9
2	B	80	LEU	4.6
2	B	76	ALA	4.5
2	B	83	ALA	3.7
1	A	93	PHE	3.7
1	A	302	HIS	3.5
2	B	85	THR	3.4
2	B	84	LEU	3.3
2	B	136	GLY	3.3
2	B	79	THR	3.2
1	A	136	GLY	3.2
2	B	81	GLN	3.2
2	B	87	ARG	3.0
1	A	156	GLU	2.9
2	B	203	ASN	2.6
2	B	121	GLN	2.5
1	A	135	ALA	2.5
2	B	159	THR	2.5
1	A	91	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	158	GLN	2.3
1	A	178	ASP	2.2
1	A	121	GLN	2.2
2	B	180	PRO	2.1
2	B	140	GLU	2.1
2	B	158	GLN	2.0
1	A	301	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	1304	6/6	0.84	0.30	55,57,58,61	0
4	ACT	B	1305	4/4	0.89	0.15	55,55,55,55	0
3	GOL	B	1302	6/6	0.90	0.20	69,72,73,73	0
3	GOL	A	1303	6/6	0.91	0.17	62,64,66,67	0
3	GOL	B	1303	6/6	0.92	0.13	56,56,57,57	0
4	ACT	A	1305	4/4	0.93	0.13	54,55,55,55	0
3	GOL	A	1304	6/6	0.93	0.15	58,59,60,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.