



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:12 PM UTC

PDB ID : 2UU7 / pdb_00002uu7
Title : Crystal structure of apo glutamine synthetase from dog (*Canis familiaris*)
Authors : Krajewski, W.W.; Jones, T.A.; Mowbray, S.L.
Deposited on : 2007-02-28
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	
1	B	381	
1	C	381	
1	D	381	
1	E	381	
1	F	381	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	381	<p>2% 77% 18% . .</p>
1	H	381	<p>5% 77% 18% . .</p>
1	I	381	<p>4% 77% 18% . .</p>
1	J	381	<p>4% 78% 17% . .</p>
1	K	381	<p>5% 80% 15% . .</p>
1	L	381	<p>6% 78% 18% . .</p>
1	M	381	<p>6% 79% 15% . . .</p>
1	N	381	<p>4% 81% 15% . . .</p>
1	O	381	<p>5% 78% 17% . .</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 44025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMINE SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	2933	1846	517	549	21	0	0	0
1	B	370	2933	1846	517	549	21	0	0	0
1	C	370	2933	1846	517	549	21	0	0	0
1	D	370	2933	1846	517	549	21	0	0	0
1	E	370	2933	1846	517	549	21	0	0	0
1	F	370	2933	1846	517	549	21	0	0	0
1	G	370	2933	1846	517	549	21	0	0	0
1	H	370	2933	1846	517	549	21	0	0	0
1	I	370	2933	1846	517	549	21	0	0	0
1	J	370	2933	1846	517	549	21	0	0	0
1	K	370	2933	1846	517	549	21	0	0	0
1	L	370	2933	1846	517	549	21	0	0	0
1	M	370	2933	1846	517	549	21	0	0	0
1	N	370	2933	1846	517	549	21	0	0	0
1	O	370	2933	1846	517	549	21	0	0	0

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0
2	G	1	Total Mg 1 1	0	0
2	H	1	Total Mg 1 1	0	0
2	I	1	Total Mg 1 1	0	0
2	J	1	Total Mg 1 1	0	0
2	K	1	Total Mg 1 1	0	0
2	L	1	Total Mg 1 1	0	0
2	M	1	Total Mg 1 1	0	0
2	N	1	Total Mg 1 1	0	0
2	O	1	Total Mg 1 1	0	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0
3	E	1	Total Cl 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total 1	Cl 1	0	0
3	G	1	Total 1	Cl 1	0	0
3	H	1	Total 1	Cl 1	0	0
3	I	1	Total 1	Cl 1	0	0
3	J	1	Total 1	Cl 1	0	0
3	K	1	Total 1	Cl 1	0	0
3	L	1	Total 1	Cl 1	0	0
3	M	1	Total 1	Cl 1	0	0
3	N	1	Total 1	Cl 1	0	0
3	O	1	Total 1	Cl 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLUTAMINE SYNTHETASE




- Molecule 1: GLUTAMINE SYNTHETASE






- Molecule 1: GLUTAMINE SYNTHETASE












- Molecule 1: GLUTAMINE SYNTHETASE




- Molecule 1: GLUTAMINE SYNTHETASE




- Molecule 1: GLUTAMINE SYNTHETASE





- Molecule 1: GLUTAMINE SYNTHETASE









• Molecule 1: GLUTAMINE SYNTHETASE




• Molecule 1: GLUTAMINE SYNTHETASE






• Molecule 1: GLUTAMINE SYNTHETASE










• Molecule 1: GLUTAMINE SYNTHETASE




• Molecule 1: GLUTAMINE SYNTHETASE




• Molecule 1: GLUTAMINE SYNTHETASE





• Molecule 1: GLUTAMINE SYNTHETASE





• Molecule 1: GLUTAMINE SYNTHETASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	183.59Å 485.60Å 192.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.99 – 3.00 29.99 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.9 (29.99-3.00) 98.8 (29.99-3.00)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 3.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.213 , 0.230 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtrriage
Anisotropy	0.324	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 26.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	44025	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/3014	0.93	3/4075 (0.1%)
1	B	0.76	1/3014 (0.0%)	0.94	1/4075 (0.0%)
1	C	0.78	1/3014 (0.0%)	0.95	3/4075 (0.1%)
1	D	0.75	1/3014 (0.0%)	0.95	4/4075 (0.1%)
1	E	0.79	1/3014 (0.0%)	0.96	2/4075 (0.0%)
1	F	0.78	0/3014	0.95	2/4075 (0.0%)
1	G	0.77	1/3014 (0.0%)	0.94	4/4075 (0.1%)
1	H	0.79	1/3014 (0.0%)	0.94	2/4075 (0.0%)
1	I	0.80	0/3014	0.96	3/4075 (0.1%)
1	J	0.79	1/3014 (0.0%)	0.95	4/4075 (0.1%)
1	K	0.79	2/3014 (0.1%)	0.95	5/4075 (0.1%)
1	L	0.78	1/3014 (0.0%)	0.97	4/4075 (0.1%)
1	M	0.78	1/3014 (0.0%)	0.94	4/4075 (0.1%)
1	N	0.83	1/3014 (0.0%)	0.92	4/4075 (0.1%)
1	O	0.80	1/3014 (0.0%)	0.93	4/4075 (0.1%)
All	All	0.78	13/45210 (0.0%)	0.95	49/61125 (0.1%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	3	THR	CA-CB	6.67	1.72	1.54
1	K	3	THR	CA-CB	6.66	1.72	1.54
1	D	3	THR	CA-CB	6.54	1.72	1.54
1	G	3	THR	CA-CB	6.28	1.71	1.54
1	B	3	THR	CA-CB	6.18	1.71	1.54

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	193	THR	CB-CA-C	-7.46	97.90	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	193	THR	CB-CA-C	-7.02	98.63	110.43
1	K	193	THR	CB-CA-C	-6.86	98.91	110.43
1	L	193	THR	CB-CA-C	-6.64	99.01	110.95
1	M	193	THR	CB-CA-C	-6.62	99.31	110.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2933	0	2796	44	0
1	B	2933	0	2796	40	0
1	C	2933	0	2796	45	0
1	D	2933	0	2796	49	0
1	E	2933	0	2796	43	0
1	F	2933	0	2796	49	0
1	G	2933	0	2796	50	0
1	H	2933	0	2796	50	8
1	I	2933	0	2796	47	2
1	J	2933	0	2796	44	2
1	K	2933	0	2796	45	2
1	L	2933	0	2796	47	6
1	M	2933	0	2796	41	4
1	N	2933	0	2796	39	0
1	O	2933	0	2796	48	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	K	1	0	0	0	0
2	L	1	0	0	0	0
2	M	1	0	0	0	0
2	N	1	0	0	0	0
2	O	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
3	M	1	0	0	0	0
3	N	1	0	0	0	0
3	O	1	0	0	0	0
All	All	44025	0	41940	627	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 627 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:276:LYS:NZ	1:L:362:ASN:HD21	1.82	0.78
1:H:193:THR:HG22	1:H:203:GLU:O	1.90	0.72
1:D:215:GLY:HA3	1:D:349:PHE:CE2	2.25	0.72
1:C:193:THR:HG22	1:C:203:GLU:O	1.93	0.69
1:A:193:THR:HG22	1:A:203:GLU:O	1.92	0.69

The worst 5 of 12 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:119:ARG:NH2	1:M:264:GLU:OE2[1_554]	1.49	0.71
1:H:106:ARG:CZ	1:L:263:GLU:OE1[3_555]	1.66	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:106:ARG:NH2	1:L:263:GLU:CB[3_555]	1.80	0.40
1:J:331:GLN:O	1:M:126:ASN:ND2[1_554]	1.85	0.35
1:H:106:ARG:NE	1:L:263:GLU:OE1[3_555]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/381 (97%)	354 (96%)	12 (3%)	2 (0%)	24	60
1	B	368/381 (97%)	350 (95%)	16 (4%)	2 (0%)	24	60
1	C	368/381 (97%)	348 (95%)	18 (5%)	2 (0%)	24	60
1	D	368/381 (97%)	352 (96%)	15 (4%)	1 (0%)	36	70
1	E	368/381 (97%)	350 (95%)	17 (5%)	1 (0%)	36	70
1	F	368/381 (97%)	350 (95%)	16 (4%)	2 (0%)	24	60
1	G	368/381 (97%)	348 (95%)	18 (5%)	2 (0%)	24	60
1	H	368/381 (97%)	350 (95%)	17 (5%)	1 (0%)	36	70
1	I	368/381 (97%)	347 (94%)	19 (5%)	2 (0%)	24	60
1	J	368/381 (97%)	350 (95%)	16 (4%)	2 (0%)	24	60
1	K	368/381 (97%)	352 (96%)	14 (4%)	2 (0%)	24	60
1	L	368/381 (97%)	351 (95%)	16 (4%)	1 (0%)	36	70
1	M	368/381 (97%)	350 (95%)	16 (4%)	2 (0%)	24	60
1	N	368/381 (97%)	351 (95%)	15 (4%)	2 (0%)	24	60
1	O	368/381 (97%)	352 (96%)	15 (4%)	1 (0%)	36	70
All	All	5520/5715 (97%)	5255 (95%)	240 (4%)	25 (0%)	24	60

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	GLY
1	B	72	GLY
1	C	72	GLY
1	D	72	GLY
1	E	72	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/320 (97%)	288 (93%)	22 (7%)	13	43
1	B	310/320 (97%)	290 (94%)	20 (6%)	15	47
1	C	310/320 (97%)	293 (94%)	17 (6%)	19	53
1	D	310/320 (97%)	291 (94%)	19 (6%)	17	49
1	E	310/320 (97%)	292 (94%)	18 (6%)	18	51
1	F	310/320 (97%)	292 (94%)	18 (6%)	18	51
1	G	310/320 (97%)	292 (94%)	18 (6%)	18	51
1	H	310/320 (97%)	293 (94%)	17 (6%)	19	53
1	I	310/320 (97%)	291 (94%)	19 (6%)	17	49
1	J	310/320 (97%)	289 (93%)	21 (7%)	14	45
1	K	310/320 (97%)	293 (94%)	17 (6%)	19	53
1	L	310/320 (97%)	290 (94%)	20 (6%)	15	47
1	M	310/320 (97%)	291 (94%)	19 (6%)	17	49
1	N	310/320 (97%)	291 (94%)	19 (6%)	17	49
1	O	310/320 (97%)	292 (94%)	18 (6%)	18	51
All	All	4650/4800 (97%)	4368 (94%)	282 (6%)	17	49

5 of 282 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	71	GLU
1	M	273	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	N	305	GLU
1	F	4	SER
1	E	364	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 112 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	304	HIS
1	O	331	GLN
1	J	304	HIS
1	O	304	HIS
1	N	226	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 30 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9	
1	A	370/381 (97%)	0.10	12 (3%)	50	29	14, 38, 80, 121	0
1	B	370/381 (97%)	0.16	16 (4%)	40	21	14, 38, 80, 121	0
1	C	370/381 (97%)	0.16	12 (3%)	50	29	14, 39, 80, 121	0
1	D	370/381 (97%)	0.10	17 (4%)	37	20	14, 38, 80, 121	0
1	E	370/381 (97%)	0.05	14 (3%)	44	24	14, 38, 80, 121	0
1	F	370/381 (97%)	0.28	15 (4%)	41	23	14, 39, 80, 121	0
1	G	370/381 (97%)	0.13	9 (2%)	59	36	14, 39, 80, 121	0
1	H	370/381 (97%)	0.21	18 (4%)	35	18	14, 38, 80, 121	0
1	I	370/381 (97%)	0.19	17 (4%)	37	20	14, 39, 80, 121	0
1	J	370/381 (97%)	0.18	16 (4%)	40	21	14, 39, 80, 121	0
1	K	370/381 (97%)	0.15	19 (5%)	33	17	14, 38, 80, 121	0
1	L	370/381 (97%)	0.38	24 (6%)	25	13	14, 39, 80, 121	0
1	M	370/381 (97%)	0.17	22 (5%)	28	14	14, 39, 80, 121	0
1	N	370/381 (97%)	0.10	17 (4%)	37	20	14, 38, 80, 121	0
1	O	370/381 (97%)	0.08	18 (4%)	35	18	13, 38, 80, 121	0
All	All	5550/5715 (97%)	0.16	246 (4%)	39	21	13, 39, 80, 121	0

The worst 5 of 246 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	126	ASN	5.6
1	H	372	LYS	5.5
1	D	369	PHE	5.3
1	K	369	PHE	5.0
1	M	305	GLU	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	J	401	1/1	0.76	0.21	29,29,29,29	0
2	MG	N	401	1/1	0.76	0.26	29,29,29,29	0
2	MG	M	401	1/1	0.79	0.20	29,29,29,29	0
2	MG	B	401	1/1	0.80	0.24	29,29,29,29	0
2	MG	E	401	1/1	0.82	0.23	29,29,29,29	0
2	MG	K	401	1/1	0.82	0.21	29,29,29,29	0
2	MG	H	401	1/1	0.83	0.20	29,29,29,29	0
2	MG	I	401	1/1	0.83	0.13	30,30,30,30	0
2	MG	A	401	1/1	0.83	0.22	29,29,29,29	0
2	MG	D	401	1/1	0.84	0.14	29,29,29,29	0
2	MG	C	401	1/1	0.85	0.15	30,30,30,30	0
2	MG	F	401	1/1	0.86	0.13	29,29,29,29	0
2	MG	L	401	1/1	0.86	0.15	29,29,29,29	0
3	CL	J	1373	1/1	0.86	0.20	66,66,66,66	0
2	MG	G	401	1/1	0.89	0.14	29,29,29,29	0
2	MG	O	401	1/1	0.91	0.12	29,29,29,29	0
3	CL	K	1373	1/1	0.91	0.11	45,45,45,45	0
3	CL	A	1373	1/1	0.92	0.12	50,50,50,50	0
3	CL	F	1373	1/1	0.92	0.20	64,64,64,64	0
3	CL	G	1373	1/1	0.93	0.14	59,59,59,59	0
3	CL	H	1373	1/1	0.93	0.13	55,55,55,55	0
3	CL	O	1373	1/1	0.93	0.11	38,38,38,38	0
3	CL	M	1373	1/1	0.94	0.07	42,42,42,42	0
3	CL	I	1373	1/1	0.94	0.08	33,33,33,33	0
3	CL	E	1373	1/1	0.95	0.08	33,33,33,33	0
3	CL	B	1373	1/1	0.95	0.15	51,51,51,51	0
3	CL	C	1373	1/1	0.95	0.11	62,62,62,62	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CL	D	1373	1/1	0.96	0.08	38,38,38,38	0
3	CL	N	1373	1/1	0.97	0.09	26,26,26,26	0
3	CL	L	1373	1/1	0.97	0.10	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.