



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 05:08 AM UTC

PDB ID : 2UUF / pdb_00002uuf
Title : Thrombin-hirugen binary complex at 1.26Å resolution
Authors : Ahmed, H.U.; Blakeley, M.P.; Cianci, M.; Cruickshank, D.W.J.; Hubbard, J.A.; Helliwell, J.R.
Deposited on : 2007-03-02
Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	36	28% 47% 19% 8% • 22%
2	B	259	34% 78% 17% • •
3	H	10	80% 80% 20%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN ALPHA THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	28	Total	C	N	O	S	0	1	0
			236	149	38	48	1			

- Molecule 2 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	252	Total	C	N	O	S	0	7	0
			2060	1315	362	368	15			

- Molecule 3 is a protein called HIRUDIN I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	10	Total	C	N	O	S	0	0	0
			94	59	10	24	1			

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	42	Total	O	0	0
			42	42		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	297	Total	O	0	0
			297	297		
6	H	7	Total	O	0	0
			7	7		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: HUMAN ALPHA THROMBIN



• Molecule 2: THROMBIN



• Molecule 3: HIRUDIN I



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	70.28Å 71.29Å 72.04Å 90.00° 99.90° 90.00°	Depositor
Resolution (Å)	49.39 – 1.26 49.39 – 1.26	Depositor EDS
% Data completeness (in resolution range)	93.0 (49.39-1.26) 94.7 (49.39-1.26)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.26Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.146 , 0.194 (Not available) , 0.264	Depositor DCC
R_{free} test set	4490 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	13.0	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2738	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	0/241	1.57	10/320 (3.1%)
2	B	1.10	9/2139 (0.4%)	1.60	36/2890 (1.2%)
3	H	1.08	1/78 (1.3%)	1.44	0/103
All	All	1.08	10/2458 (0.4%)	1.59	46/3313 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	247	GLU	C-O	22.14	1.67	1.23
2	B	186(A)	ASP	N-CA	12.04	1.62	1.46
2	B	186	PRO	C-N	-11.86	1.06	1.34
2	B	33[A]	LEU	C-N	-7.18	1.23	1.33
2	B	33[B]	LEU	C-N	-7.18	1.23	1.33
2	B	32	MET	C-N	-7.01	1.23	1.33
2	B	186(A)	ASP	C-O	6.95	1.33	1.24
2	B	168	CYS	CA-CB	6.20	1.63	1.53
2	B	91	HIS	CG-ND1	-6.16	1.31	1.38
3	H	61	GLU	C-N	5.87	1.41	1.33

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	91	HIS	ND1-CG-CD2	-10.42	95.68	106.10
2	B	92	PRO	CA-C-N	9.95	138.13	122.60
2	B	92	PRO	C-N-CA	9.95	138.13	122.60
2	B	91	HIS	ND1-CE1-NE2	-9.05	99.35	108.40
2	B	91	HIS	CG-ND1-CE1	8.86	124.36	109.30
2	B	110	LYS	O-C-N	8.63	127.23	121.71
2	B	186(D)	LYS	CA-C-O	-8.33	112.14	121.89
2	B	75	ARG	NE-CZ-NH1	-8.32	113.18	121.50
2	B	75	ARG	CB-CA-C	-8.26	96.01	109.72
2	B	92	PRO	O-C-N	-8.18	111.17	122.30
2	B	232	PHE	CA-CB-CG	-7.95	105.85	113.80
2	B	75	ARG	CG-CD-NE	7.89	129.35	112.00
2	B	173	ARG	NE-CZ-NH1	-7.78	113.72	121.50
2	B	205[A]	ASN	CA-CB-CG	-7.71	104.89	112.60
2	B	205[B]	ASN	CA-CB-CG	-7.71	104.89	112.60
2	B	186(D)	LYS	O-C-N	7.57	131.53	122.68
2	B	101	ARG	NE-CZ-NH2	7.13	125.62	119.20
2	B	247	GLU	CA-C-O	-6.83	109.20	120.80
2	B	127	GLU	CB-CG-CD	6.69	123.97	112.60
1	A	7	PHE	CA-CB-CG	-6.49	107.31	113.80
2	B	230	HIS	CA-CB-CG	6.37	120.17	113.80
1	A	9[A]	LYS	CA-C-N	6.33	133.64	121.54
1	A	9[A]	LYS	C-N-CA	6.33	133.64	121.54
1	A	9[B]	LYS	CA-C-N	6.33	133.64	121.54
1	A	9[B]	LYS	C-N-CA	6.33	133.64	121.54
2	B	38	GLN	CB-CG-CD	6.30	123.32	112.60
1	A	14(K)	ILE	CB-CA-C	-6.17	104.16	111.94
2	B	33[A]	LEU	O-C-N	6.15	130.22	123.27
2	B	33[B]	LEU	O-C-N	6.15	130.22	123.27
2	B	34	PHE	CA-CB-CG	5.98	119.78	113.80
1	A	1(A)	ASP	CA-CB-CG	-5.83	106.77	112.60
2	B	77(A)	ARG	O-C-N	-5.69	116.23	123.06
2	B	49	ASP	CA-CB-CG	5.69	118.29	112.60
2	B	217	GLU	CB-CG-CD	5.55	122.03	112.60
2	B	192	GLU	CB-CG-CD	5.48	121.92	112.60
1	A	14(K)	ILE	CA-C-O	5.29	126.33	120.46
2	B	33[A]	LEU	CA-C-N	-5.21	114.71	122.74
2	B	33[A]	LEU	C-N-CA	-5.21	114.71	122.74
2	B	33[B]	LEU	CA-C-N	-5.21	114.71	122.74
2	B	33[B]	LEU	C-N-CA	-5.21	114.71	122.74
2	B	178	ASP	O-C-N	-5.17	115.50	122.43
1	A	14(D)	ARG	NE-CZ-NH2	-5.13	114.58	119.20
2	B	76	TYR	CA-CB-CG	-5.13	104.67	113.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	91	HIS	CB-CG-CD2	5.07	137.79	131.20
2	B	60(B)	PRO	N-CA-C	5.04	116.85	110.70
1	A	1(A)	ASP	N-CA-CB	-5.00	103.24	110.70

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	14(L)	ASP	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	14(J)	TYR	Mainchain
1	A	9[A]	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	236	0	242	23	0
2	B	2060	0	2039	25	0
3	H	94	0	73	0	0
4	B	1	0	0	0	0
5	B	1	0	0	0	0
6	A	42	0	0	7	0
6	B	297	0	0	8	1
6	H	7	0	0	0	0
All	All	2738	0	2354	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:247:GLU:C	2:B:247:GLU:O	1.67	1.35

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:33[B]:LEU:HD23	2:B:42[B]:CYS:SG	2.01	1.00
1:A:5:PRO:HA	1:A:9[B]:LYS:HD2	1.49	0.95
1:A:9[B]:LYS:HE3	6:A:1012:HOH:O	1.67	0.92
1:A:14(K):ILE:HG13	1:A:14(L):ASP:N	1.90	0.87
2:B:33[B]:LEU:CD2	2:B:42[B]:CYS:SG	2.66	0.84
2:B:60(I):THR:HG23	6:B:1083:HOH:O	1.80	0.80
1:A:14(K):ILE:O	1:A:14(L):ASP:OD1	2.07	0.72
1:A:14(K):ILE:O	1:A:14(L):ASP:CB	2.43	0.65
2:B:217:GLU:HG3	6:B:1278:HOH:O	1.96	0.64
1:A:5:PRO:HA	1:A:9[B]:LYS:CD	2.26	0.62
1:A:14(K):ILE:CG1	1:A:14(L):ASP:N	2.62	0.61
2:B:186(C):GLY:HA2	6:B:1230:HOH:O	2.00	0.61
1:A:9[B]:LYS:CE	6:A:1012:HOH:O	2.39	0.59
1:A:14(K):ILE:O	1:A:14(L):ASP:HB2	2.03	0.58
1:A:1(A):ASP:OD1	1:A:9[B]:LYS:NZ	2.36	0.58
2:B:195:SER:OG	6:B:1244:HOH:O	2.17	0.55
1:A:1(B):ALA:CA	6:A:1002:HOH:O	2.55	0.54
1:A:9[A]:LYS:HE2	6:A:1010:HOH:O	2.05	0.54
1:A:14(K):ILE:C	1:A:14(L):ASP:OD1	2.51	0.53
1:A:1(B):ALA:N	6:A:1002:HOH:O	2.35	0.52
2:B:66:VAL:CG2	2:B:83:SER:HB2	2.40	0.51
2:B:42[B]:CYS:SG	2:B:58:CYS:HB3	2.50	0.51
2:B:151:GLN:HG2	6:B:1174:HOH:O	2.11	0.50
2:B:145:LYS:NZ	6:B:1176:HOH:O	2.38	0.49
1:A:1(B):ALA:C	6:A:1002:HOH:O	2.55	0.48
2:B:33[B]:LEU:HD22	2:B:42[B]:CYS:SG	2.52	0.48
1:A:14(K):ILE:O	1:A:14(L):ASP:CG	2.58	0.46
2:B:175:ARG:NH2	6:B:1212:HOH:O	2.45	0.46
2:B:35:ARG:O	2:B:38:GLN:HA	2.17	0.45
1:A:4:ARG:O	1:A:9[B]:LYS:HG3	2.16	0.44
2:B:77(A):ARG:O	2:B:78:ASN:HB2	2.17	0.44
2:B:33[A]:LEU:CD2	2:B:66:VAL:HG12	2.48	0.43
2:B:204(B):ASN:O	2:B:205[B]:ASN:HB3	2.19	0.43
1:A:14(L):ASP:CG	6:A:1027:HOH:O	2.61	0.43
2:B:17:VAL:O	2:B:188:GLY:HA2	2.20	0.42
2:B:211:GLY:HA2	2:B:229:THR:O	2.19	0.42
1:A:5:PRO:CA	1:A:9[B]:LYS:HD2	2.35	0.41
2:B:247:GLU:O	2:B:247:GLU:CA	2.60	0.41
1:A:14(K):ILE:C	1:A:14(L):ASP:CG	2.82	0.41
2:B:244:GLN:HG2	2:B:245:PHE:CE1	2.56	0.41
2:B:94:TYR:CZ	2:B:96:TRP:HB3	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:127:GLU:HG3	6:B:1159:HOH:O	2.20	0.41
2:B:42[B]:CYS:HB2	2:B:195:SER:O	2.21	0.40
2:B:66:VAL:O	2:B:66:VAL:HG23	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:1098:HOH:O	6:B:1098:HOH:O[2_657]	0.99	1.21

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/36 (75%)	25 (93%)	1 (4%)	1 (4%)	2	0
2	B	255/259 (98%)	247 (97%)	8 (3%)	0	100	100
3	H	7/10 (70%)	7 (100%)	0	0	100	100
All	All	289/305 (95%)	279 (96%)	9 (3%)	1 (0%)	30	14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	27/31 (87%)	27 (100%)	0	100	100
2	B	226/225 (100%)	221 (98%)	5 (2%)	45	10
3	H	9/9 (100%)	9 (100%)	0	100	100
All	All	262/265 (99%)	257 (98%)	5 (2%)	48	13

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	64	LEU
2	B	151	GLN
2	B	182	CYS
2	B	236	LYS
2	B	240	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	62	ASN
2	B	244	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TYS	H	63	3	15,16,17	1.94	4 (26%)	15,22,24	1.49	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	H	63	3	-	0/10/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	63	TYS	OH-CZ	-4.85	1.34	1.42
3	H	63	TYS	CE1-CD1	-3.05	1.33	1.38
3	H	63	TYS	CE2-CD2	-2.62	1.34	1.38
3	H	63	TYS	CE2-CZ	-2.46	1.34	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	63	TYS	O3-S-O2	4.24	123.39	108.56
3	H	63	TYS	OH-S-O1	-2.39	100.28	107.56
3	H	63	TYS	OH-S-O2	-2.21	100.83	107.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	186:PRO	C	186(A):ASP	N	1.06

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	28/36 (77%)	2.06	10 (35%)  	10, 15, 36, 122	1 (3%)
2	B	252/259 (97%)	1.79	87 (34%)  	8, 18, 38, 66	7 (2%)
3	H	9/10 (90%)	3.54	8 (88%)  	35, 39, 67, 68	0
All	All	289/305 (94%)	1.87	105 (36%)  	8, 18, 40, 122	8 (2%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14(L)	ASP	7.5
2	B	147	THR	7.2
1	A	9[A]	LYS	5.9
2	B	247	GLU	5.4
3	H	64	LEU	5.3
2	B	60(I)	THR	4.7
1	A	14(K)	ILE	4.7
3	H	58	GLU	4.4
3	H	55	ASP	4.3
2	B	75	ARG	4.2
2	B	60(D)	TRP	4.2
3	H	56	PHE	4.2
3	H	59	ILE	4.0
2	B	150	GLY	3.9
2	B	37(A)	PRO	3.9
3	H	60	PRO	3.9
2	B	113	ALA	3.7
2	B	41	LEU	3.6
1	A	1(B)	ALA	3.5
2	B	74	THR	3.5
2	B	204(A)	PHE	3.4
1	A	1(A)	ASP	3.3
2	B	62	ASN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	64	LEU	3.2
2	B	154	VAL	3.1
2	B	186(D)	LYS	3.0
2	B	138	VAL	3.0
2	B	151	GLN	3.0
2	B	66	VAL	2.9
2	B	144	LEU	2.9
2	B	109	LYS	2.9
2	B	34	PHE	2.9
2	B	129(C)	LEU	2.8
2	B	130[A]	LEU	2.8
2	B	153	SER	2.8
2	B	129(A)	ALA	2.8
2	B	40	LEU	2.8
2	B	96	TRP	2.8
2	B	76	TYR	2.8
2	B	134	TYR	2.8
2	B	241	VAL	2.8
2	B	108	LEU	2.8
2	B	127	GLU	2.8
2	B	112	VAL	2.7
2	B	234	LEU	2.7
2	B	207	TRP	2.7
2	B	186(C)	GLY	2.7
2	B	97(A)	GLU	2.7
2	B	29	TRP	2.7
2	B	65	LEU	2.6
2	B	85	LEU	2.6
1	A	1	CYS	2.6
2	B	38	GLN	2.5
2	B	155	LEU	2.5
2	B	79	ILE	2.5
2	B	242	ILE	2.5
1	A	7	PHE	2.5
2	B	132	ALA	2.5
2	B	141	TRP	2.5
2	B	245	PHE	2.4
2	B	24	ILE	2.4
2	B	231	VAL	2.4
2	B	37	SER	2.4
2	B	68	ILE	2.3
1	A	14(J)	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	129(B)[A]	SER	2.3
2	B	205[A]	ASN	2.3
2	B	204	PRO	2.3
2	B	36	LYS	2.3
2	B	110	LYS	2.3
2	B	60(H)	PHE	2.3
2	B	162	ILE	2.3
2	B	105	LEU	2.3
2	B	145	LYS	2.3
2	B	86	GLU	2.3
2	B	82	ILE	2.3
2	B	193	GLY	2.3
2	B	244	GLN	2.2
2	B	35	ARG	2.2
2	B	31	VAL	2.2
2	B	191	CYS	2.2
2	B	17	VAL	2.2
3	H	61	GLU	2.1
2	B	118	ILE	2.1
2	B	208	TYR	2.1
2	B	121	VAL	2.1
2	B	213	VAL	2.1
1	A	14(A)	LYS	2.1
2	B	114	PHE	2.1
2	B	90	ILE	2.1
2	B	103	ILE	2.1
2	B	117	TYR	2.1
2	B	225	TYR	2.1
3	H	62	GLU	2.1
2	B	42[A]	CYS	2.1
2	B	44	ALA	2.1
2	B	129	ALA	2.1
2	B	174	ILE	2.1
2	B	240	LYS	2.1
2	B	72	SER	2.0
2	B	60(G)	ASN	2.0
1	A	14(B)	THR	2.0
2	B	47	ILE	2.0
2	B	94	TYR	2.0
2	B	140	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TYS	H	63	16/17	0.91	0.13	33,38,42,44	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NA	B	2248	1/1	0.94	0.09	18,18,18,18	0
5	CA	B	2249	1/1	0.95	0.14	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.