



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:05 AM UTC

PDB ID : 2UUK / pdb_00002uuk
Title : Thrombin-hirugen-gw420128 ternary complex at 1.39Å resolution
Authors : Ahmed, H.U.; Blakeley, M.P.; Cianci, M.; Cruickshank, D.W.J.; Hubbard, J.A.; Helliwell, J.R.
Deposited on : 2007-03-03
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	36	53% 50% 25% 22%
2	B	259	64% 81% 14% ...
3	H	10	80% 60% 40%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN ALPHA THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	28	Total	C	N	O	S	0	3	0
			248	156	40	51	1			

- Molecule 2 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	252	Total	C	N	O	S	0	9	0
			2073	1322	364	373	14			

- Molecule 3 is a protein called HIRUDIN I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	10	Total	C	N	O	S	0	0	0
			94	59	10	24	1			


- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		

- Molecule 6 is N-[3-(TERT-BUTYLAMINO)-3-OXOPROPYL]-N-ISOPROPYL-3-METHYL-5-[[[(2S)-2-(PYRIDIN-4-YLAMINO)PROPYL]OXY}BENZAMIDE (CCD ID: 897) (formula: C₂₆H₃₈N₄O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			33	26	4	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	44	Total	O	0	0
			44	44		
7	B	294	Total	O	0	0
			294	294		
7	H	7	Total	O	0	0
			7	7		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HUMAN ALPHA THROMBIN



• Molecule 2: THROMBIN



• Molecule 3: HIRUDIN I



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	70.35Å 71.44Å 72.12Å 90.00° 100.03° 90.00°	Depositor
Resolution (Å)	18.55 – 1.39 18.55 – 1.40	Depositor EDS
% Data completeness (in resolution range)	95.9 (18.55-1.39) 95.9 (18.55-1.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.40Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.118 , 0.173 0.226 , 0.252	Depositor DCC
R_{free} test set	3345 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2795	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 897, CA, TYS, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/259	1.59	6/343 (1.7%)
2	B	1.92	4/2155 (0.2%)	1.50	27/2912 (0.9%)
3	H	1.94	1/78 (1.3%)	1.57	0/103
All	All	1.84	5/2492 (0.2%)	1.51	33/3358 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	247	GLU	C-O	78.32	2.80	1.23
3	H	64	LEU	C-O	16.64	1.56	1.23
2	B	157[A]	VAL	C-N	15.21	1.51	1.33
2	B	157[B]	VAL	C-N	15.21	1.51	1.33
2	B	128	THR	C-N	-9.60	1.21	1.33

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	75	ARG	NE-CZ-NH1	-14.45	107.05	121.50
1	A	14(D)	ARG	CD-NE-CZ	12.86	142.41	124.40
2	B	91	HIS	ND1-CG-CD2	-9.98	96.12	106.10
2	B	92	PRO	CA-C-N	9.78	137.85	122.60
2	B	92	PRO	C-N-CA	9.78	137.85	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	91	HIS	ND1-CE1-NE2	-8.78	99.62	108.40
2	B	92	PRO	O-C-N	-8.43	110.83	122.30
2	B	91	HIS	CG-ND1-CE1	8.28	123.37	109.30
2	B	50	ARG	CG-CD-NE	7.68	128.90	112.00
2	B	36	LYS	CA-C-O	7.43	128.29	120.42
2	B	101	ARG	NE-CZ-NH2	7.37	125.83	119.20
2	B	75	ARG	NH1-CZ-NH2	7.21	128.67	119.30
2	B	92	PRO	CA-C-O	6.77	128.47	119.00
2	B	186(A)	ASP	CA-CB-CG	6.63	119.23	112.60
2	B	232	PHE	CA-CB-CG	-6.53	107.27	113.80
1	A	1(A)	ASP	CA-CB-CG	-6.51	106.09	112.60
1	A	14(K)	ILE	CB-CA-C	-6.45	104.14	111.55
2	B	186(D)	LYS	CA-C-O	-5.99	114.75	121.81
2	B	63	ASP	CA-CB-CG	5.92	118.52	112.60
2	B	74	THR	CA-C-O	5.87	125.60	119.14
2	B	156	GLN	O-C-N	5.74	130.40	123.16
2	B	178	ASP	O-C-N	-5.65	114.86	122.43
2	B	75	ARG	NE-CZ-NH2	5.64	124.28	119.20
2	B	37(A)	PRO	N-CD-CG	-5.54	97.15	103.80
1	A	14(K)	ILE	CA-C-O	5.42	126.28	120.32
2	B	173	ARG	NE-CZ-NH1	-5.32	116.18	121.50
2	B	205[A]	ASN	CB-CA-C	-5.26	102.47	111.26
2	B	205[B]	ASN	CB-CA-C	-5.26	102.47	111.26
2	B	243	ASP	CA-CB-CG	5.19	117.79	112.60
2	B	143	ASN	CA-CB-CG	5.15	117.75	112.60
2	B	206	ARG	CD-NE-CZ	5.12	131.57	124.40
1	A	10[A]	LYS	CA-CB-CG	5.07	124.24	114.10
1	A	10[B]	LYS	CA-CB-CG	5.07	124.24	114.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	14(L)	ASP	CA

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	248	0	256	15	0
2	B	2073	0	2041	23	0
3	H	94	0	74	2	0
4	B	1	0	0	0	0
5	B	1	0	0	0	0
6	B	33	0	38	2	0
7	A	44	0	0	3	0
7	B	294	0	0	13	0
7	H	7	0	0	0	0
All	All	2795	0	2409	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:2250:897:H12	7:B:1278:HOH:O	1.55	1.02
1:A:14(K):ILE:HG13	1:A:14(L):ASP:N	1.90	0.87
1:A:9[B]:LYS:HE3	7:A:1010:HOH:O	1.85	0.77
1:A:14(K):ILE:O	1:A:14(L):ASP:OD1	2.07	0.72
1:A:9[B]:LYS:HG2	7:A:1009:HOH:O	1.95	0.67
2:B:53:LEU:HD23	7:B:1258:HOH:O	1.94	0.67
2:B:151:GLN:HG2	7:B:1174:HOH:O	1.95	0.66
1:A:14(K):ILE:O	1:A:14(L):ASP:CB	2.43	0.65
2:B:81:LYS:HG3	7:B:1146:HOH:O	1.97	0.64
2:B:53:LEU:CD2	7:B:1258:HOH:O	2.45	0.63
2:B:247:GLU:OE1	2:B:247:GLU:HA	2.00	0.62
2:B:209:GLN:CD	7:B:1258:HOH:O	2.42	0.61
1:A:14(K):ILE:CG1	1:A:14(L):ASP:N	2.63	0.61
1:A:14(A):LYS:HB2	7:B:1029:HOH:O	2.04	0.58
1:A:14(K):ILE:O	1:A:14(L):ASP:HB2	2.03	0.58
1:A:9[B]:LYS:NZ	7:A:1009:HOH:O	2.35	0.57
2:B:22:ALA:HB2	2:B:157[B]:VAL:HG23	1.86	0.57
2:B:75:ARG:NH1	7:B:1103:HOH:O	2.38	0.56
2:B:33[A]:LEU:CD2	2:B:66:VAL:HG12	2.37	0.55
2:B:35:ARG:HD2	2:B:37(A):PRO:O	2.07	0.55
2:B:139:THR:HG22	2:B:157[B]:VAL:HG22	1.88	0.54
1:A:14(K):ILE:C	1:A:14(L):ASP:OD1	2.51	0.53
6:B:2250:897:C12	7:B:1278:HOH:O	2.29	0.53
2:B:84:MET:HE3	7:B:1048:HOH:O	2.08	0.52
2:B:75:ARG:CZ	7:B:1103:HOH:O	2.58	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:30[B]:GLN:NE2	2:B:139:THR:OG1	2.43	0.48
1:A:13[B]:GLU:HG3	1:A:14(C):GLU:CD	2.38	0.48
2:B:22:ALA:HB2	2:B:157[B]:VAL:CG2	2.44	0.48
2:B:75:ARG:HG2	3:H:57:GLU:OE1	2.13	0.47
1:A:14(A):LYS:HD2	7:B:1029:HOH:O	2.14	0.47
2:B:66:VAL:CG2	2:B:83:SER:HB2	2.45	0.47
1:A:14(K):ILE:O	1:A:14(L):ASP:CG	2.58	0.46
2:B:60(F):LYS:NZ	7:B:1079:HOH:O	2.49	0.45
3:H:58:GLU:H	3:H:58:GLU:CD	2.22	0.44
2:B:75:ARG:HG2	2:B:76:TYR:N	2.33	0.43
1:A:3:LEU:O	1:A:9[B]:LYS:HE2	2.19	0.43
2:B:60(I):THR:HG22	2:B:63:ASP:OD2	2.19	0.42
2:B:75:ARG:HE	2:B:75:ARG:HB3	1.51	0.42
2:B:204(B):ASN:O	2:B:205[B]:ASN:HB3	2.20	0.41
2:B:211:GLY:HA2	2:B:229:THR:O	2.20	0.41
1:A:14(K):ILE:C	1:A:14(L):ASP:CG	2.82	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/36 (81%)	28 (97%)	1 (3%)	0	100	100
2	B	257/259 (99%)	249 (97%)	8 (3%)	0	100	100
3	H	7/10 (70%)	7 (100%)	0	0	100	100
All	All	293/305 (96%)	284 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/31 (94%)	29 (100%)	0	100	100
2	B	228/225 (101%)	225 (99%)	3 (1%)	61	30
3	H	9/9 (100%)	9 (100%)	0	100	100
All	All	266/265 (100%)	263 (99%)	3 (1%)	63	37

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	64	LEU
2	B	75	ARG
2	B	247	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TYS	H	63	3	15,16,17	1.01	1 (6%)	15,22,24	1.16	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	H	63	3	-	0/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	63	TYS	OH-S	-2.36	1.53	1.58

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	63	TYS	OH-S-O1	-3.84	95.85	107.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	897	B	2250	-	33,34,34	1.91	8 (24%)	43,47,47	2.18	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	897	B	2250	-	-	0/31/31/31	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	2250	897	C5-C4	-4.19	1.43	1.51
6	B	2250	897	C6-C5	-3.87	1.44	1.52
6	B	2250	897	C18-C17	-3.84	1.33	1.39
6	B	2250	897	C7-N3	-3.65	1.43	1.47
6	B	2250	897	O2-C21	3.43	1.29	1.22
6	B	2250	897	C2-C3	3.05	1.59	1.52
6	B	2250	897	C15-C19	2.64	1.43	1.39
6	B	2250	897	C16-C17	2.26	1.42	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	2250	897	C13-C12-N4	-6.82	111.97	123.60
6	B	2250	897	C12-N4-C11	5.78	130.09	116.86
6	B	2250	897	C10-C11-N4	-3.59	117.47	123.60
6	B	2250	897	O1-C4-N1	-3.47	117.59	123.09
6	B	2250	897	C17-C18-C19	3.46	124.48	121.16
6	B	2250	897	C14-C15-C19	3.32	123.23	119.67
6	B	2250	897	C12-C13-C9	3.15	123.02	118.88
6	B	2250	897	C18-C19-C15	-3.12	114.65	118.11
6	B	2250	897	C3-N1-C4	2.86	129.79	126.14
6	B	2250	897	O2-C21-C17	-2.60	115.13	120.29
6	B	2250	897	C5-C4-N1	2.44	119.84	115.77
6	B	2250	897	C20-C19-C18	2.36	124.12	120.92

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	2250	897	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	28/36 (77%)	2.53	19 (67%) 0 0	11, 15, 41, 122	3 (10%)
2	B	252/259 (97%)	2.42	165 (65%) 0 0	8, 18, 39, 69	9 (3%)
3	H	9/10 (90%)	3.44	8 (88%) 0 0	32, 37, 60, 71	0
All	All	289/305 (94%)	2.46	192 (66%) 0 0	8, 18, 42, 122	12 (4%)

All (192) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	37(A)	PRO	5.9
2	B	147	THR	5.9
1	A	14(L)	ASP	5.8
2	B	60(D)	TRP	5.5
1	A	14(K)	ILE	5.2
3	H	55	ASP	4.8
3	H	59	ILE	4.8
3	H	60	PRO	4.6
2	B	186(C)	GLY	4.6
2	B	186(A)	ASP	4.4
2	B	24	ILE	4.2
2	B	132	ALA	4.2
2	B	204(A)	PHE	4.1
2	B	207	TRP	4.1
2	B	75	ARG	4.0
2	B	128	THR	4.0
2	B	66	VAL	3.9
1	A	1(B)	ALA	3.9
2	B	29	TRP	3.9
1	A	7	PHE	3.9
3	H	56	PHE	3.8
2	B	183	ALA	3.8
3	H	64	LEU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	140	GLY	3.8
2	B	113	ALA	3.7
2	B	130	LEU	3.7
2	B	79	ILE	3.7
2	B	64	LEU	3.7
2	B	154	VAL	3.7
2	B	82	ILE	3.6
2	B	174	ILE	3.6
2	B	191	CYS	3.5
2	B	60(I)	THR	3.5
2	B	96	TRP	3.5
2	B	38	GLN	3.5
1	A	1	CYS	3.5
2	B	245	PHE	3.5
2	B	237	TRP	3.5
2	B	193	GLY	3.4
2	B	186	PRO	3.4
2	B	155	LEU	3.4
2	B	76	TYR	3.4
2	B	162	ILE	3.4
1	A	1(A)	ASP	3.4
2	B	134	TYR	3.4
2	B	208	TYR	3.4
2	B	150	GLY	3.4
2	B	141	TRP	3.4
2	B	153[A]	SER	3.3
2	B	168	CYS	3.3
2	B	186(D)	LYS	3.3
2	B	65	LEU	3.2
2	B	176[A]	ILE	3.2
2	B	205[A]	ASN	3.2
2	B	215	TRP	3.2
2	B	117	TYR	3.2
2	B	60(H)	PHE	3.2
2	B	46	LEU	3.2
2	B	129(C)	LEU	3.2
2	B	234	LEU	3.2
2	B	114	PHE	3.1
2	B	220	CYS	3.1
2	B	144	LEU	3.1
2	B	190	ALA	3.1
2	B	112	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	242	ILE	3.0
2	B	142	GLY	3.0
2	B	60(G)	ASN	3.0
2	B	20	SER	3.0
2	B	225	TYR	3.0
1	A	9[A]	LYS	3.0
2	B	167	VAL	3.0
2	B	213	VAL	3.0
2	B	40	LEU	3.0
2	B	94	TYR	3.0
2	B	31	VAL	3.0
2	B	231	VAL	3.0
2	B	226	GLY	3.0
2	B	232	PHE	3.0
1	A	6	LEU	3.0
2	B	51	TRP	2.9
2	B	120	PRO	2.9
2	B	199	PHE	2.9
2	B	238	ILE	2.9
2	B	74	THR	2.9
2	B	121	VAL	2.9
2	B	200	VAL	2.9
2	B	33[A]	LEU	2.9
2	B	103	ILE	2.9
2	B	247	GLU	2.8
2	B	161	PRO	2.8
2	B	127	GLU	2.8
2	B	204	PRO	2.8
2	B	44	ALA	2.8
2	B	99	LEU	2.8
2	B	34	PHE	2.8
2	B	181	PHE	2.8
2	B	90	ILE	2.8
2	B	145	LYS	2.7
1	A	12	LEU	2.7
1	A	14(G)	LEU	2.7
2	B	36	LYS	2.7
2	B	47	ILE	2.7
3	H	61	GLU	2.7
2	B	19	GLY	2.7
2	B	151	GLN	2.7
2	B	138	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	163	VAL	2.7
3	H	58	GLU	2.7
2	B	41	LEU	2.7
2	B	88	ILE	2.6
1	A	5	PRO	2.6
2	B	77(A)	ARG	2.6
2	B	89	TYR	2.6
2	B	158	VAL	2.6
2	B	241	VAL	2.6
2	B	123	LEU	2.6
1	A	14(I)	SER	2.6
2	B	72	SER	2.6
2	B	124	PRO	2.6
2	B	244	GLN	2.6
2	B	53	LEU	2.6
2	B	60	LEU	2.6
2	B	122	CYS	2.6
2	B	216	GLY	2.6
2	B	118	ILE	2.6
1	A	14(J)	TYR	2.5
2	B	25	GLY	2.5
2	B	219	GLY	2.5
2	B	108	LEU	2.5
2	B	182	CYS	2.5
2	B	111	PRO	2.5
2	B	166	PRO	2.5
2	B	110	LYS	2.5
1	A	14(F)	LEU	2.5
2	B	78	ASN	2.5
2	B	131	GLN	2.5
2	B	178	ASP	2.5
2	B	184(A)	TYR	2.5
1	A	14(D)	ARG	2.4
2	B	143	ASN	2.4
1	A	14(A)	LYS	2.4
2	B	228	TYR	2.4
2	B	71	HIS	2.4
2	B	115	SER	2.4
3	H	62	GLU	2.4
2	B	188	GLY	2.4
2	B	204(B)	ASN	2.4
2	B	85	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	17	VAL	2.4
2	B	69	GLY	2.4
1	A	3	LEU	2.3
2	B	60(A)	TYR	2.3
2	B	152	PRO	2.3
2	B	58	CYS	2.3
2	B	104	ALA	2.3
2	B	139	THR	2.3
2	B	229	THR	2.3
2	B	160	LEU	2.3
2	B	22	ALA	2.3
2	B	243	ASP	2.3
2	B	227	PHE	2.3
2	B	68	ILE	2.3
2	B	83	SER	2.2
2	B	186(B)	GLU	2.2
2	B	172	THR	2.2
2	B	133	GLY	2.2
2	B	60(C)	PRO	2.2
2	B	201	MET	2.2
2	B	105	LEU	2.2
2	B	16	ILE	2.2
2	B	55	ALA	2.2
2	B	129	ALA	2.2
2	B	84	MET	2.2
2	B	48	SER	2.2
2	B	239	GLN	2.1
2	B	26	MET	2.1
2	B	62	ASN	2.1
2	B	63	ASP	2.1
2	B	45	SER	2.1
2	B	109	LYS	2.1
2	B	177	THR	2.1
2	B	230	HIS	2.1
2	B	59	LEU	2.1
1	A	13[A]	GLU	2.1
2	B	98	ASN	2.0
2	B	106	MET	2.0
2	B	42	CYS	2.0
2	B	212	ILE	2.0
2	B	136	GLY	2.0
1	A	14	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	180	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TYS	H	63	16/17	0.86	0.18	34,37,39,41	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	B	2249	1/1	0.91	0.18	23,23,23,23	0
6	897	B	2250	33/33	0.91	0.12	9,13,32,34	0
4	NA	B	2248	1/1	0.95	0.10	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.