



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:33 PM UTC

PDB ID : 6UUH / pdb_00006uuu
Title : Crystal structure of broad and potent HIV-1 neutralizing antibody 438-B11
Authors : Kumar, S.; Wilson, I.A.
Deposited on : 2019-10-30
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	3% 86% 11% .
1	C	235	3% 83% 11% ..
2	B	215	% 91% 9%
2	D	215	4% 86% 13% .
3	E	66	3% 85% 6% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	66	5% 85% 6% 9%
4	G	8	62% 38%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B11 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	225	Total	C	N	O	S	0	0	0
			1706	1086	289	323	8			
1	A	228	Total	C	N	O	S	0	0	0
			1724	1095	292	329	8			

- Molecule 2 is a protein called B11 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	214	Total	C	N	O	S	0	0	0
			1643	1027	282	330	4			
2	B	215	Total	C	N	O	S	0	0	0
			1649	1030	283	331	5			

- Molecule 3 is a protein called Immunoglobulin G-binding protein G.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	60	Total	C	N	O	0	0	0
			459	289	72	98			
3	F	60	Total	C	N	O	0	0	0
			459	289	72	98			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	1	MET	-	initiating methionine	UNP P19909
E	62	HIS	-	expression tag	UNP P19909
E	63	HIS	-	expression tag	UNP P19909
E	64	HIS	-	expression tag	UNP P19909
E	65	HIS	-	expression tag	UNP P19909
E	66	HIS	-	expression tag	UNP P19909
F	1	MET	-	initiating methionine	UNP P19909
F	62	HIS	-	expression tag	UNP P19909

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	63	HIS	-	expression tag	UNP P19909
F	64	HIS	-	expression tag	UNP P19909
F	65	HIS	-	expression tag	UNP P19909
F	66	HIS	-	expression tag	UNP P19909

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	8	Total	C	N	O	0	0	0
			96	54	3	39			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: B11 Fab Heavy Chain




- Molecule 1: B11 Fab Heavy Chain




- Molecule 2: B11 Fab Light Chain




- Molecule 2: B11 Fab Light Chain






- Molecule 3: Immunoglobulin G-binding protein G



- Molecule 3: Immunoglobulin G-binding protein G



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.16Å 72.75Å 108.72Å 90.00° 101.17° 90.00°	Depositor
Resolution (Å)	48.22 – 2.70 48.22 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.4 (48.22-2.70) 96.3 (48.22-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.227 , 0.260 0.226 , 0.261	Depositor DCC
R_{free} test set	1455 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	57.4	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7750	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/1769	0.41	1/2410 (0.0%)
1	C	0.14	0/1751	0.41	1/2386 (0.0%)
2	B	0.13	0/1684	0.37	0/2287
2	D	0.17	0/1678	0.43	0/2279
3	E	0.23	0/466	0.30	0/635
3	F	0.22	0/466	0.31	0/635
All	All	0.16	0/7814	0.39	2/10632 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	100(B)	PRO	N-CA-CB	7.08	110.68	103.25
1	A	100(B)	PRO	N-CA-CB	7.03	110.63	103.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1724	0	1691	13	0
1	C	1706	0	1676	15	0
2	B	1649	0	1598	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1643	0	1593	22	0
3	E	459	0	447	2	0
3	F	459	0	447	2	0
4	G	96	0	82	0	0
5	D	14	0	13	0	0
All	All	7750	0	7547	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:61:ARG:HG2	2:D:61:ARG:HH11	1.30	0.97
1:C:85:ASP:OD1	1:C:85:ASP:N	2.19	0.74
2:D:61:ARG:NH1	2:D:82:ASP:OD2	2.23	0.68
2:D:37:GLN:HB2	2:D:47:LEU:HD11	1.78	0.66
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.80	0.61
2:D:39:ARG:NH2	2:D:42:GLN:OE1	2.36	0.59
3:E:13:ASN:HB2	3:E:60:THR:HG22	1.84	0.58
2:D:61:ARG:HG3	2:D:76:THR:O	2.05	0.57
2:D:6:GLN:HG3	2:D:100:PRO:HD2	1.87	0.56
1:C:144:ASP:OD1	1:C:171:GLN:NE2	2.38	0.56
1:A:148:GLU:HG3	1:A:149:PRO:HA	1.88	0.56
1:C:12:ARG:HG3	1:C:18:VAL:HB	1.90	0.54
2:D:61:ARG:HG2	2:D:61:ARG:NH1	2.08	0.54
3:F:13:ASN:HB2	3:F:60:THR:HG22	1.91	0.53
1:C:34:ILE:HD13	1:C:94:ARG:HB3	1.91	0.51
2:D:183:LYS:O	2:D:187:GLU:HG2	2.10	0.51
1:C:1:GLN:HG2	1:C:96:ARG:NH2	2.24	0.51
2:B:54:ARG:HH21	2:B:63:THR:HG22	1.76	0.51
2:D:61:ARG:HH11	2:D:61:ARG:CG	2.09	0.50
2:D:54:ARG:HH21	2:D:63:THR:HG22	1.77	0.49
1:C:171:GLN:HG2	1:C:175:LEU:O	2.13	0.49
3:E:15:LYS:HG2	3:E:61:GLU:OE1	2.12	0.49
1:C:13:LYS:HD3	1:C:113:ALA:HA	1.95	0.48
1:A:168:ALA:HB2	1:A:178:LEU:HD23	1.96	0.48
2:D:61:ARG:NH1	2:D:61:ARG:CG	2.73	0.48
1:A:168:ALA:HA	1:A:178:LEU:HB3	1.95	0.47
2:D:76:THR:HG23	2:D:77:ARG:HD2	1.97	0.47
2:D:187:GLU:O	2:D:211:ARG:NH1	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:108:ARG:NH1	2:D:109:THR:HG23	2.31	0.46
1:C:168:ALA:HA	1:C:178:LEU:HB3	1.97	0.46
1:A:27:GLY:HA2	1:A:100:MET:HA	1.97	0.46
2:D:113:PRO:HB3	2:D:139:PHE:HB3	1.98	0.45
2:B:6:GLN:HG3	2:B:100:PRO:HD2	1.99	0.45
2:B:50:ALA:O	2:B:52:SER:N	2.45	0.45
1:A:162:GLY:O	1:A:182:VAL:HA	2.16	0.45
2:B:21:LEU:HD23	2:B:102:THR:HB	1.98	0.45
1:C:32:PHE:CZ	1:C:97:GLY:HA3	2.51	0.44
2:B:210:ASN:HB3	2:B:214:CYS:HB3	1.98	0.44
2:D:78:LEU:HD12	2:D:83:PHE:CE2	2.52	0.44
1:A:47:TRP:CG	2:B:96:PHE:HB2	2.53	0.44
3:F:48:TRP:HA	3:F:58:THR:O	2.18	0.44
1:C:119:PRO:HB3	1:C:145:TYR:HB3	1.99	0.43
2:D:61:ARG:CG	2:D:76:THR:O	2.66	0.43
2:D:189:HIS:O	2:D:211:ARG:NH1	2.45	0.43
1:A:148:GLU:HG3	1:A:149:PRO:CA	2.48	0.43
1:A:13:LYS:HD3	1:A:113:ALA:HA	2.00	0.42
2:D:14:SER:HB2	2:D:17:GLU:HG3	2.00	0.42
2:B:187:GLU:O	2:B:211:ARG:NH2	2.51	0.42
1:A:171:GLN:HG2	1:A:175:LEU:O	2.19	0.42
1:C:168:ALA:HB2	1:C:178:LEU:HD23	2.01	0.42
1:A:108:LYS:HG2	1:A:110:ILE:HD11	2.01	0.42
1:A:20:ILE:HD12	1:A:80:LEU:HD22	2.01	0.42
1:C:148:GLU:HG3	1:C:149:PRO:HA	2.01	0.42
2:D:5:THR:OG1	2:D:24:ARG:HB3	2.19	0.42
2:D:54:ARG:HG2	2:D:58:ILE:HB	2.02	0.41
2:B:83:PHE:CZ	2:B:106:VAL:HG22	2.56	0.41
2:D:50:ALA:O	2:D:52:SER:N	2.48	0.41
1:A:93:THR:HB	1:A:100(M):SER:HB3	2.03	0.41
2:B:183:LYS:O	2:B:187:GLU:HG2	2.21	0.41
1:C:148:GLU:HG3	1:C:149:PRO:CA	2.50	0.41
2:B:10:THR:HG23	2:B:103:GLN:HG2	2.03	0.41
1:C:146:PHE:HA	1:C:147:PRO:HA	1.84	0.40
1:C:162:GLY:O	1:C:182:VAL:HA	2.20	0.40
1:A:119:PRO:HB3	1:A:145:TYR:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/235 (95%)	216 (96%)	7 (3%)	1 (0%)	30	54
1	C	221/235 (94%)	213 (96%)	7 (3%)	1 (0%)	24	48
2	B	213/215 (99%)	208 (98%)	5 (2%)	0	100	100
2	D	212/215 (99%)	205 (97%)	7 (3%)	0	100	100
3	E	58/66 (88%)	57 (98%)	1 (2%)	0	100	100
3	F	58/66 (88%)	57 (98%)	1 (2%)	0	100	100
All	All	986/1032 (96%)	956 (97%)	28 (3%)	2 (0%)	43	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	100(B)	PRO
1	A	100(B)	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/201 (96%)	189 (98%)	4 (2%)	47	75
1	C	190/201 (94%)	185 (97%)	5 (3%)	40	70
2	B	185/185 (100%)	185 (100%)	0	100	100
2	D	184/185 (100%)	180 (98%)	4 (2%)	45	74
3	E	49/55 (89%)	49 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	F	49/55 (89%)	49 (100%)	0	100	100
All	All	850/882 (96%)	837 (98%)	13 (2%)	57	81

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	85	ASP
1	C	98	CYS
1	C	115	SER
1	C	199	ASN
1	C	207	VAL
2	D	3	VAL
2	D	61	ARG
2	D	77	ARG
2	D	109	THR
1	A	98	CYS
1	A	150	VAL
1	A	199	ASN
1	A	207	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	1	2,4	14,14,15	0.34	0	17,19,21	0.58	0
4	NAG	G	2	4	14,14,15	0.18	0	17,19,21	0.52	0
4	BMA	G	3	4	11,11,12	0.45	0	15,15,17	0.74	0
4	MAN	G	4	4	11,11,12	0.53	0	15,15,17	1.03	1 (6%)
4	MAN	G	5	4	11,11,12	0.99	1 (9%)	15,15,17	1.20	2 (13%)
4	MAN	G	6	4	11,11,12	0.77	1 (9%)	15,15,17	0.95	2 (13%)
4	NAG	G	7	4	14,14,15	0.50	0	17,19,21	0.42	0
4	FUC	G	8	4	10,10,11	0.93	0	14,14,16	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	BMA	G	3	4	-	2/2/19/22	0/1/1/1
4	MAN	G	4	4	-	2/2/19/22	0/1/1/1
4	MAN	G	5	4	-	2/2/19/22	0/1/1/1
4	MAN	G	6	4	-	1/2/19/22	0/1/1/1
4	NAG	G	7	4	-	2/6/23/26	0/1/1/1
4	FUC	G	8	4	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	5	MAN	C1-C2	2.53	1.58	1.52
4	G	6	MAN	C1-C2	2.03	1.57	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	5	MAN	C1-O5-C5	3.16	116.43	112.19
4	G	4	MAN	C1-O5-C5	3.03	116.25	112.19
4	G	5	MAN	O2-C2-C3	-2.41	105.17	110.15
4	G	6	MAN	C1-O5-C5	2.34	115.33	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	6	MAN	O2-C2-C3	-2.02	105.96	110.15

There are no chirality outliers.


All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	1	NAG	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	G	5	MAN	C4-C5-C6-O6
4	G	3	BMA	O5-C5-C6-O6
4	G	4	MAN	O5-C5-C6-O6
4	G	5	MAN	O5-C5-C6-O6
4	G	4	MAN	C4-C5-C6-O6
4	G	3	BMA	C4-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6
4	G	7	NAG	C4-C5-C6-O6
4	G	7	NAG	O5-C5-C6-O6
4	G	6	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	D	301	2	14,14,15	0.37	0	17,19,21	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	301	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	301	NAG	C4-C5-C6-O6
5	D	301	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	228/235 (97%)	0.11	6 (2%)	57	54	29, 49, 76, 90	0
1	C	225/235 (95%)	0.26	6 (2%)	56	53	33, 56, 79, 92	0
2	B	215/215 (100%)	0.27	2 (0%)	81	80	34, 54, 87, 106	0
2	D	214/215 (99%)	0.56	8 (3%)	45	41	44, 68, 96, 105	0
3	E	60/66 (90%)	0.46	2 (3%)	49	45	40, 60, 80, 99	0
3	F	60/66 (90%)	0.72	3 (5%)	34	30	40, 64, 94, 117	0
All	All	1002/1032 (97%)	0.33	27 (2%)	56	53	29, 58, 87, 117	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	154	LEU	4.0
2	B	214	CYS	3.9
2	B	60	ASP	3.4
3	F	3	PRO	3.3
3	F	2	THR	3.3
1	A	128	SER	3.1
1	C	126	PRO	3.0
2	D	212	GLY	2.9
3	F	26	VAL	2.9
3	E	3	PRO	2.8
1	C	100(B)	PRO	2.7
2	D	191	VAL	2.6
3	E	2	THR	2.6
1	A	64	HIS	2.5
1	A	113	ALA	2.5
1	C	100(A)	CYS	2.4
2	D	193	ALA	2.4
1	C	214	LYS	2.4
2	D	184	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	100	MET	2.2
2	D	78	LEU	2.2
1	A	100(B)	PRO	2.1
1	A	133	GLY	2.1
1	C	75	MET	2.1
2	D	202	SER	2.0
1	A	42	GLY	2.0
2	D	13	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	G	7	14/15	0.61	0.23	56,58,59,59	0
4	FUC	G	8	10/11	0.65	0.18	130,131,132,132	0
4	MAN	G	5	11/12	0.67	0.13	87,89,90,91	0
4	MAN	G	6	11/12	0.69	0.14	76,78,79,79	0
4	NAG	G	1	14/15	0.74	0.16	126,127,128,128	0
4	MAN	G	4	11/12	0.78	0.18	126,127,129,129	0
4	NAG	G	2	14/15	0.79	0.17	127,129,132,132	0
4	BMA	G	3	11/12	0.84	0.18	125,126,129,129	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	D	301	14/15	0.67	0.18	136,137,138,139	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAG D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.