



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:12 PM UTC

PDB ID : 2UVF / pdb_00002uvf
Title : Structure of Yersinia enterocolitica Family 28 Exopolysaccharide Synthase in Complex with Digalacturonic Acid
Authors : Abbott, D.W.; Boraston, A.B.
Deposited on : 2007-03-09
Resolution : 2.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	608	 9% 67% 22% •• 6%
1	B	608	 2% 75% 16% •• 7%
2	C	2	 50% 50%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 9493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXOPOLYGALACTURONASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	571	Total	C	N	O	S	0	4	0
			4459	2795	781	868	15			
1	B	567	Total	C	N	O	S	0	0	0
			4392	2760	766	851	15			

- Molecule 2 is an oligosaccharide called alpha-D-galactopyranuronic acid-(1-4)-alpha-D-galactopyranuronic acid.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	Total	C	O	0	0	0
			25	12	13			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total Ni 4 4	0	0
4	B	6	Total Ni 6 6	0	0

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	4	2	2	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	B	1	7	4	3	0	0
6	B	1	7	4	3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	184	184	184	0	0
7	B	303	303	303	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EXOPOLYGALACTURONASE



• Molecule 1: EXOPOLYGALACTURONASE





- Molecule 2: alpha-D-galactopyranuronic acid-(1-4)-alpha-D-galactopyranuronic acid

Chain C: 50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.02Å 79.66Å 98.49Å 90.00° 103.90° 90.00°	Depositor
Resolution (Å)	19.72 – 2.10 19.72 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.1 (19.72-2.10) 98.9 (19.72-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.09Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.232 , 0.284 0.225 , 0.271	Depositor DCC
R_{free} test set	3976 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9493	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NI, ACT, SO4, ADA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	1/4552 (0.0%)	0.99	17/6179 (0.3%)
1	B	0.91	5/4482 (0.1%)	1.03	9/6083 (0.1%)
All	All	0.85	6/9034 (0.1%)	1.01	26/12262 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	1	5
All	All	1	10

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	357	ILE	C-O	-6.20	1.17	1.24
1	B	599	VAL	CA-CB	5.73	1.61	1.54
1	B	238	ILE	CA-CB	5.37	1.60	1.54
1	A	601	VAL	N-CA	5.21	1.52	1.46
1	B	294	VAL	CA-CB	5.17	1.59	1.54

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	604	LYS	CA-C-N	12.47	135.42	119.84
1	B	604	LYS	C-N-CA	12.47	135.42	119.84
1	A	604	LYS	CA-C-N	11.88	134.69	119.84
1	A	604	LYS	C-N-CA	11.88	134.69	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	446	ALA	N-CA-C	-11.81	95.29	110.53

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	442	SER	CA

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	440	THR	Peptide
1	A	502	ALA	Peptide
1	A	600	LYS	Peptide
1	A	602	ASP	Peptide
1	A	604	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4459	0	4340	114	0
1	B	4392	0	4289	74	0
2	C	25	0	15	1	0
3	A	30	0	0	0	0
3	B	60	0	0	0	0
4	A	4	0	0	0	0
4	B	6	0	0	0	0
5	B	16	0	12	1	0
6	B	14	0	20	2	0
7	A	184	0	0	3	0
7	B	303	0	0	9	0
All	All	9493	0	8676	188	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:ALA:HB3	1:A:503:ASP:HA	1.30	1.10
1:B:422:LYS:HD3	7:B:2226:HOH:O	1.63	0.98
1:B:601:VAL:O	1:B:603:GLY:HA2	1.67	0.93
1:A:502:ALA:CB	1:A:503:ASP:HA	1.98	0.92
1:B:113:GLN:H	1:B:428:ASN:HD21	1.10	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	573/608 (94%)	522 (91%)	41 (7%)	10 (2%)	7 3
1	B	563/608 (93%)	527 (94%)	31 (6%)	5 (1%)	14 10
All	All	1136/1216 (93%)	1049 (92%)	72 (6%)	15 (1%)	10 6

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	600	LYS
1	A	601	VAL
1	B	605	PRO
1	A	503	ASP
1	A	507	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	481/507 (95%)	449 (93%)	32 (7%)	15	12
1	B	472/507 (93%)	451 (96%)	21 (4%)	25	26
All	All	953/1014 (94%)	900 (94%)	53 (6%)	19	18

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	574	ASP
1	B	121	LYS
1	B	548	LYS
1	A	599	VAL
1	B	36	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	418	GLN
1	B	563	ASN
1	B	428	ASN
1	B	531	ASN
1	A	428	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADA	C	1	2	13,13,13	1.47	2 (15%)	18,19,19	1.68	3 (16%)
2	ADA	C	2	2	12,12,13	1.28	0	14,17,19	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADA	C	1	2	-	0/4/24/24	0/1/1/1
2	ADA	C	2	2	-	0/4/21/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	ADA	O5-C5	3.32	1.49	1.43
2	C	1	ADA	O1-C1	2.11	1.46	1.39

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	ADA	C1-O5-C5	3.00	116.64	112.22
2	C	1	ADA	O5-C1-C2	-2.92	105.16	110.30
2	C	1	ADA	C4-C3-C2	-2.86	105.81	110.83

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	ADA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 34 ligands modelled in this entry, 10 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	1615	-	4,4,4	0.28	0	6,6,6	0.19	0
5	ACT	B	1628	-	3,3,3	0.85	0	3,3,3	0.74	0
3	SO4	A	1611	-	4,4,4	0.25	0	6,6,6	0.29	0
3	SO4	A	1612	-	4,4,4	0.23	0	6,6,6	0.15	0
3	SO4	B	1611	-	4,4,4	0.26	0	6,6,6	0.10	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	1610	-	4,4,4	0.22	0	6,6,6	0.28	0
3	SO4	B	1609	-	4,4,4	0.22	0	6,6,6	0.10	0
3	SO4	B	1616	-	4,4,4	0.22	0	6,6,6	0.15	0
3	SO4	B	1614	-	4,4,4	0.25	0	6,6,6	0.29	0
3	SO4	B	1612	-	4,4,4	0.22	0	6,6,6	0.15	0
3	SO4	B	1618	-	4,4,4	0.26	0	6,6,6	0.30	0
5	ACT	B	1627	-	3,3,3	0.76	0	3,3,3	0.89	0
3	SO4	A	1608	-	4,4,4	0.29	0	6,6,6	0.15	0
5	ACT	B	1629	-	3,3,3	0.73	0	3,3,3	1.04	0
3	SO4	B	1617	-	4,4,4	0.25	0	6,6,6	0.22	0
3	SO4	A	1610	-	4,4,4	0.28	0	6,6,6	0.31	0
3	SO4	A	1609	-	4,4,4	0.21	0	6,6,6	0.20	0
6	PEG	B	1630	-	6,6,6	0.35	0	5,5,5	0.94	0
6	PEG	B	1631	-	6,6,6	0.57	0	5,5,5	0.16	0
3	SO4	B	1619	-	4,4,4	0.56	0	6,6,6	0.60	0
3	SO4	B	1608	-	4,4,4	0.25	0	6,6,6	0.19	0
5	ACT	B	1626	-	3,3,3	0.87	0	3,3,3	0.99	0
3	SO4	B	1613	-	4,4,4	0.26	0	6,6,6	0.20	0
3	SO4	A	1607	-	4,4,4	0.30	0	6,6,6	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	1630	-	-	3/4/4/4	-
6	PEG	B	1631	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	1630	PEG	O1-C1-C2-O2
6	B	1631	PEG	O2-C3-C4-O4
6	B	1630	PEG	O2-C3-C4-O4
6	B	1631	PEG	C1-C2-O2-C3
6	B	1631	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1630	PEG	1	0
6	B	1631	PEG	1	0
5	B	1626	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	571/608 (93%)	0.61	54 (9%) 14 14	8, 39, 64, 86	6 (1%)
1	B	567/608 (93%)	-0.04	12 (2%) 63 66	7, 27, 42, 58	1 (0%)
All	All	1138/1216 (93%)	0.29	66 (5%) 29 30	7, 32, 56, 86	7 (0%)

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	502	ALA	8.8
1	B	508	ILE	6.6
1	A	546	ALA	6.6
1	A	545	THR	6.3
1	A	606	VAL	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ADA	C	1	13/13	0.79	0.12	40,47,51,52	0
2	ADA	C	2	12/13	0.84	0.12	43,46,49,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	1613	5/5	0.50	0.16	87,87,87,87	0
3	SO4	A	1612	5/5	0.65	0.16	104,104,105,105	0
5	ACT	B	1626	4/4	0.66	0.20	44,44,44,44	0
3	SO4	B	1618	5/5	0.69	0.16	73,73,74,74	0
3	SO4	B	1611	5/5	0.69	0.12	112,112,112,112	0
5	ACT	B	1628	4/4	0.72	0.15	42,42,42,43	0
5	ACT	B	1627	4/4	0.74	0.19	48,48,48,49	0
3	SO4	A	1610	5/5	0.76	0.20	82,82,83,83	0
4	NI	B	1622	1/1	0.79	0.13	93,93,93,93	0
3	SO4	A	1611	5/5	0.82	0.20	73,73,74,74	0
3	SO4	B	1615	5/5	0.83	0.13	97,97,97,98	0
3	SO4	B	1612	5/5	0.83	0.15	97,97,97,98	0
3	SO4	B	1614	5/5	0.85	0.11	77,78,78,78	0
3	SO4	B	1619	5/5	0.85	0.33	40,41,42,42	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PEG	B	1631	7/7	0.85	0.10	46,48,50,50	0
4	NI	B	1624	1/1	0.86	0.13	85,85,85,85	0
3	SO4	B	1608	5/5	0.88	0.14	75,75,76,76	0
6	PEG	B	1630	7/7	0.88	0.16	33,37,43,43	0
3	SO4	B	1616	5/5	0.88	0.16	71,71,72,73	0
3	SO4	B	1617	5/5	0.89	0.11	60,60,60,61	0
5	ACT	B	1629	4/4	0.90	0.17	36,36,36,36	0
3	SO4	A	1607	5/5	0.90	0.13	52,54,56,56	0
3	SO4	A	1608	5/5	0.90	0.10	61,62,63,63	0
4	NI	B	1625	1/1	0.92	0.14	75,75,75,75	0
3	SO4	A	1609	5/5	0.94	0.09	53,53,55,55	0
3	SO4	B	1609	5/5	0.94	0.09	47,48,49,49	0
4	NI	A	1616	1/1	0.95	0.16	75,75,75,75	0
3	SO4	B	1610	5/5	0.95	0.11	41,42,42,46	0
4	NI	B	1621	1/1	0.96	0.06	50,50,50,50	0
4	NI	A	1615	1/1	0.96	0.20	49,49,49,49	0
4	NI	B	1623	1/1	0.98	0.15	40,40,40,40	0
4	NI	A	1613	1/1	0.98	0.13	32,32,32,32	0
4	NI	B	1620	1/1	0.98	0.08	28,28,28,28	0
4	NI	A	1614	1/1	0.99	0.07	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.