



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:39 AM UTC

PDB ID : 3UV7 / pdb_00003uv7
Title : Ec_IspH in complex with buta-2,3-dienyl diphosphate (1300)
Authors : Span, I.; Wang, K.; Wang, W.; Zhang, Y.; Bacher, A.; Eisenreich, W.; Schulz, C.; Oldfield, E.; Groll, M.
Deposited on : 2011-11-29
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	 80% 12% 5%
1	B	324	 80% 13% . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2383	1487	427	459	10			
1	B	310	Total	C	N	O	S	0	0	0
			2390	1492	428	460	10			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	expression tag	UNP P62623
A	-6	HIS	-	expression tag	UNP P62623
A	-5	HIS	-	expression tag	UNP P62623
A	-4	HIS	-	expression tag	UNP P62623
A	-3	HIS	-	expression tag	UNP P62623
A	-2	HIS	-	expression tag	UNP P62623
A	-1	GLY	-	expression tag	UNP P62623
A	0	SER	-	expression tag	UNP P62623
B	-7	HIS	-	expression tag	UNP P62623
B	-6	HIS	-	expression tag	UNP P62623
B	-5	HIS	-	expression tag	UNP P62623
B	-4	HIS	-	expression tag	UNP P62623
B	-3	HIS	-	expression tag	UNP P62623
B	-2	HIS	-	expression tag	UNP P62623
B	-1	GLY	-	expression tag	UNP P62623
B	0	SER	-	expression tag	UNP P62623

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is buta-2,3-dien-1-yl trihydrogen diphosphate (CCD ID: 0CN) (formula: $C_4H_8O_7P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			13	4	7	2		
3	B	1	Total	C	O	P	0	0
			13	4	7	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	236	Total 236	O 236	0	0
4	B	239	Total 239	O 239	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase

Chain A: 



- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.40Å 80.51Å 111.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 50.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-1.60) 99.7 (50.00-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.12 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.160 , 0.210 0.160 , 0.209	Depositor DCC
R_{free} test set	4204 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	15.5	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5290	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.34 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7890e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, OCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.58	23/2418 (1.0%)	1.46	7/3273 (0.2%)
1	B	1.52	18/2425 (0.7%)	1.48	17/3283 (0.5%)
All	All	1.55	41/4843 (0.8%)	1.47	24/6556 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	230	ARG	CD-NE	-10.08	1.32	1.46
1	A	121	HIS	CE1-NE2	8.54	1.41	1.32
1	A	124	HIS	CE1-NE2	8.29	1.40	1.32
1	B	225	SER	N-CA	7.73	1.55	1.46
1	B	230	ARG	CD-NE	-7.40	1.35	1.46
1	B	37	HIS	CG-CD2	7.25	1.43	1.35
1	A	124	HIS	CG-CD2	7.12	1.43	1.35
1	B	124	HIS	CG-CD2	7.06	1.43	1.35
1	A	103	HIS	ND1-CE1	6.56	1.39	1.32
1	B	57	ILE	N-CA	6.44	1.53	1.46
1	B	237	ARG	CD-NE	-6.37	1.37	1.46
1	A	264	VAL	C-O	6.23	1.30	1.24
1	A	103	HIS	CE1-NE2	6.11	1.38	1.32
1	B	41	HIS	CG-CD2	6.11	1.42	1.35
1	B	41	HIS	CE1-NE2	6.09	1.38	1.32
1	A	230	ARG	CZ-NH2	6.07	1.41	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	184	ARG	CZ-NH2	6.07	1.41	1.33
1	A	164	MET	N-CA	6.03	1.53	1.45
1	B	121	HIS	CG-CD2	6.03	1.42	1.35
1	A	75	GLY	N-CA	5.99	1.51	1.45
1	A	262	VAL	N-CA	5.99	1.53	1.46
1	B	41	HIS	ND1-CE1	5.86	1.38	1.32
1	A	303	GLU	N-CA	5.83	1.53	1.45
1	A	121	HIS	CG-CD2	5.79	1.42	1.35
1	A	276	GLN	CD-OE1	5.73	1.34	1.23
1	A	2	GLN	CA-C	5.68	1.60	1.52
1	A	95	THR	C-O	5.59	1.30	1.23
1	A	121	HIS	ND1-CE1	5.50	1.38	1.32
1	A	41	HIS	CE1-NE2	5.44	1.38	1.32
1	B	262	VAL	N-CA	5.40	1.52	1.46
1	B	121	HIS	CE1-NE2	5.38	1.38	1.32
1	A	121	HIS	CB-CG	5.38	1.57	1.50
1	A	225	SER	N-CA	5.36	1.52	1.46
1	A	176	ASP	CA-C	5.33	1.59	1.52
1	B	160	LYS	C-O	-5.32	1.17	1.23
1	B	40	VAL	C-O	5.28	1.29	1.23
1	B	230	ARG	CZ-NH2	5.26	1.40	1.33
1	B	144	VAL	C-O	5.17	1.29	1.24
1	B	3	ILE	N-CA	5.17	1.52	1.46
1	A	7	ASN	CB-CG	-5.13	1.39	1.52
1	A	101	LYS	N-CA	5.04	1.52	1.46

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	230	ARG	NE-CZ-NH1	-21.34	100.16	121.50
1	B	230	ARG	NE-CZ-NH2	16.82	134.34	119.20
1	B	230	ARG	NE-CZ-NH1	-16.82	104.68	121.50
1	A	230	ARG	NE-CZ-NH2	15.84	133.45	119.20
1	B	230	ARG	CD-NE-CZ	10.12	138.57	124.40
1	A	230	ARG	CD-NE-CZ	9.59	137.82	124.40
1	B	230	ARG	CG-CD-NE	-9.15	91.87	112.00
1	B	237	ARG	NE-CZ-NH2	9.10	127.39	119.20
1	B	237	ARG	CB-CG-CD	8.64	131.18	111.30
1	B	52	ARG	NE-CZ-NH2	7.67	126.10	119.20
1	B	184	ARG	NE-CZ-NH1	-7.58	113.92	121.50
1	B	285	LEU	N-CA-C	-7.29	103.96	112.92
1	B	281	ARG	NE-CZ-NH1	6.94	128.44	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	230	ARG	CG-CD-NE	-6.36	98.01	112.00
1	A	7	ASN	CA-CB-CG	-6.00	106.60	112.60
1	B	203	ARG	CA-CB-CG	-5.95	102.20	114.10
1	A	141	MET	CG-SD-CE	5.85	113.77	100.90
1	B	281	ARG	CD-NE-CZ	5.75	132.46	124.40
1	B	281	ARG	NE-CZ-NH2	-5.73	114.04	119.20
1	B	237	ARG	NE-CZ-NH1	-5.73	115.77	121.50
1	A	233	GLU	CB-CG-CD	-5.67	102.96	112.60
1	B	184	ARG	NE-CZ-NH2	5.52	124.17	119.20
1	B	155	VAL	N-CA-CB	-5.35	103.48	112.47
1	B	52	ARG	NE-CZ-NH1	-5.04	116.46	121.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	7	ASN	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2383	0	2395	30	1
1	B	2390	0	2404	31	1
2	A	8	0	0	0	0
2	B	8	0	0	0	0
3	A	13	0	5	3	0
3	B	13	0	6	1	0
4	A	236	0	0	9	0
4	B	239	0	0	5	0
All	All	5290	0	4810	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:HIS:HD2	4:A:566:HOH:O	0.85	1.18
1:A:36:ARG:HH21	1:A:36:ARG:HG2	1.19	1.02
1:A:36:ARG:HG2	1:A:36:ARG:NH2	1.78	0.87
1:B:108:ARG:NH2	1:B:108:ARG:HB2	1.94	0.83
1:B:42:ASN:HD22	1:B:45:VAL:H	1.26	0.80
1:A:121:HIS:CD2	4:A:566:HOH:O	1.75	0.79
1:B:108:ARG:HB2	1:B:108:ARG:CZ	2.17	0.75
1:B:112:ARG:HG3	1:B:112:ARG:HH21	1.51	0.74
1:B:51:GLU:OE2	1:B:52:ARG:HD2	1.88	0.73
1:A:42:ASN:HD22	1:A:45:VAL:H	1.34	0.72
1:A:103:HIS:HD2	4:A:478:HOH:O	1.71	0.72
1:B:233:GLU:OE2	4:B:451:HOH:O	2.09	0.71
1:B:68:ILE:HD13	1:B:90:THR:HB	1.74	0.69
1:A:73:ALA:CB	3:A:318:OCN:H11	2.23	0.68
1:B:103:HIS:HD2	4:B:424:HOH:O	1.77	0.68
1:A:36:ARG:HH21	1:A:36:ARG:CG	1.95	0.67
1:B:310:VAL:HG12	1:B:310:VAL:O	1.93	0.66
1:B:272:ASP:OD2	4:B:580:HOH:O	2.14	0.65
1:A:4:LEU:HD22	1:A:261:CYS:SG	2.37	0.65
1:B:112:ARG:HB3	1:B:114:GLU:HG3	1.80	0.64
1:A:73:ALA:HB3	3:A:318:OCN:H11	1.83	0.60
1:A:52:ARG:NH1	4:A:494:HOH:O	2.34	0.60
1:A:66:GLY:H	1:A:88:ASP:HB3	1.66	0.59
1:B:136:ASN:ND2	1:B:138:GLU:H	2.00	0.59
1:A:103:HIS:HE1	4:A:324:HOH:O	1.88	0.57
1:A:136:ASN:HD22	1:A:136:ASN:C	2.15	0.55
1:B:108:ARG:CZ	1:B:108:ARG:CB	2.85	0.53
1:B:74:HIS:CD2	3:B:318:OCN:H11	2.43	0.53
1:B:112:ARG:CB	1:B:114:GLU:HG3	2.39	0.53
1:B:161:LEU:HB3	1:B:188:ILE:HG13	1.94	0.50
1:B:299:ASN:HD22	1:B:299:ASN:C	2.20	0.49
1:B:136:ASN:HD22	1:B:138:GLU:H	1.60	0.49
1:B:103:HIS:HE1	4:B:327:HOH:O	1.97	0.48
1:B:310:VAL:O	1:B:310:VAL:CG1	2.61	0.48
1:B:136:ASN:HD22	1:B:136:ASN:C	2.22	0.47
1:A:136:ASN:ND2	1:A:138:GLU:H	2.11	0.47
1:A:73:ALA:HB1	3:A:318:OCN:H11	1.95	0.47
1:A:1:MET:N	4:A:471:HOH:O	2.49	0.46
1:A:236:GLN:HG3	1:A:242:ALA:HB3	1.97	0.46
1:A:36:ARG:HB3	4:A:441:HOH:O	2.16	0.46
1:B:119:ILE:HD12	1:B:165:THR:HG22	1.98	0.45
1:A:161:LEU:HD13	1:A:188:ILE:HD12	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:VAL:HB	1:B:301:VAL:HB	1.98	0.45
1:A:60:ILE:CD1	1:A:69:LEU:HD11	2.48	0.44
1:A:59:GLN:HB3	4:A:479:HOH:O	2.17	0.44
1:B:42:ASN:ND2	1:B:45:VAL:H	2.04	0.43
1:B:230:ARG:HA	1:B:230:ARG:HD3	1.68	0.43
1:A:42:ASN:ND2	1:A:45:VAL:H	2.10	0.43
1:A:230:ARG:HD3	1:A:230:ARG:HA	1.83	0.43
1:A:7:ASN:OD1	4:A:564:HOH:O	2.21	0.42
1:A:5:LEU:HD12	1:A:264:VAL:HB	2.01	0.42
1:A:42:ASN:HD21	1:A:44:TYR:HB3	1.85	0.42
1:B:42:ASN:HD21	1:B:44:TYR:HB3	1.84	0.42
1:B:108:ARG:NH2	1:B:108:ARG:CB	2.75	0.42
1:B:120:GLY:O	1:B:145:GLU:HG2	2.20	0.42
1:A:136:ASN:HD22	1:A:138:GLU:H	1.68	0.41
1:B:136:ASN:HD21	1:B:138:GLU:HB2	1.86	0.41
1:B:230:ARG:HD3	4:B:451:HOH:O	2.21	0.40
1:A:108:ARG:CZ	1:A:108:ARG:HB2	2.51	0.40
1:B:8:PRO:HG2	1:B:202:ASN:HB3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:GLU:OE1	1:B:52:ARG:NH1[4_545]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/324 (95%)	304 (99%)	3 (1%)	0	100	100
1	B	308/324 (95%)	305 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	615/648 (95%)	609 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/271 (95%)	245 (95%)	12 (5%)	23	6
1	B	258/271 (95%)	242 (94%)	16 (6%)	16	3
All	All	515/542 (95%)	487 (95%)	28 (5%)	20	4

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	5	LEU
1	A	57	ILE
1	A	69	LEU
1	A	138	GLU
1	A	153	LEU
1	A	161	LEU
1	A	187	LYS
1	A	230	ARG
1	A	258	GLU
1	A	294	GLU
1	A	306	LYS
1	B	5	LEU
1	B	108	ARG
1	B	112	ARG
1	B	138	GLU
1	B	148	ASP
1	B	153	LEU
1	B	155	VAL
1	B	159	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	161	LEU
1	B	230	ARG
1	B	253	GLU
1	B	273	ILE
1	B	281	ARG
1	B	285	LEU
1	B	294	GLU
1	B	299	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	103	HIS
1	A	121	HIS
1	A	133	GLN
1	A	136	ASN
1	A	213	GLN
1	A	277	ASN
1	A	283	GLN
1	B	42	ASN
1	B	103	HIS
1	B	121	HIS
1	B	133	GLN
1	B	136	ASN
1	B	213	GLN
1	B	283	GLN
1	B	299	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OCN	B	318	-	11,12,12	2.27	5 (45%)	13,17,17	1.45	3 (23%)
2	SF4	B	317	1,4	0,12,12	-	-	-		
2	SF4	A	317	1,4	0,12,12	-	-	-		
3	OCN	A	318	-	11,12,12	2.09	5 (45%)	13,17,17	1.49	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OCN	B	318	-	-	1/10/12/12	-
3	OCN	A	318	-	-	1/10/12/12	-
2	SF4	B	317	1,4	-	-	0/6/5/5
2	SF4	A	317	1,4	-	-	0/6/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	318	OCN	C30-C27	4.75	1.39	1.29
3	A	318	OCN	C30-C27	4.10	1.38	1.29
3	A	318	OCN	P17-O20	-3.27	1.42	1.54
3	B	318	OCN	P13-O15	-2.72	1.42	1.55
3	B	318	OCN	P13-O14	-2.72	1.41	1.50
3	A	318	OCN	P17-O18	-2.64	1.45	1.54
3	A	318	OCN	C13-C30	2.48	1.40	1.30
3	B	318	OCN	P17-O19	-2.29	1.43	1.50
3	B	318	OCN	C13-C30	2.10	1.39	1.30
3	A	318	OCN	P13-O15	-2.01	1.46	1.55

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	318	0CN	O15-P13-O16	3.50	116.73	107.27
3	B	318	0CN	O15-P13-O16	3.10	115.64	107.27
3	B	318	0CN	C28-C27-C30	-2.56	118.12	124.49
3	B	318	0CN	O16-P13-O14	-2.34	103.67	110.70
3	A	318	0CN	C28-C27-C30	-2.17	119.09	124.49

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	318	0CN	C27-C28-O29-P13
3	B	318	0CN	C27-C28-O29-P13

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	318	0CN	1	0
3	A	318	0CN	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/324 (95%)	-0.28	1 (0%) 90 91	11, 20, 40, 65	0
1	B	310/324 (95%)	-0.22	1 (0%) 90 91	11, 21, 44, 69	0
All	All	619/648 (95%)	-0.25	2 (0%) 90 91	11, 20, 42, 69	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	VAL	3.5
1	A	27	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OCN	A	318	13/13	0.99	0.05	8,14,16,19	0
3	OCN	B	318	13/13	0.99	0.04	8,13,16,16	0
2	SF4	A	317	8/8	1.00	0.02	11,12,15,16	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SF4	B	317	8/8	1.00	0.02	11,12,15,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.