



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 07:22 PM UTC

PDB ID : 4UV8 / pdb_00004uv8
Title : LSD1(KDM1A)-CoREST in complex with 1-Benzyl-Tranylcypromine
Authors : Vianello, P.; Botrugno, O.; Cappa, A.; Ciossani, G.; Dessanti, P.; Mai, A.;
Mattevi, A.; Meroni, G.; Minucci, S.; Thaler, F.; Tortorici, M.; Trifiro, P.;
Valente, S.; Villa, M.; Varasi, M.; Mercurio, C.
Deposited on : 2014-08-05
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	872	5% 69% 7% 24%
2	B	482	6% 25% • 72%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSINE-SPECIFIC HISTONE DEMETHYLASE 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	666	5217	3324	906	967	20	0	0	0


There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	171	PRO	ALA	conflict	UNP O60341
A	?	-	ASP	deletion	UNP O60341
A	?	-	THR	deletion	UNP O60341
A	?	-	VAL	deletion	UNP O60341
A	?	-	LYS	deletion	UNP O60341

- Molecule 2 is a protein called REST COREPRESSOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	133	1076	676	194	203	3	0	0	0


- Molecule 3 is [[(2R,3S,4S)-5-[(4aS,10aS)-4a-[(1S)-3-azanylidene-1,4-diphenyl-butyl]-7,8-dimethyl-2,4-bis(oxidanylidene)-5,10a-dihydro-1H-benzo[g]pteridin-10-yl]-2,3,4-tris(oxidanyl)pen toxy]-oxidanyl-phosphoryl] [(2R,3S,4R,5R)-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methyl hydrogen phosphate (CCD ID: D69) (formula: C₄₃H₅₂N₁₀O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			70	43	10	15	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.85Å 179.21Å 235.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.96 – 2.80 58.96 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (58.96-2.80) 99.2 (58.96-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.7.0027	Depositor
R, R_{free}	0.226 , 0.237 0.224 , 0.235	Depositor DCC
R_{free} test set	1221 reflections (1.97%)	wwPDB-VP
Wilson B-factor (Å ²)	71.3	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6367	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D69

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/5331	0.77	0/7232
2	B	0.43	0/1091	0.79	0/1471
All	All	0.53	0/6422	0.77	0/8703

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5217	0	5252	38	0
2	B	1076	0	1091	12	0
3	A	70	0	48	3	0
4	A	4	0	0	0	0
All	All	6367	0	6391	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:900:D69:NAA	3:A:900:D69:HAJ	1.95	0.80
1:A:384:ARG:HB3	2:B:314:MET:CE	2.21	0.69
1:A:384:ARG:HB3	2:B:314:MET:HE1	1.78	0.66
1:A:548:SER:O	1:A:552:TRP:HB3	1.97	0.63
1:A:374:LYS:HE3	1:A:525:ASP:OD1	1.98	0.62
1:A:801:GLU:HG2	1:A:809:ALA:H	1.69	0.57
1:A:384:ARG:HB3	2:B:314:MET:HE3	1.89	0.55
1:A:374:LYS:C	1:A:376:GLU:H	2.15	0.54
1:A:435:VAL:HG13	2:B:349:ILE:HG13	1.89	0.53
1:A:463:LYS:O	1:A:467:GLU:HG2	2.09	0.52
1:A:441:LEU:HD23	2:B:356:ASN:HD22	1.75	0.51
3:A:900:D69:NAA	3:A:900:D69:CAJ	2.67	0.50
1:A:442:LYS:HE3	2:B:355:THR:HG21	1.94	0.49
1:A:801:GLU:HG2	1:A:809:ALA:N	2.29	0.47
1:A:419:GLN:HE22	2:B:315:PHE:H	1.61	0.47
1:A:776:MET:HE2	1:A:802:HIS:HD2	1.78	0.47
1:A:601:GLU:HA	1:A:616:TYR:O	2.14	0.46
1:A:662:VAL:HB	1:A:705:ALA:HB3	1.98	0.46
1:A:716:GLU:HG2	1:A:750:ARG:HG2	1.95	0.46
1:A:661:LYS:HD3	1:A:704:LEU:HD21	1.97	0.46
1:A:364:GLU:HA	1:A:681:VAL:HB	1.98	0.46
1:A:720:ASP:O	1:A:724:VAL:HG23	2.16	0.46
1:A:435:VAL:O	1:A:439:GLU:HB2	2.15	0.45
1:A:435:VAL:CG1	2:B:349:ILE:HG13	2.47	0.44
1:A:566:THR:HG21	1:A:697:LEU:HD13	1.99	0.44
1:A:592:GLN:HB3	1:A:603:ILE:HD12	2.00	0.44
1:A:209:VAL:HG13	1:A:242:TYR:HD1	1.83	0.43
1:A:801:GLU:CG	1:A:809:ALA:HA	2.47	0.43
1:A:353:LEU:HB3	1:A:565:LEU:HD22	2.00	0.43
1:A:331:ALA:HB2	3:A:900:D69:H10	2.01	0.42
1:A:485:ARG:HG3	2:B:407:ASP:HB2	2.01	0.42
1:A:501:GLN:O	1:A:505:GLU:HB2	2.20	0.41
1:A:270:ILE:O	1:A:272:PRO:HD3	2.21	0.41
1:A:761:TYR:CD1	1:A:809:ALA:HB1	2.56	0.41
1:A:180:GLN:HA	1:A:339:GLY:HA2	2.02	0.41
2:B:317:SER:O	2:B:321:VAL:HG23	2.21	0.40
1:A:495:ASP:OD1	2:B:371:ARG:NH2	2.55	0.40
1:A:484:HIS:CD2	2:B:372:LEU:HD13	2.56	0.40
1:A:374:LYS:C	1:A:376:GLU:N	2.78	0.40
1:A:541:ALA:O	1:A:657:GLY:HA3	2.22	0.40
1:A:625:LEU:HD22	1:A:629:VAL:HG11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/872 (76%)	643 (97%)	18 (3%)	3 (0%)	24	55
2	B	131/482 (27%)	124 (95%)	7 (5%)	0	100	100
All	All	795/1354 (59%)	767 (96%)	25 (3%)	3 (0%)	30	60

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	801	GLU
1	A	375	ASP
1	A	468	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	566/711 (80%)	553 (98%)	13 (2%)	44	78
2	B	117/395 (30%)	113 (97%)	4 (3%)	32	68
All	All	683/1106 (62%)	666 (98%)	17 (2%)	42	76

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	429	GLU
1	A	438	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	449	VAL
1	A	458	LEU
1	A	469	LYS
1	A	492	LYS
1	A	624	THR
1	A	633	GLN
1	A	645	GLU
1	A	659	LEU
1	A	684	THR
1	A	704	LEU
2	B	316	LEU
2	B	332	THR
2	B	337	GLN
2	B	349	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	GLN
1	A	380	GLN
1	A	410	GLN
1	A	419	GLN
1	A	438	GLN
1	A	459	HIS
1	A	484	HIS
1	A	742	GLN
2	B	318	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	D69	A	900	-	72,77,77	1.99	13 (18%)	90,116,116	1.53	14 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	D69	A	900	-	-	3/45/101/101	0/8/8/8

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	900	D69	C5X-C9A	8.34	1.49	1.40
3	A	900	D69	CAP-CAM	-6.87	1.41	1.51
3	A	900	D69	CAN-CAQ	-6.12	1.41	1.51
3	A	900	D69	C5A-C4A	4.70	1.47	1.39
3	A	900	D69	C8-C7	3.70	1.49	1.40
3	A	900	D69	C4X-N5	3.55	1.50	1.45
3	A	900	D69	C5A-C6A	2.87	1.49	1.41
3	A	900	D69	CAO-NAA	2.82	1.34	1.27
3	A	900	D69	C8A-N7A	2.47	1.36	1.31
3	A	900	D69	C5X-N5	-2.32	1.35	1.39
3	A	900	D69	P-O3P	2.26	1.61	1.59
3	A	900	D69	PA-O3P	2.21	1.61	1.59
3	A	900	D69	CAL-CAO	2.13	1.54	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	900	D69	C5A-C4A-N3A	-5.37	119.33	126.72
3	A	900	D69	N3A-C4A-N9A	4.30	134.48	127.17
3	A	900	D69	N3-C2-N1	3.82	119.89	116.06
3	A	900	D69	N3A-C2A-N1A	-3.80	122.83	128.58
3	A	900	D69	C2A-N3A-C4A	3.78	121.06	111.83
3	A	900	D69	C4-N3-C2	-3.51	120.80	126.04
3	A	900	D69	C4A-C5A-N7A	-3.40	106.69	110.58
3	A	900	D69	CAL-CAM-CAP	-3.40	108.12	112.27
3	A	900	D69	C4A-N9A-C8A	3.01	108.90	105.74
3	A	900	D69	C5A-N7A-C8A	2.49	107.37	103.45
3	A	900	D69	C6A-C5A-N7A	2.42	136.76	132.09
3	A	900	D69	C2A-N1A-C6A	2.19	122.33	118.73
3	A	900	D69	N9A-C8A-N7A	-2.18	110.85	113.94
3	A	900	D69	CAN-CAQ-CAK	-2.11	117.79	120.89

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	900	D69	C4-C4X-CAM-CAL
3	A	900	D69	CAL-CAM-CAP-CAH
3	A	900	D69	CAL-CAM-CAP-CAI

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	900	D69	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	666/872 (76%)	0.47	47 (7%)	22 16	44, 73, 109, 143	0
2	B	133/482 (27%)	1.16	28 (21%)	2 2	68, 104, 141, 148	0
All	All	799/1354 (59%)	0.59	75 (9%)	14 10	44, 80, 120, 148	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	836	LEU	9.7
1	A	171	PRO	7.0
2	B	376	ILE	6.0
1	A	377	MET	5.4
1	A	275	THR	4.8
1	A	833	MET	4.4
1	A	786	ILE	3.7
2	B	340	MET	3.6
2	B	337	GLN	3.5
1	A	510	GLU	3.5
1	A	269	ARG	3.5
1	A	467	GLU	3.5
2	B	312	LYS	3.5
1	A	835	THR	3.5
1	A	271	LYS	3.4
1	A	273	LEU	3.4
1	A	272	PRO	3.3
2	B	382	ARG	3.3
2	B	329	THR	3.2
1	A	358	GLN	3.0
1	A	834	TYR	2.9
1	A	403	ASN	2.9
2	B	308	ARG	2.9
1	A	506	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	375	VAL	2.9
1	A	684	THR	2.8
1	A	373	GLU	2.7
1	A	744	LYS	2.7
2	B	374	GLU	2.7
2	B	440	GLU	2.7
1	A	787	PRO	2.7
1	A	372	LYS	2.6
2	B	415	VAL	2.6
2	B	379	CYS	2.6
2	B	377	GLN	2.6
1	A	492	LYS	2.6
1	A	667	ASP	2.6
2	B	378	LYS	2.5
2	B	431	ASP	2.5
2	B	414	VAL	2.5
2	B	399	GLY	2.4
1	A	652	GLN	2.4
2	B	439	ALA	2.4
2	B	427	ARG	2.4
1	A	357	LYS	2.3
1	A	394	HIS	2.3
1	A	203	PRO	2.3
2	B	424	TYR	2.3
1	A	511	LEU	2.3
1	A	374	LYS	2.3
1	A	503	LYS	2.3
2	B	425	ARG	2.3
1	A	433	LYS	2.3
2	B	313	GLY	2.3
1	A	518	ASP	2.2
2	B	362	LYS	2.2
1	A	499	GLU	2.2
2	B	350	GLN	2.2
1	A	332	MET	2.2
2	B	386	GLU	2.2
1	A	204	GLN	2.2
1	A	519	VAL	2.2
2	B	402	PHE	2.1
1	A	668	ARG	2.1
1	A	237	GLN	2.1
2	B	383	TRP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	517	SER	2.1
1	A	698	TYR	2.1
1	A	270	ILE	2.1
2	B	438	GLU	2.1
1	A	172	SER	2.0
1	A	241	PRO	2.0
1	A	350	ASN	2.0
1	A	274	PRO	2.0
1	A	771	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	D69	A	900	70/70	0.97	0.09	44,61,88,90	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D69 A 900:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.