



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:20 PM UTC

PDB ID : 5UV8 / pdb_00005uv8
Title : Interleukin-3 Receptor Complex
Authors : Broughton, S.E.; Parker, M.W.
Deposited on : 2017-02-19
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	20% 87% • 10%
1	G	288	9% 86% 5% 8%
2	B	122	11% 83% • 14%
2	I	122	7% 78% 10% • 11%
3	C	3	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	4	 25%75%
5	E	4	 100%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-3 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2105	1331	378	381	15			
1	G	265	Total	C	N	O	S	0	0	0
			2143	1357	379	391	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	212	GLN	ASN	engineered mutation	UNP P26951
A	299	VAL	ALA	engineered mutation	UNP P26951
G	212	GLN	ASN	engineered mutation	UNP P26951
G	299	VAL	ALA	engineered mutation	UNP P26951


- Molecule 2 is a protein called Interleukin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	105	Total	C	N	O	S	0	0	0
			845	532	152	157	4			
2	I	108	Total	C	N	O	S	0	0	0
			871	551	155	161	4			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	13	TYR	TRP	engineered mutation	UNP P08700
I	13	TYR	TRP	engineered mutation	UNP P08700

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	G	1	Total	C	N	O	0	0
			14	8	1	5		
6	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	G	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	G	1	Total	C	O	0	0
			6	3	3		
7	G	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	33	Total	O	0	0
			33	33		
8	B	9	Total	O	0	0
			9	9		
8	G	45	Total	O	0	0
			45	45		
8	I	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-3 receptor subunit alpha




- Molecule 1: Interleukin-3 receptor subunit alpha



- Molecule 2: Interleukin-3



- Molecule 2: Interleukin-3



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%

MAG1
MAG2
BMA3

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  25%  75%

MAG1
MAG2
BMA3
FUC4

- Molecule 5: beta-D-mannopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2
BMA3
FUC4

4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.04Å 132.04Å 210.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	114.35 – 2.70 114.35 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (114.35-2.70) 99.9 (114.35-2.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.11 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.242 , 0.286 0.245 , 0.291	Depositor DCC
R_{free} test set	1535 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6261	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL, FUC, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2157	0.80	0/2919
1	G	0.56	0/2198	0.81	2/2979 (0.1%)
2	B	0.58	0/860	0.85	0/1164
2	I	0.58	0/888	0.87	1/1204 (0.1%)
All	All	0.57	0/6103	0.82	3/8266 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	281	PHE	N-CA-CB	6.34	119.30	110.17
1	G	213	MET	N-CA-C	5.22	117.32	109.23
2	I	108	ARG	CB-CA-C	-5.09	102.34	110.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2031	0	0
1	G	2143	0	2065	7	0
2	B	845	0	848	4	0
2	I	871	0	875	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	39	0	34	0	0
4	D	49	0	43	0	0
5	E	49	0	43	0	0
6	A	14	0	13	0	0
6	G	28	0	26	0	0
7	G	18	0	24	0	0
8	A	33	0	0	0	0
8	B	9	0	0	0	0
8	G	45	0	0	0	0
8	I	13	0	0	0	0
All	All	6261	0	6002	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:87:ASN:CA	1:G:88:PRO:N	2.43	0.82
2:B:29:GLN:HG3	2:I:29:GLN:HG3	1.73	0.70
2:I:30:PRO:HG3	2:I:104:TRP:HE1	1.60	0.67
2:I:30:PRO:O	2:I:108:ARG:NH1	2.31	0.64
2:I:29:GLN:O	2:I:31:PRO:HD3	1.99	0.63
2:B:29:GLN:HG3	2:I:29:GLN:CG	2.33	0.57
2:I:30:PRO:HG3	2:I:104:TRP:NE1	2.22	0.54
2:I:16:CYS:CB	2:I:84:CYS:HG	2.17	0.54
2:I:29:GLN:N	2:I:30:PRO:HD2	2.24	0.51
1:G:224:MET:HE3	1:G:270:VAL:HG21	1.92	0.51
2:I:30:PRO:HG2	2:I:108:ARG:NH1	2.26	0.51
1:G:275:ARG:HD3	1:G:280:GLU:OE1	2.13	0.49
1:G:224:MET:CE	1:G:270:VAL:HG21	2.44	0.47
2:B:29:GLN:HG3	2:I:29:GLN:CD	2.40	0.47
2:B:55:ARG:HB3	2:B:56:PRO:HD3	1.97	0.46
2:I:25:THR:O	2:I:29:GLN:HG2	2.18	0.44
2:I:55:ARG:HB3	2:I:56:PRO:HD3	2.01	0.42
1:G:224:MET:HE1	1:G:240:LEU:HD23	2.02	0.42
1:G:214:THR:HG22	1:G:215:ALA:N	2.35	0.41
1:G:149:TYR:CD2	1:G:173:LEU:HD11	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/288 (86%)	238 (96%)	10 (4%)	1 (0%)	30	54
1	G	257/288 (89%)	246 (96%)	11 (4%)	0	100	100
2	B	101/122 (83%)	101 (100%)	0	0	100	100
2	I	104/122 (85%)	102 (98%)	2 (2%)	0	100	100
All	All	711/820 (87%)	687 (97%)	23 (3%)	1 (0%)	48	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/257 (90%)	224 (97%)	7 (3%)	36	66
1	G	237/257 (92%)	233 (98%)	4 (2%)	53	79
2	B	95/110 (86%)	94 (99%)	1 (1%)	65	85
2	I	98/110 (89%)	95 (97%)	3 (3%)	35	65
All	All	661/734 (90%)	646 (98%)	15 (2%)	44	73

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	57	ASP
1	A	171	SER
1	A	187	ARG
1	A	200	VAL
1	A	217	CYS
1	A	293	CYS
2	B	15	ASN
1	G	200	VAL
1	G	217	CYS
1	G	281	PHE
1	G	293	CYS
2	I	34	LEU
2	I	69	GLN
2	I	94	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	147	GLN
1	A	153	HIS
1	A	204	GLN
2	B	15	ASN
1	G	233	ASN
2	I	18	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.43	0	17,19,21	1.14	1 (5%)
3	NAG	C	2	3	14,14,15	0.70	0	17,19,21	1.42	1 (5%)
3	BMA	C	3	3	11,11,12	0.57	0	15,15,17	1.09	2 (13%)
4	NAG	D	1	1,4	14,14,15	0.44	0	17,19,21	1.84	5 (29%)
4	NAG	D	2	4	14,14,15	0.54	0	17,19,21	1.35	2 (11%)
4	BMA	D	3	4	11,11,12	0.46	0	15,15,17	0.74	0
4	FUC	D	4	4	10,10,11	0.55	0	14,14,16	0.85	1 (7%)
5	NAG	E	1	1,5	14,14,15	0.46	0	17,19,21	1.20	2 (11%)
5	NAG	E	2	5	14,14,15	0.43	0	17,19,21	1.10	2 (11%)
5	BMA	E	3	5	11,11,12	0.30	0	15,15,17	0.78	1 (6%)
5	FUC	E	4	5	10,10,11	0.47	0	14,14,16	1.83	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
4	NAG	D	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	BMA	D	3	4	-	2/2/19/22	0/1/1/1
4	FUC	D	4	4	-	-	0/1/1/1
5	NAG	E	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	0/2/19/22	0/1/1/1
5	FUC	E	4	5	-	-	0/1/1/1

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C4-C3-C2	4.78	118.03	111.02
5	E	4	FUC	C1-C2-C3	4.35	115.98	109.64
5	E	4	FUC	C1-O5-C5	3.74	121.80	112.97
4	D	1	NAG	C1-O5-C5	3.48	116.84	112.19
4	D	2	NAG	C1-O5-C5	3.44	116.79	112.19
4	D	2	NAG	O5-C1-C2	-3.32	106.15	111.29
4	D	1	NAG	C2-N2-C7	3.02	126.94	122.90
5	E	3	BMA	C1-O5-C5	2.60	115.67	112.19
5	E	2	NAG	C3-C4-C5	2.58	114.90	110.23
4	D	1	NAG	C8-C7-N2	2.54	120.33	116.12
4	D	1	NAG	C6-C5-C4	2.53	119.24	113.02
3	C	3	BMA	O5-C5-C6	2.52	112.57	107.66
5	E	4	FUC	O5-C1-C2	2.45	116.62	110.79
4	D	1	NAG	O5-C1-C2	-2.44	107.52	111.29
5	E	2	NAG	O5-C1-C2	-2.34	107.68	111.29
5	E	1	NAG	C4-C3-C2	2.29	114.37	111.02
3	C	1	NAG	C3-C4-C5	-2.27	106.11	110.23
3	C	3	BMA	C1-O5-C5	2.23	115.18	112.19
5	E	1	NAG	C1-C2-N2	-2.17	107.02	110.43
4	D	4	FUC	C1-O5-C5	2.03	117.76	112.97

There are no chirality outliers.


All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
4	D	3	BMA	O5-C5-C6-O6
3	C	3	BMA	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
4	D	3	BMA	C4-C5-C6-O6
5	E	1	NAG	C4-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6
5	E	2	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
5	E	1	NAG	O5-C5-C6-O6
5	E	2	NAG	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
3	C	2	NAG	C3-C2-N2-C7


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	G	401	-	5,5,5	0.36	0	5,5,5	0.88	0
6	NAG	A	404	1	14,14,15	0.78	0	17,19,21	1.62	1 (5%)
6	NAG	G	409	1	14,14,15	0.42	0	17,19,21	1.11	2 (11%)
7	GOL	G	403	-	5,5,5	0.40	0	5,5,5	0.36	0
7	GOL	G	402	-	5,5,5	0.32	0	5,5,5	0.28	0
6	NAG	G	408	1	14,14,15	0.78	1 (7%)	17,19,21	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	G	401	-	-	0/4/4/4	-
6	NAG	A	404	1	-	2/6/23/26	0/1/1/1
6	NAG	G	409	1	-	2/6/23/26	0/1/1/1
7	GOL	G	403	-	-	2/4/4/4	-
7	GOL	G	402	-	-	3/4/4/4	-
6	NAG	G	408	1	-	3/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	408	NAG	C1-C2	2.46	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	404	NAG	C1-O5-C5	5.67	119.78	112.19
6	G	408	NAG	C2-N2-C7	2.42	126.14	122.90
6	G	408	NAG	C4-C3-C2	2.26	114.33	111.02
6	G	409	NAG	O5-C1-C2	-2.13	108.00	111.29
6	G	409	NAG	C1-O5-C5	2.09	114.99	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	G	403	GOL	C1-C2-C3-O3
7	G	403	GOL	O2-C2-C3-O3
6	G	409	NAG	O5-C5-C6-O6
6	A	404	NAG	O5-C5-C6-O6
6	G	408	NAG	O5-C5-C6-O6
6	A	404	NAG	C4-C5-C6-O6
6	G	409	NAG	C4-C5-C6-O6
6	G	408	NAG	C4-C5-C6-O6
7	G	402	GOL	C1-C2-C3-O3
7	G	402	GOL	O1-C1-C2-C3
6	G	408	NAG	C3-C2-N2-C7
7	G	402	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/288 (89%)	1.23	58 (22%) 2 2	20, 43, 84, 97	0
1	G	265/288 (92%)	0.70	25 (9%) 14 12	19, 35, 71, 86	0
2	B	105/122 (86%)	1.01	14 (13%) 7 6	25, 43, 71, 102	0
2	I	108/122 (88%)	0.84	9 (8%) 17 14	22, 40, 66, 103	0
All	All	737/820 (89%)	0.95	106 (14%) 6 5	19, 40, 79, 103	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	34	LEU	5.9
2	I	35	LEU	5.6
2	B	34	LEU	5.5
2	B	33	PRO	5.5
2	I	33	PRO	5.5
2	B	35	LEU	5.3
1	A	266	GLY	5.2
1	A	47	VAL	5.1
1	A	85	VAL	5.0
1	A	28	THR	4.8
1	A	220	THR	4.6
1	A	293	CYS	4.5
1	G	217	CYS	4.4
1	A	94	ILE	4.3
1	A	249	VAL	4.3
1	A	222	SER	4.2
1	A	221	HIS	4.2
1	A	223	PHE	4.1
1	A	42	ASP	4.0
2	I	31	PRO	4.0
1	A	245	ARG	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	29	GLN	3.9
1	A	265	PRO	3.9
2	B	30	PRO	3.8
1	A	41	TRP	3.7
1	A	45	ARG	3.7
1	A	90	PHE	3.6
1	A	83	VAL	3.6
2	I	36	ASP	3.6
1	A	244	LYS	3.5
1	G	90	PHE	3.5
2	B	29	GLN	3.5
1	A	267	THR	3.5
1	A	88	PRO	3.5
1	G	87	ASN	3.4
1	G	175	SER	3.3
1	G	47	VAL	3.3
1	G	27	ILE	3.3
1	A	91	SER	3.3
1	G	89	PRO	3.3
2	B	88	ALA	3.3
1	A	219	LYS	3.1
1	A	92	THR	3.0
1	A	39	LEU	3.0
1	A	263	LEU	3.0
2	I	37	PHE	3.0
1	A	68	CYS	3.0
1	A	89	PRO	3.0
2	B	36	ASP	3.0
1	A	240	LEU	2.9
1	G	84	ARG	2.9
1	A	217	CYS	2.9
2	B	37	PHE	2.9
1	G	147	GLN	2.9
1	G	43	LEU	2.9
1	A	248	PRO	2.8
1	A	147	GLN	2.8
1	A	93	TRP	2.8
1	G	25	PRO	2.8
1	A	212	GLN	2.8
1	G	290	ARG	2.8
1	A	260	PHE	2.7
1	A	250	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	28	THR	2.7
1	A	65	ASN	2.7
1	A	62	ALA	2.7
1	A	252	GLU	2.7
1	G	88	PRO	2.6
1	G	77	GLU	2.6
1	A	30	LEU	2.5
1	A	29	ASN	2.5
1	A	292	GLU	2.5
1	A	215	ALA	2.5
1	A	172	ARG	2.5
1	A	63	VAL	2.5
1	A	31	ARG	2.4
1	G	245	ARG	2.4
1	G	294	ASP	2.4
1	A	165	CYS	2.4
1	A	291	PHE	2.4
1	A	40	THR	2.4
1	A	163	ILE	2.4
2	B	89	THR	2.4
2	I	59	GLU	2.3
1	A	218	ASN	2.3
1	A	242	ILE	2.3
1	A	33	LYS	2.3
1	G	214	THR	2.3
1	G	149	TYR	2.3
2	I	121	ALA	2.2
2	B	90	ALA	2.1
2	B	92	PRO	2.1
1	A	247	GLN	2.1
1	G	176	GLY	2.1
2	B	59	GLU	2.1
1	G	247	GLN	2.1
1	A	57	ASP	2.1
1	A	146	ARG	2.1
1	G	26	PRO	2.1
1	A	141	ASN	2.1
2	B	39	ASN	2.1
1	G	57	ASP	2.0
1	G	256	ASP	2.0
1	A	261	GLN	2.0
2	B	69	GLN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	142	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	1	14/15	-	-	75,78,80,80	0
3	NAG	C	2	14/15	-	-	75,83,86,87	0
3	BMA	C	3	11/12	-	-	76,84,89,93	0
4	NAG	D	1	14/15	-	-	64,70,74,76	0
4	NAG	D	2	14/15	-	-	72,76,77,80	0
4	BMA	D	3	11/12	-	-	73,84,88,89	0
4	FUC	D	4	10/11	-	-	77,82,84,84	0
5	BMA	E	3	11/12	0.36	0.37	79,83,84,86	0
5	NAG	E	2	14/15	0.55	0.22	66,73,78,80	0
5	NAG	E	1	14/15	0.65	0.19	53,57,63,66	0
5	FUC	E	4	10/11	-	-	62,63,64,64	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	G	408	14/15	0.46	0.21	66,72,75,75	0
6	NAG	A	404	14/15	0.67	0.19	62,67,69,70	0
6	NAG	G	409	14/15	0.71	0.16	60,63,64,66	0
7	GOL	G	403	6/6	0.74	0.18	58,60,61,61	0
7	GOL	G	401	6/6	0.79	0.27	37,39,41,41	0
7	GOL	G	402	6/6	0.89	0.16	48,50,50,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.